



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



4 106 316 268

Bot. Gard
48
Z 5.9



HARVARD UNIVERSITY

LIBRARY

OF THE

GRAY HERBARIUM

Received 13 Feb. 1914

Bought
Bound 10 May, 1919

Gray Herbarium
Harvard University
13 Feb. 1914

CONSPECTUS
PLANTARUM IN HORTO BOTANICO
ET
PLANTATIONE
UNIVERSITATIS LUNDENSIS

PRÆCIPUE ANNIS 1834—1837 OBVIARUM,

QUEM
PROPOSUIT

an
JOH. WILH. ZETTERSTEDT,
HORT. BOT. LUND. H. T. PRÆFECT.

LUNDÆ,
TYPIS EXCUDIT CAROLUS FR. BERLING,
UNIVERSITATIS TYPOGRAPHUS.
MDCCCXXXVIII.

Bot. Gard

Z LIBRARY OF THE GRAY HERBARIUM

HARVARD UNIVERSITY.

BOUGHT.

Digitized by Google

1898

~~1894~~

W. 20

coll. compl. m. w.

Abbreviationes et Signa.

- Auctores citati:* Ag. *Agardh*; Ait. *Aiton*; All. *Allioni*; Anders. *Andersson*; Andr. *Andrews*; Andr. *Andrasiowski*; Ard. *Arduino*; Aubl. *Aubl.*
- Balb. *Balbis*; Bart. *Barton*; Bauh. *Bauhinus*; Benth. *Bentham*; Berg. *Bergius*; Bernh. *Bernhardi*; Bertol. *Bertolone*; Bess. *Besser*; Bivon. *Bivona*; Booth. *Booth et Söhne*, Hamb. 1837; Brot. *Brotero*; P. Br. *Patr. Browne*; R. Br. *Rob. Brown*; Buchan. *Buchanan*, deinde *Hamilton*; Burch. *Burchell*.
- Cand. de *Candolle*; Cassin. *Cassini*; Cav. *Cavanilles*; Cerv. *Cervantes*; Chais. *Cham. Chamisso*; Corr. *Correa*; Crantz; Curt. *Curtis*; Cuss. *Cusson*; Cyr. *Cyrillo*.
- Delil. *Delile*; Desf. *Desfontaines*; Desrouss. *Desrousseaux*; Desv. *Desvaux*; Dietr. *Dietrich*; Don; Donn; Dougl. *Douglas*; Dufresn. *Dufresne*; Dun. *Dunal*; d. Roi *du Roi*.
- Ehrh. *Ehrhart*; Estrch. *Estreicher*; Etling. *Etlinger*.
- Feuill. *Feuilles*; Fisch. *F. Fischer*; Forst. *Forster*; Fr. El. *Fries*; Friedl. *Friedland*.
- Glox. *Glosin*; Gmel. *Gmelin*; Gochn. *Gochwat*; Gooden. *Goodenough*; Gouan; Grah. *Graham*; Guers. *Guersent*; Guss. *Gussone*; Gärtn. *Gärtner*.
- Haw. *Haworth*; Henck. *Henckel*; Herit. *l'Heritier*; Herm. *Hermann*; Hilar. *Aug. S. Hilaire*; Höck. *Höckert*; Hoffm. *Hoffmann*; Hopp. *Hoppe*; Hornem. *Hornemann* (Hort. Hafn. 1813); Hornsch. *Hornschuch*; Host; Huds. *Hudson*; Humb. *Humboldt*.
- Jacqv. a *Jacquin*; Juss. *Jussieu*.
- Kämpf. *Kämpfer*; Ker. *Gawler, dein Kery*; Kit. *Kitabel*; Koch; Kön. *König*; Kunth; Kunz. *Kunze*.
- Labill. *Labillardière*; Lag. *Lagasca*; Lam. *Lamarck*; Lapeyr. *Lapeyrouse*; Ledb. *Ledebour*; Leers; Lehm. *Lehmann*; Lepech. *Lepechin*; Liljebl. *Liljebld*; Lindl. *Lindley*; Linck; L. a *Linne*; L. fil. a *Linne, filius*; Lois. *Loiseleur*; Lour. *Loureiro*; Lumnitz. *Lumnitzer*.
- M. B. *Marschall a Bieberstein*; Mart. *Martius*; Mass. *Masson*; Med. *Medicus*; Mey. *Meyer*; Mx. *Michaux*; Mill. *Miller*; Mönch; Mörch. *O. J. N.*; Molin. *Molina*; Morett. *Moretti*; Moris. *Morison*; Mühl. *Mühlenberg*; Murr. *Murray*.
- Nees. *Nees ab Esenbeck*; Nocc. *Nocca*; Nolt. *Nolte*; Nutt. *Nuttall*.
- Oed. ab *Oeder*; Ort. *Ortega*; Outh; Ott. *Otto*.
- P. B. *Patient de Beauvois*; Pall. *Pallas*; Pers. *Persoon*; Pohl; Poir. *Poiret*; Poll. *Pollich*; Pollin. *Pollini*; Pourr. *Pourret*; Presl; Pursh.

Rafin. *Rafinesque*; Ram. *Ramond*; Red. *Redouté*; Reich. *Reichardt*; Reichb. *Reichenbach*; Retz. *A. J. Retzius*; Rich. *Richard*; R. et Sch. *Römer et Schultes*; Roch. *Rochel*; Rosc. *Roscoe*; Roth; Roxb. *Roxburgh*; R. et P. *Ruiz et Pavon*.

Salisb. *Salisbury*; Salm; Sav. *Savi*; Schousb. *Schousboe*; Schröd. *Schrader*; Schrb. *Schreber*; Schweigg. *Schweigger*; Scop. *Scopoli*; Sibth. *Sibthorp*; Sieb. *Sieber*; Sims; Sm. *Smith*; Sol. *Solander*; Spr. *Curt. Sprengel*; Steph. *Stephan*; Stern. a *Sternberg*; Stev. *Steven*; Sw. *Swartz*; Sweet.

Tausch; Tenor. *Tenore*; Thuill. *Thuillier*; Thb. *Thunberg*; Torr. *Torrey*; Tournef. *Tournefort*; Trev. *Treviranus*; Trin. *Trinius*.

Vahl; Vent. *Ventenat*; Vig. *Viguier*; Vill. *Villars*.

Wahl. *Wahlenberg*; Wall. *Wallich*; Wallr. *Wallroth*; Walt. *Walter*; Weib. *Weike*; Wendl. *Wendland*; Wib. *Wibel*; Wilbr. *Wilbrand* (1837); W. *Willdenow*; Wither. *Withering*; Wulf. *Wulfen*.

Zuccar. *Zuccarini*.

Horti Botanici, e quibus semina aut plantæ communicata fuerunt;

Berolinensis, *Bonnensis*, *Chiswicensis* (Angliæ), *Cracoviensis*, *Edinburgensis*, *Gryphicus*, *Hafniensis*, *Hamburgensis*, *Holmiensis*, *Kewensis*, *Kiliensis*, *Lipsiensis*, *Londinensis*, *Parisiensis* et *Petropolitæ*.

☉.	signific. plant. annuam (ett-årig).	C. —	Caldarium (Varmhus; Drifthus hvari medeltemperaturen är 10–18 graders varme).
♂.	— — biennem (tuårig).	D. —	Sub Dio l. Diu (under bar himmel; på fritt land).
♀.	— — perennem (flerårig).	F. —	Frigidarium (Kallhus, Kylhus: 1–5 graders varme under vintertiden).
♂.	— suffruticem, fruticem aut arborem (busk- vext, buske eller träd).	T. —	Tepidar. (Ljumbus: 5–10 grad. varme under vintertiden).

Acacia

- acanthocarpa. W. \bar{p} . T.
alata. R. Br. \bar{p} . T. 1 ulnam
alta.
decipiens. R. Br. \bar{p} . T.
decurrens W. \bar{p} . T. Intr. 1837.
ex Anglia.
floribunda W. (*Mimosa* Vent.)
 \bar{p} . T.
linifolia. W. \bar{p} . T.
lophantha W. (*Mimosa dista-*
chya Vent.) \bar{p} . T.
prostrata \bar{p} . T. $1\frac{1}{2}$ uln. alt.
Introd. ex horto Hamburg.
1833.
undulata W. \bar{p} . T. 12 pollic.
alt. Intr. ex hort. Hafniæ
1833.
venusta W. \bar{p} . T. Ex. Ang-
lia intr. 1837, curante
Clar. J. Agardh.
viscosa Wendl. \bar{p} . T.
species \bar{p} . T.

Acæna

- adscendens Vahl. \bar{q} . F. D. Intr.
1837 ex h. Hafniæ.
ovalifolia R. et Pav. \bar{q} . F.
intr. 1837 ex h. Hafniæ.

Acalypha

- caroliniana. Walt. (*pauciflora*
Hornem.) \odot C. intr. 1837
ex h. Parisino, Ag.
rubra W. \bar{q} . C. intr. 1837 ex
h. Parisino.

Acer (Lönn).

- campestre L. \bar{p} . D.
creticum L. \bar{p} . F. intr. 1837.
ex h. Paris.
dasycarpum Ehrh. \bar{p} . D. in
nov. nostr. arboretum in-
trod. 1836, hodie 20 uln.
alt.
monspessulanum L. \bar{p} . D. 10
12 uln. alt.
Negundo L. \bar{p} . D. in planta-
tione Paradislyckan 22
uln. alt.
obtusifolium Sm. \bar{p} . D. intr.
1837 ex h. Hafn.
Opalus Ait. olim in horto no-
stro bene viguit; ibi diu-
tius non exstat.
pensylvanicum L. (*striatum* du
Roi.) \bar{p} . D. in arbor, ubi
quotannis floret; 5 uln. alt.
Platanoides L. \bar{p} . D.
 β . laciniatum du Roi. in
arb. 8. uln. alt.
Pseudoplatanus L. \bar{p} . D.
saccharinum L. \bar{p} . D. in arb.
6 uln. alt.
spicatum Lam. (*montanum* Ait.)
 \bar{p} . D. 4 uln. alt.
tataricum L. \bar{p} . D. 8. uln. alt.

Achania

- Malvaviscus Sv. (*Hibiscus* L.)
 \bar{p} . T.

Achillea

- albicaulis \bar{q} . D. ex h. Petrop.
1836.
alpina L. \bar{q} D.
asplenifolia Vent. \bar{q} . D. ex
Ht. Gryph, 1836.

bahnatica W. 2. D. ex. h. Gr.
 1837.
compacta W. 2. D. ex. horto
 Gr. 1837.
coronopifolia W. 2. ex. h.
 Gr. 1837.
decolorans Schrad. 2. ex. h.
 Gr. 1837.
filipendula Lam. (*A. Eupato-*
rium et filicifolia M. B.)
 2. D. ex. h. h. Petr. et
 Paris. 1837.
impatiens L. 2. D.
leptophylla M. B. 2. D. 1819.
magna L. 2. D.
Millefolium L. 2. D. spont.
nobilis L. 2. D.
pauciflora Lam. 2. D. ex. h.
 Paris. 1836. *Ag.*
Ptarmica L. 2. D.
 β. fl. pleno.
prostrata 2. D. ex. horto Gr.
 1837.
salicifolia Host. 2. ex. h. Gr.
 1837.
Santolina L. Spr. 2. T. ex. h.
 Petr. 1836.
serrata Retz. 2. D. 1826.
speciosa Henckel. 2. D. ex. h.
 Gr. 1837.
taurica M. B. 2. ex. h. Gr.
 1837.
vermicularis 2. F. ex. h. Petr.
 1836. sub hoc nom.

Achyranthes

brachiata L. (*Illecebrum* W.)
 ☉. T. ex. h. Hafn. 1837.
lappacea L. vide *Desmochata*
atropurpurea.
spinosa Hornem v. *Illecebrum*.

Achyronia

arborea Spr. v. *Carmichaelia*
australis.

Achyrophorus

hispidus Horn v. *Hypochaeris*.

Aconitum (Stormhatt).

Cammarum L. Fries Fl. Scan.
 p. 25. (*A. Störkeanum* Rei-
 chenb.) 2. D. (Obs. *Adon.*
Napellus Linn. genuinum,
 paullo humilior, florib. mi-
 nus pure caeruleis, capsu-
 lis divergentibus, in hor-
 to nostro non colitur, sed
 in nemore ad Hasslemölla
 Scan. etc. spontan. mens.
 Aug. legitur).
cernuum Wulf. 2. D. ex hort.
 Cracoviensi 1837, in no-
 stro adhuc non floruit.
excelsum 2. T. sub hoc nom.
 ex h. Petrop. 1836, in no-
 stro adhuc non floruit.
galectonum Reichb. 2. D. ex
 h. Kiliensi 1837.
ochroleucum W. 2. D. 1825
 1826.
pyrenaicum Lam. 2. D. ex h. C.
 1837, in nostro non flor.
rostratum Cand. 2. D. ex h.
 Crac. 1837.
tauricum Wulf. 2. D. ex. h.
 Crac. 1837.
vulgare 2. D. ex h. Crac. 1837.

Acorus (Kalmus).

Calamus L. 2. D.

Achrotriche

ovatifolia R. Br. v. *Styphelia*.

Actæa

spicata L. 2. D.

Adelia*ligustrina* Mx. v. *Berga*.**Adiantum***Capillus Veneris* L. 2. F. 1814.**Adonis***æstivalis* L. (*miniata* Jacqv.)

⊙. D.

autumnalis L. ⊙. ♂. D.*vernalis* L. 2. D.**Adoxa***moschatellina* L. 2. D. *spont.***Aegilops***caudata* L. (*cylindrica* Host.)

⊙. ♂. D.

ovata L. ⊙. D.*squarrosa* L. ⊙. D. ex horto

Hafn. 1837.

triuncialis L. (*triaristata* W.)

⊙. D. intr. 1835.

Aegopodium*Podagraria* L. (*Sison* Spr.) 2. D
*sponte.***Aesculus** (Kastaniefräd).*flava* Ait. ♂. D. intr. 1827,
*hodie 4 uln. alt.**Hippocastanum* L. ♂. D.*Pavia* L. (*Pavia rubra* Lam.)

♂. D.

rubicunda Cand. ♂. D.**Aethionema***cappadocicum* Spr. (*Aeth. Byz-*
baunii Cand. *heris arabi-**ca* L.) 2. F. ex h. Grac.
1837.*saxatile* R. Br. (*Thlaspi* L.) ⊙.

♂. (2. Wilbr.) F. 1819.

Aethusa*Cynapium* L. ⊙. ♂. D. in h.
*vitium.*β. *elata* Hoffm. Fisch. var.
major ex h. Paris. 1836.*Meum* Ait. v. *Meum athaman-*
*ticum.***Agapanthus***umbellatus* Herit. (*Crinum afri-*
canum L.) 2. T.**Agave***americana* L. 2. T.**Ageratum***coeruleum* Desf. ⊙. C. ex h.
Paris. 1837.*conyzoides* L. ⊙. C. 1822.**Agrimonia***Eupatoria* L. 2. D.*odorata* Mill. 2. D. 1826.*pilosa* 2. D. ex h. Hamburg.
1835.*repens* L. 2. D.*svaveolens* Pursh. 2. D.**Agrostemma***Coeli rosa* L. (*Lychnis* Spr.)

⊙. D.

coronaria L. (*Lychnis* Spr.)

♂. D.

β. *bicolor.*

γ. fl. albo.

Githago L. (*Lychnis* Scop, Spr.)

⊙. D.

Agrostis (Hven).*alba* L. 2. D.*Cinna* Lam. v. *Cinna arundi-*
*naea.**mexicana* L. 2. D.*paniculata* W. v. *Polypogon mon-*
*speliensis.**procumbens* R. et Sch. 2. D.
ex h. Hafn. 1837.*retrofracta* W. (*amula* R. Br.)

2. T. ex h. Hafn. 1837.

- rubra* L. 2. D.
stolonifera L. 2. D.
tenacissima Hornem. 2. C. ex
 h. Hafn. 1837.
varia 2. D. ex h. Hafn. 1837.
verrucosa 2. D. ex h. Crac.
 1837.
versicolor Stev. 2. F. ex h.
 Hafn. 1837.

Ailanthus

- glandulosa* Desf. (*Rhus Caco-*
dendron Ehrh.) 2. D. intr.
 1832. hodie 3 $\frac{1}{4}$ uln. alt.

Aira (Tatel).

- cristata* L. (*Poa* Host.) 2. D.
 ex h. Hafn. 1837.
junceae Vill. 2. F. ex h. Paris.
 1837.
pulchella W. 2. F. ex h. Hafn.
 1837.

Ajuga

- genevensis* L. 2. D.
reptans L. 2. D. 1826.

Aizoon

- hispanicum* L. ☉. T.

Alcea L. v. *Althaea*.

Alchemilla (Dagškápa).

- alpina* L. 2. D.
Aphanes Leers. (*Aphanes ar-*
vensis L. *Alchemilla arven-*
sis Scop.) ☉. D.
pubescens M. B. Lam. 2. F.
 ex h. Hafn. 1837.
vulgaris L. 2. D.
 2. *montana* W.

Allionia

- incarnata* L. ☉. T. ex horto
 Hafn. 1835.
nyctaginea Mx. 2. D.

Allium (Lökö).

- albidum* Fisch. 2. D.
atropurpureum Kit. 2. D.
atroviolaceum 2. D. ex hort.
 Hafn. 1837.
baicalense W. (*senescens* L?)
 2. D. ex h. Paris. 1837.
Cepa L. ♂. D.
descendens L. 2. D.
Fischeri 2. F. ex h. Hafn. 1837.
fistulosum L. 2. D.
fragrans Vent. 2. F. intr. 1835.
lineare L. 2. D.
Moly L. 2. D.
nigrum W. 2. D.
nutans L. 2. D.
obliquum L. 2. D.
odorum L. 2. D. 1826.
oleraceum L. 2. D.
pallens Gouan. (*paniculatum*
 Spr?) 2. D. 1822.
Porrum L. ♂. 2. D.
rubens Hornem. 2. D. 1826.
Schoenoprasum L. 2. D.
spirale W. 2. D. ex h. Hafn.
 1837.
ursinum L. 2. D.
victoriale L. 2. D. 1808.

Alnus (Ahl).

- cordifolia* Tenor. 2. D. in novo
 arboret. 5 uln. alt.
incana 2. pinnatifida Wahl.
 2. D. Paradistlyckan.
plicata Hort. Hamburg. 2. D.
 in arbor. 9 uln. alt.
serrulata W. 2. D. in arbor.
 6 uln. alt.

Aloe

- arborescens* Cand. 2. T.

attenuata Haw. 2. T.
cymbæfolia Hornem. 2. T.
humilis Thunb. 2. T.
Lingva W. 2. T.
 β. angustifolia Hornem.
picta Cand. 2. T.
prolifera Haw. (*brevifolia* W.)
 2. T.
pulchra Jacqu. (*obliqua* Cand.)
 2. T.
retusa L. 2. T.
spiralis L. 2. T.
variegata L. 2. T.
verrucosa L. (*carinata* Cand.)
 2. T.
viscosa L. (*Havorthia concinna*
 Hav.) 2. T.

Alopecurus

nigricans Hornem. 2. D. 1812,
 iterum ex h. Hafn. 1837.
pratensis L. 2. D. spont.
utriculatus Pers. (*Phalaris* L.)
 ⊙. D. ex h. Paris. 1837.

Alpinia

nutans Rosc. Spr. (*Globba* W.
 Renealmia Andr.) 2. C.

Alsine

globulosa T. Sub hoc nomin.
 ex h. Petropol. 1836.
media L. v. *Stellaria*.

Alstroemeria

pittacina Lehm. 2. T. ex h.
 Kiliens. 1836.

Althæa

ficifolia Cav. (*Alcea* L.) ♂. 2.
 D. ex Hafn. 1836.

Froloviana ⊙. ♂. D. ex hort.
 Hafnæ 1835.
narbonensis Jacqu. 2. D.
officinalis L. 2. D.
pallida Kit. ♂. D. 1812.
plebeja p. F. 2 uln. alt. 1835
 1837.
rosa Cav. (*Alcea* L.) ⊙. ♂.
 2. D.
 β. fl. plen. et color. var.
 species.

Alyssum

argenteum W. p. F. ex hort.
 Paris. 1837. cur. J. Ag.
calycinum L. ⊙. D. ex hort.
 Crac. 1837.
caucasicum 2. D.
cheirifolium Hort. Cracov.
 D. 1837.
creticum L. v. *Vesicaria*.
dentatum H. Cracov. D. 1837.
Draba W. ♂. D. an var. *A.*
 hirsuti?
edentulum Kit. ♂. D. 1819.
hirsutum M. B. ♂. ⊙. ex h.
 Crac. 1836.
incanum L. (*Farselia* R. Br.)
 ♂. D. etiam ex h. Crac.
 1837.
maritimum Lam. v. *Clypeola*.
montanum L. 2. F. D. 1808.
sativum Scop. v. *Camelina*.
tortuosum Kit. 2. D.
saxatile L. p. F. D. ex hort.
 Crac. 1837.
sinuatum L. v. *Vesicaria*.
spinosum L. (*Draba* Lam.) p.
 F. ex h. Paris. 1837.
utriculatum L. v. *Vesicaria*.
vernale Hornem. ♂. D. 1819.
vesicarioides 2. D. Sub hoc
 nom. ex h. Crac. 1837.

Alyxia

daphnoides p. T. intr. 1837.
 sub hoc nom. ex h. Kew.
 communicat. per Cl. J. Ag.

Amaranthus

- albus L. (C). D.
 aureus Dietr. (C). D. ex hort.
 Crac. 1837, et Kil. 1836.
 hæticus? H. Crac.; commun.
 1837. (*hecticus* videt.).
 Berchtholdi Hort. Crac.; com-
 mun. 1837.
 bicolor W. (C). C.
 Blitum L. (C). D.
 caudatus L. (C). D.
 cauliflorus Link. (*persicarioides*
 Poir.) (C). D. ex hort.
 Crac. 1837.
 deflexus L. (C). D. ex h. Crac.
 1837.
 flavus L. (*chlorostachys* W.) (C).
 D. ex h. Paris. 1837.
 gangeticus L. (C). C. ex hort.
 Crac. 1837.
 græcizans L. (C). D. ex h. Pa-
 ris. 1837.
 hybridus L. (*hecticus* W.) (C).
 D. semel introd. sponte dis-
 seminatur.
 hypochondriacus L. (C). D.
 inamoenus W. Hornem. (C). D.
 ex h. Crac. 1837.
 incompertus W. Hornem. (C). D.
 ex h. Crac. 1837.
 lætus W. (C). D.
 maximus Mill. (C). D. ex h.
 Crac. 1837.
 paniculatus L. (C). D. ex hort.
 Crac. 1837.
 polygamus L. (C). C. 1822.
 polygonoides L. (C). C.
 recurvatus (C). D. ex h. Paris.
 1837.
 retroflexus L. (C). D. ex hort.
 Crac. 1837.
 rigidus Hort. Hafn. ex h. Pa-
 ris. 1837, cura J. Ag.
 sanguineus L. (C). D.
 scandens L. (C). P. C. 1820
 1822.
 speciosus Ker. (C). C. ex hort.
 Kil. 1836, Crac. 1837.
 spinosus L. (C). C.
 tenuifolius W. (C). C. ex h.
 Paris. 1837.

tricolor L. (C). C.
 virens Hort. Crac. (C).

Amaryllis

- aurea Ait. 2. C. T.
 Belladonna L. (*variabilis* Jacqv.)
 2. C.
 chloroleuca Ker. 2. T.
 coranica Burch. 2. C.
 crocata Ker. 2. C.
 curvifolia Jacqv. 2. T.
 cylindracea Herit, v. *Cyrtanthus*
angustifolius.
 equestris Jacqv. (*dubia* L.) 2. F.
 formosissima L. 2. C.
 gigantea Ait. (*Crinum* Andr.)
 2. C.
 Johnsonii Hort. Hamb. 1836.
 2. C.
 lutea L. v. *Sternbergia*.
 minuta Humb. Spr. (*Zephyran-
 thes grandiflora* Spr.) 2. T.
 psittacina Ker. 2. C.
 pulverulenta Hort. Hamb. 2.
 C. 1836.
 purpurea Ait. (*Crinum specio-
 sum* Th. L.) 2. C.
 Radula Jacqv. (*Brunsvigia* H.
 Hamb.) 2. T. C.
 Regina L. (*brasiliensis* Andr.)
 2. C.
 revoluta Ait. (*Crinum lineare*
 Thunb.) 2. C.
 undulata Jacqv. 2. C. T.
 vittata Ait. 2. C.

Ambrosia

- artemisiæfolia L. (C). D. 1808
 1812; iterum ex h. Paris.
 1837.
 maritima L. (C). D. ex h. Pa-
 ris. 1837 cur J. Ag.
 trifida L. (C). D. ex h. Paris.
 1837.

Amellus

pedunculatus Ort. v. *Balbisia elongata*.

Amethystea

coerulea L. ☉. D.

Ammannia

egyptiaca W. (*viridis* Hornem.) ☉. C. 1819.

baccifera L. (*indica* Lam.) ☉. C. ex h. Hafn. 1836.

caspica M. B. ☉. C. ex h. Hafn. 1836. an var. *egyptiaca*?

debilis Ait. ☉. C. ex h. Hafn. 1836.

diffusa W. ☉. C. ex h. Hafn. 1836.

latifolia L. ☉. C. ex h. Kil. 1836.

multiflora Roxb. ☉. C. ex h. Hafn. 1836.

sangvinolenta Sw. ☉. C. ex h. Hafn. 1836.

senegalensis Lam. ☉. C. ex h. Hafn. 1836.

Ammi

majus L. (*A. Böberi* Höckert) ☉. D.

Ammobium

alatum R. Br. v. *Isodia Ammobium*.

Amomum

Zingiber L. v. *Zingiber officinale*.

Amorpha

fruticosa L. ♀. D. Ante 40 annos in horto Lundensi sine successu sub Dio plantata (cfr. Act. Holm. 1798); ibi vero hodie bene vigens, 2 uln. alt.

Lewinii Hort. Hamb. ♀. D. in nov. arb. intr. 1834.

species.

Ampelopsis (Vitis)

hirsuta ♀. D. sub hoc nom. ex h. Hamb. 1834 communicat.

hederacea Mx. v. *Vitis*.

Amsinkia

lycopsoides Lehm. 2? F. ex h.

Paris. 1837. cur. Dom. J. Ag.

Amsonia

salicifolia Pursh. 2. T. D. semina ex h. Kiliensi a cel. Profes. Nolte mihi missa 1836.

Amygdalus (Mandelträd).

communis L. ♀. D. in arbor. intr. 1827; hodie 8 uln. alt.

nana L. ♀. D.

pedunculata ♀. D. in nov. arb. intr. 1834 ex h. Petrop.; jam $\frac{1}{2}$ uln. alt.

persica L. ♀. D.

pumila L. D. ♀. in nov. arbor.

Anacyclus

aureus L. ☉. 2. D. 1819.

bicolor ☉. D. 1836.

creticus L. ☉. D. 1819.

valentinus L. ☉. D.

Anagallis

arvensis L. ☉. D.

caerulea Schreb. ☉. D. ex h. Hafn. et Cracov. 1837.

latifolia L. ☉. D.

Anchusa

altissima 2? D. ex h. Paris. 1837.

angustifolia L. 2. D. ex hort. Paris. 1837.

arvensis Spr. et Fr. v. *Lycopsis*.

capensis Thb. 2. ex hh. Hafn. et Paris. 1837.

hybrida Tenor. ex h. Hafn. 1837.

latifolia 2. D. sub hoc nom.
ex h. Hafn. 1837.

lutea M. B. v. *Lycopsis*.
ochroleuca M. B. 2. D. ex h.
Paris. 1837.

officinalis L. 2. D.
paniculata Ait. (*italica* Retz.)

♂. 2. D.

procera Bess. ♂. D.

sempervirens L. 2. D.

tenella Hornem. ☉. T. ex h.
Hafn. 1837.

undulata L. 2. D. ex h. Hafn.
1837.

Ancistrum

repens 2. F. ex h. Paris. 1837.
sub h. nomine.

Andersonia

Sprengelioides R. Br. ♂. T. flo-
ruit d. 9 Jun. 1835., $\frac{1}{2}$
uln. alt.

Andromeda

calycantha ♂. F. D. sub hoc
nom. ex h. Hafniæ com-
mun. 1837.

paniculata L. v. *Lyonia*.

Andropogon

Gryllus L. v. *Pollinia*.

Ischæmum Kunth (*virens* Spr.)
2. D. ex h. Paris. 1837.

Sorghum Brot. v. *Sorghum vul-*
gare.

Androsace

elongata L. ☉. D. in horto
jam sponte disseminat.

lactea L. 2. F. olim in horto
cultā, ibi diutius non ex-
stat.

lactiflora Fisch. (*A. alismoides*
Hornem.) ♂. ☉. F. D. ex
h. Hafn. 1837.

Androsæmum

officinale All. (*Hypericum An-*
droseum L.) ♂. D.

Andryala

candidissima 2? T. ex hort.
Paris. 1837.

Anemone (Sippa).

altaica Fisch. 2. F. ex hort.

Gryph. a D. Prof. *Horn-*
schuch nobis 1837 commun.

capensis ex h. Paris. 1837 sub
hoc nom. commun.

dichotoma L. 2. D. olim horti
incola, hodie ibi ignota.

hepatica L. 2. D. var. flor.
simplic., plen., alb. rubr.,
cærul.

narcissiflora L. 2. F. ex hort.
Petropol. 1836.

nemorosa L. 2. D.

pratensis et *Pulsatilla* v. *Pulsa-*
tilla.

ranunculoides L. 2. D.

sibirica L. 2. D. ex h. Crac.
1837.

stellata Lam. (*hortensis* L.) 2. D.
sylvestris L. (*ochotensis* Fisch.)

2. D.

Anethum

foeniculum L. (*Meum* Spr.) ♂.
2. D.

gravecolens L. (*Pastinaca An-*
ethum Spr.) ☉. D.

Angelica

atropurpurea L. 2. D. 1826.

Carvifolia Spr. v. *Selinum*.

Angelonia

salicornioides T. sub hoc nom.
ex h. Paris. 1837.

Anoda

Dilleniana Cav. v. *Sida*.

incarnata Kunth v. *Sida*.

Anthemis

arabica L. ☉. D. 10-14-poll.
licar. flor. flav. Jul.

artemisiifolia W. v. *Chrysanthemum indicum*.

arvensis L. ☉. D. sponte.

austriaca Jacq. ☉. D. ex h. Paris. 1837.

chamomilla W. 2. F. 1822.

clavata Desf. ☉. D. 1811.

Cota L. (*ulissima* Spr.) ☉. D. 1808. iterum ex h. Paris. 1837.

Cotula L. ☉. D. sponte.

rigescens W. (*caucasica* Hornem.) 2. D. 1812-22. et jam ex h. Paris. 1837.

secundiramea Bivon. ☉. D. ex h. Paris. 1837.

tinctoria L. 2. D.

Antherium

annuum L. v. *Bulbine*.

cirrhatum Forst. v. *Arthropodium*.

Liliago L. 2. D. bipedal. flor. niv. Jul. Aug.

paniculatum Andr. v. *Arthropodium*.

pendulum Hornem. W. v. *Arthropodium*.

ramosum L. 2. D.

Anthoxantum

(Vårbrodd).

odoratum L. 2. D.

ovatum Lag. ☉. D. ex h. Paris. 1837.

Anthriscus

vulgaris Pers. (*Scandix Anthriscus* L.) ☉. D.

fumarioides Spr. (*Scandix* Kit.) 2. D. ex h. Paris. 1837.

Anthyllis

Barba Jovis L. 2. T. 1820 et iterum ex h. Kil. 1836.

Hermannia L. 2. T. fere 3 pedal. flor. parv. flav. Maj. maritima Schweigg. 2? T. ex h. Bonn. 1836.

tetraphylla L. ☉. D.

vulneraria L. 2. D.

Antirrhinum (Cfr. etiam *Linaria*).

Majus L. 2. D. flor. color. var.

molle L. 2. T. ex h. Paris. 1837.

Orontium L. ☉. D.

Anychia

dichotoma Mx. (*Oxeria canadensis* L.) ☉. D.

Apargia

aspera Kit. 2. D. 21 poll. alt. fl. flav. Jun.

autumnalis Hoffm. (*Leontodon* L.) 2. D. spont.

caucasica M. B. 2. F. 1822, et ex h. Paris. 1837.

crispa W. 2. F. 1822.

hastilis Host. (*Leontodon* L.) 2. D. pedalis et ultra.

hirta Scop. v. *Thrinicia*.

hispida W. (*Leontodon* L. *Aparg. dubia* Hopp.) 2. D. 1812 1826.

incana Scop. (*Hieracium* L.) 2. D. 1824.

tuberosa W. v. *Thrinicia*.

Villarsii W. 2. D. sesquipedal. flor. flav. Jun.

Apium (Persilja).

graveolens L. 2. D.

Petroselinum L. 2. D. 2. *crispum*.

Apocynum

hypericifolium L. 2. D. ex h. Paris. 1837 cur J. Ag.

venetum L. 2. D. in h. s. Dio
plantat, quotannis floret,
2 nln. alt. flor. parv. sa-
voalb. inodor. Aug. ex-
plicat.

rosea Cand. 2. ♂. D. ex hort.
Crac. 1837.
stricta Huds. 2. ♂. D. 1805
in horto culta, hodie exsult.
Thaliana L. ☉. D.

Aquilegia (Ackleja).

alpina L. ♂. 2. D. bipedal.
floret Maj. — Aug.
alata Ledeb. 2. D. a Cel. Nol-
te ex h. Kil. 1836. nob.
commun.
viridiflora Pall. 2. D. 9-pol-
lic. floret Maj. Jul.
vulgaris L. 2. D. fl. col. var.
β. *speciosa* 1824.
γ. fl. plen.

Arabis

albida Stev. (*caucasica* W.
Cheiranth. mollis Horn.) 2.
D. 1818.
auriculata Lam. (*aspera* All.)
☉. 2. F. ex hort. Crac.
1837.
collina Tenor. 2. D. 1833
intr. ex hort. Berol. 7-8
poll. alt.
conferta 2. F. D. ex h. Crac.
1837.
crispata W. 2. D. 4-5 pol-
lic. alt.
Halleri L. ♂. D. ex h. Crac.
1837.
hirsuta Scop. v. *Turritis*.
incana Roth. ♂. D. ex hort.
Crac. 1837.
leptocarpa H. Crac. 2. D. a
Cel. *Estreicher* nob. com-
mun. sub hoc nom. 1837.
lucida L. v. *Turritis glabra*.
muralis Bertol. 2. D. ex h.
Paris. 1837.
Patriniana Cand. 2. D. ex h.
Paris. 1837.
pendula L. ☉. ♂. D. ex h.
Paris. 1837.
patens Stern. 2. D. ex hort.
Crac. 1837.

Aralia

racemosa L. 2. D. 26 pollic.
alt. flor. alb. Aug.
spinosa L. ♀. D.

Arbutus

Unedo L. ♀. F. etiam sub dio
non sine successu planta-
ta, fere 2 uln. alt. floret
Januar. Mart.

Arctium (Karrborre).

Bardana W. Fries. ♂. D.
Arctotis v. *Cryptostemma*.

Ardisia

crenulata Vent. 1828 ex hort.
Hafn. commun., hodie pro-
fuga.

Arenaria

balearica L. 2. F. ex h. Pa-
ris. 1837 cur. cl. J. Ag.
cæspitosa Ehrh. (*verna* L.) 2.
F. ex h. Crac. 1837.
fasciculata Gouan. ☉. D. ex
h. Paris. 1837.
graminifolia W. 2. D. ex h.
Crac. a Prof. *Estreicher*
communic. 1837.
montana L. 2. F. ex h. Paris.
1837.
nardiflora (*nardifolia*?) Led.;
T. ex h. Crac. 1837.
rostrata Kit. (*striata* Spr.?) 2.
F. ex h. Paris. 1837.
serpyllifolia L. ☉. D. spont.
triflora L. (*grandiflora* Spr.)
2. F. ex h. Paris. 1837.
trinervia L. ☉. ♂. D.

Argemone

grandiflora Sweet; a Cel. Nolte.
ex h. Kil. nobis commun.
1836.

intermedia ♂. D. ex h. Paris.
1837.

mexicana L. ♂. D. 1837.

ochroleuca Lindl. ♂. ex hort.
Kil. 1836.

Aristolochia

Clematitis L. ♀. D. in horto
nostro fere vitium, sed ob
singularem foecundationis
modum per *Ceratomyzom*
pennicornem libenter ser-
vata.

Armeniaca

valgaris v. *Prunus armeniaca*.

Armeria

alliacea W. (*Statice* Cav. Sm.)
♀. F. ex Hafn. 1836.

fasciculata W. ♀. 1836.

maritima W. (*Statice Armeria*
L.) ♀. D.

plantaginea W. ♀. F.

vulgaris W. ♀. D.

Arnebia

cornuta ex h. Petrop. 1836.
an *Lithospermum*?

Arnica

Arnopogon

picroides W. v. *Urospermum*.

Artemisia

Abrotanum L. (Åbrodd) ♀. D.

Absinthium L. ♀. D. (Malört).

annua L. ♂. ♂. D. intr. 1835.
5-pedal. floret Sept. Nov.

austriaca Jaqv. ♀. D.

campestris L. ♀. D.

chamæmelifolia Vill. ♀. F. ex
h. Petr. 1836.

fasciculata M. B. ♀. F. C.
1835.

leucanthemifolia H. Par. (*des-*
sertorum Spr.) ♀. D. ex h.
Paris. 1837.

maritima L. ♀. D. 1826.

Marschalliana Spr. (*inodora*
M. B.) ♀. F.

monogyna Kit. ♀. F. ex hort.
Paris. 1837.

pontica L. ♀. D. 1811.

procera W. (*paniculata* Lam.)

♀. F. ex h. Paris. 1837

quingufida Reitz. pugill. (*ar-*
gentea? Ait.) ♀. F. D. in
horto Bot. 1810 culta, ho-

die ibi ignota.

scoparia Kit. ♂. ♂. D. 1811.

et iterum 1835.

tanacetifolia L. ♀. D. 1826.

variabilis ex h. Paris. 1837.

violacea ex h. Paris. 1837.

vulgaris L. ♀. D. (Gräbo). spont.

Arthropodium

cirrhatum R. Br. (*Anthericum*
Forst.) ♀. T. ex h. Kew.
1837 cur. Cl. J. Ag.

paniculatum R. Br. (*Antheric.*
Andr.) ♀. T.

pendulum Spreng. (*Phalangium*
Red. *Anthericum* Hornem.
W.) ♀. T.

Arum

Colocasia L. ♀. C.

divaricatum L. ♀. C.

maculatum L. ♀. D. in Para-
distyckan in umbrosis co-
piose. Folia numquam ma-
culata ut in speciminibus a
Cel. Achario datis, in hor-
to Klosteroslunda Ostrog.
cultis.

Sagittifolium L. v. *Caladium*.

tenuifolium L. ♀. D. 1826.

Arundo (Rör).

colorata W. v. *Phalaris arundi-*
nacea β. *picta*.

Asarum

canadense L. ♀. D. flor. mens.
Maj.

europæum L. ♀. D. flor. Maj.

Asclepias

arborescens L. (*Gomphocarpus*
R. Br.) ♀. ♀. T.

carnosa L. v. *Hoya*.

curassavica L. ♀. C.

fuscata W. v. *Cynanchum*.

incarnata L. ♀. D. 1½ uln. alt.

nigra L. v. *Cynanchum*.

nivea L. 2. ♀. C. ex h. Bonn.
1836, 21 poll. alt.
pulchra Ehrh. 2. D. F. 3 uln.
alta.
syriaca L. 2. D. 3 uln. alt.
Vincetoxicum L. v. *Cynanchum*.

Asparagus (Sparris).

officinalis L. 2. D. spont.
verticillaris L. Bieb. 2. F. ex
h. Paris. 1837.

Asperugo

procumbens L. ☉. D. spont.

Asperula (Madra).

aristata W. (*longiflora* Kit.
Spr.) 2. D. ex h. Hafn.
1837.

hirta Ramond. 2. F. D.
lævigata L. 2. D. 1822.
odorata L. 2. D.
taurina L. 2. D. 1826.
trichodes 2. F. sub hoc nom.
ex h. Petr. 1836 commun.

Asphodelus

luteus L. 2. F. D. ex h. Pa-
ris. 1837.
tauricus M. B. 2. F. ex hort.
Hafn. 1836.

Aspidium

bulbiferum Sw. (*Polypodium* L.)
2. D.
cristatum Sw. (*Polypodium* L.)
2. D.
Filixfemina Sw. (*Polypod.* L.
Asplen. Bernh.) 2. D.
Filix mas Sw. (*Polypod.* L.)
2. D.
fragile Sw. (*Polypod.* L.) 2. D.
regium Sw. (*Polypod.* L.) 2. D.
1808.

Asplenium

Scolopendrium L. v. *Scolopen-
drium*.

Aster

acris L. 2. D. 2 uln. alt.

adulterinus W. 2. D. 2½ uln.
altus.
amelloides Bess. (*elegans* W.)
2. D. ex h. Gryph. 1837.
annuus L. v. *Erigeron hetero-
phyllus*.
argyrophyllus Labill. ♀. T. ex
h. Hafn. 1836.
bellidifolius 2. D. ex. hort.
Cracov. sub h. nom. 1837.
chinensis L. ☉. D. flores co-
lore et plenit. mire ludunt.
concinus W. 2. D. 1826.
cordifolius L. 2. D. 32 poll.
altus.
dracunculoides Lam. 2. D.
1826.
dumosus L. 2. D. 2 uln. alt.
eminus W. 2. D. 2 uln. alt.
fastigiatus 2. D. ex h. Crac.
1837.
firmus Nees. (*puniceus* Spr.) 2.
D. ex h. Crac. 1837.
floribundus W. (*Novi Belgii*
L.) 2. D. 2 uln. alt.
glabrus sub h. n. ex h. Crac.
a Cel. *Estreicher* 1837 nob.
commun.
grandiflorus L. 2. D. flor. in
h. mensib. Oct. Nov.
hyssopifolius L. 2. D. 1812.
1837. 2 uln. alt.
incisus Fisch. v. *Grindelia*.
juncus Ait. 2. D. 1826. ho-
die hort. profug.
lævis L. 2. D. 2 uln. alt.
leptophyllus Desf. 2. T.
mutabilis L. 2. D.
punctatus Kit. 2. D. 2 uln.
altus.
sibiricus L. 2. D. 1811. 1826.
spectabilis Ait. 2. D. 1½ uln.
altus.
tenellus L. ☉. T.
Tradescanti L. 2. D. 1820.
Tripolium L. ♂. 2. D.
umbellatus Ait. (*Amygdalinus*
Lam.) 2. D. 1826.
species.

Asterocephalus Spreng.

- (*Scabiosa* species L.).
africanus Spr. 2. T. ex hort.
 Paris. 1837.
ambiguus Tenor. ♂. 2. D.
amoenus Spr. 2. D. ex hort.
 Gr. 1836.
argenteus Spr. F. ex hort.
 Paris. 1836.
etropurpureus Spr. ♂. D.
bannaticus Spr. ♂. D. ex h.
 Paris. 1837.
caucasicus Spr. (*Scabiosa* M.
 B.) 2. D.
Columbaria Spr. 2. D.
glauca 2. D.
graminifolius Spr. 2. F.
holosericeus Spr. (*Sc. pyrenai-*
ca All.) 2. D.
maritimus Spr. ♂. F.
ochroleucus Spr. ♂. D. flor.
 pallide flav.
palæstinus Spr. ♂. D. ex h.
 Paris. 1837.
prolifer Spr. ♂. D. ex h. Pa-
 ris. 1837.
siculus Spr. ♂. D. ex h. Hafn.
 1835. 1-2-pedal.
stellatus Spr. ♂. D.

Astragalus (Klo-ärt).

- aduncus* W. 2. D. ex h. Pe-
 trop. 1836.
asper Jaqv. 2. D. 1822 et ex
 h. Kil. 1837.
aureus W. 2. ex h. Petrop.
 1836 sub hoc nom. miss.
bæticus L. ♂. D.
carolinianus L. 2. D. 1800.
Cicer L. 2. D. 1820-26.
compactus Lam. 2. ex h. Pe-
 trop. 1836.
contortuplicatus L. ♂. D. ex
 h. Paris. 1837.
cylindricus Cand. 2. D. ex
 h. Petrop. 1836.
depressus L. 2. D. ex h. Pa-
 ris. 1837.
galegiformis L. 2. D.
glycyphyllus L. 2. D.

hispidulus Cand. (*A. Bucorus*
 W.) ♂. D.

Juvenalis; T. ex h. Par. 1837.
leucophæus Sm. 2. D. 1819.

multicaulis H. Hamb. 2. D.
 ex h. Petr. 1836.

odoratus Lam. 2. D. ex hort.
 Petr. 1836.

persicus, sub hoc nom. ex h.
 Petr. commun. 1836.

podocarpus H. Hamb. 2. D. ex
 h. Petr. 1836.

ponticus Pall. 2. D. ex hort.
 Paris. 1837.

Pseudo-Cicer 2. D. ex hort.
 Paris. 1837.

refractus H. Hamb. 2. D. ex
 h. Petrop. 1836.

sesameus L. ♂. D. ex h. Pa-
 ris. 1837.

Stevonianus Spr. (*gracilis* St.)
 2. D. ex h. Petr. 1836.

strictus 2. D. 2 uln. alt. flor.
 albido-flav. 1835.

sulcatus L. 2. D. ex h. Pa-
 ris. 1837.

Tragacantha L. 2. T. 1/2 uln.
 altus.

virescens Ait. (*falcatus* Lam.
 Pall.) 2. D. 1826.

Astrantia

- caucasica* W. 2. D.
Epipactis L. v. *Dondia*.
major L. 2. D. 3-pedal. flo-
 ret Jun. — Sept.

Athamanta

- caucasica* 2. D. ex h. Paris.
 1837.
condensata L. (var. *A. Libanot.*
 Spr.) 2. D. ex h. Paris.
 1837.
macedonica Spr. v. *Bubon.*
meum L. v. *Meum athamanticum.*
sibirica L. (var. *Libanot.* Spr.)
 2. D. ex h. Paris. 1837.

- sicala L. 2. D. ex h. Paris. 1837.
 stricta H. Hamb. 2. D. 1836.
- Athanasia**
 annua L. (*Achillea inodora* L. sp.) ☉. T. 1805-1814.
- Atraphaxis**
 spinosa L. 2. F. D. intr. 1834 ex h. Petrop. cur. Cel. C. Agardh.
- Atriplex** (Molla).
 hortensis L. ☉. D.
 patula L. ☉. D. spont.
 portulacoides L. 2. T. F.
 sibirica L. ☉. D. 1806-1820.
- Atropa**
 Belladonna L. (Vargbär) 2. D.
 frutescens L. (*Physalis* Cand.) 2. T.
physaloides L. v. *Nicandra*.
- Aubrietia**
 deltoidea Cand. (*Alyssum* L. *Fursetia* R. Br.) 2. F. D. ex h. Paris. 1837.
- Aucuba**
 japonica L. 2. T. 2½ uln. alt. intr. 1820.
- Aulacospermum**
 species ex h. Petrop. 1836.
- Avena** (Hafre).
 elatior L. v. *Holcus avenaceus*.
 fatua L. ☉.
 flavescens L. 2. D. in Paradis-lyckan.
 nuda L. ☉.
 orientalis Schreb. ☉. D. 1812.
 pyramidalis W. ex. h. Crac. 1837.
 sativa L. ☉. D.
- Axyris**
 amaranthoides L. ☉. D. 1811. 1820 et nuper etiam ex h. Paris. communic.
- Ayenia**
 pusilla L. ☉. C. ex h. Hafn. 1836.
- Azalea**
 indica L. 2. F. D. ex hort. Hafn. 1837; nuper etiam nob. donat. a Generos. L. Bar. *Gyllenkrok*.
 pontica L. 2. T. 1½ uln. alt. quotannis floret.
 viscosa L. 2. F. D. 1½ uln. alt. floret quotannis.
- Azara**
 dentata R. Br. 2. T. ex Anglia intr. 1837 cur. Cl. Bot. Adj. J. Agardh.
- Baccharis**
 species.
- Bäckea**
 saxicola H. Hamb. 2. T. ex h. Kew. 1837 cur. J. Ag. virgata Ker. (*Leptospermum* Forst.) 2. T.
- Bæria**
 crysostoma F. et M. ☉. ex h. Kil. 1836, et Parisiens. 1837.
- Balbisia**
 elongata W. (*Tridax procumbens* L. *Amellus pedunculatus* Ort.) C. 1805 et jam ex h. Paris. 1837.
- Ballota**
 foetida Lam. Fr. (*alba* L.) 2. D.
 ruderalis Sw. Fr. (*nigra* L. sp. ed. 2.) 2. D.
- Balsamita**
 ageratoides ex h. Paris. 1837 s. h. nom. communic.
 vulgaris W. (*Tanacetum Balsamita* L.) 2. D.
- Banksia**
 ericæfolia L. fil. 2. T.
- Baptisia**
 australis R. Br. (*Sophora* L. *Podaliria* W.) 2. F. D.

Barbarea

- præcox R. Br. Fr. ♂. D.
vulgaris R. Br. (*Erysimum Bar-*
baræa L.) ♀. D.

Barosma

- serratifolia* W. v. *Diosma*.

Bartonia

- auræa; T. ex h. Paris. 1837
sub h. nom. communis.

Bartsia

- Odontites Huds. (*Euphrasia* L.)
⊙. D.

Basella

- cordifolia Lam. ⊙. C. 1821.
rubra L. ⊙. C. ex hort. Kil.
1836.

Bauhinia

- brachycarpa Hort. Hamb. ♀.
C. 1836.

Beaufortia

- decussata R. Br. ♀. C. 1833;
etiam 1837 ex h. Kew.
communis.

Beckmannia

- erucæformis Host. ♀. D. 1826;
quoque 1837 ex h. Hafn.
miss.

Begonia

- discolor R. Br. Horn. ♀. C.
sub nom. *B. picta*, quæ
alia est, etiam communi-
cata.

- hirtella Link. ♀. C. 6 pollic.
alta.

- nitida Ait. (*obliqua* Herit. mi-
nor Jaqv.) ♀. C.

- semperflorens H. Hamb. ♀. C.
12-18 poll. alt.

- spathulata W. ♀. C. ex hort.
Kil. 1836.

Belis

- jaculifolia Salisb. (*Pinus lancoso-*
lata L. *Abies* Lour.) ♀. T.

Bellis

- perennis L. ♀. D.
β. hortensis.
γ. fistulosa.

Berberis

- Aquifolium Parsh. (*Mahonia*
Nutt.) ♀. T. intr. 1837,
ex Angl. per Cl. J. Agardh
communis.

- asiatica Roxb. ♀. F. intr. 1837,
ex Anglia.

- canadensis Mill. ♀. F. D.
coriaria (*coriacea* Hilar.?) ♀.
F. sub hoc nom. ex Angl.
1837.

- repens Booth. Hamb. ♀. T. ex
Angl. 1837.

- sibirica Pall. ♀. D. 1819.

- vulgaris L. ♀. D. baccis semi-
ne foetis. Paradislyckan.

- β. baccis apyrenis.
species ♀. T. ex Anglia 1837.

Beta

- Cicla L. ♂. D.

- β. altissima.

- maritima L. ♂. D. ex h. Pa-
ris. 1837.

- trigyna Kit. ♀. D. 2 uln. alt.
flor. alb. Jun.

- vulgaris L. ♂. D.

Betonica

- grandiflora W. ♀. D.

- hirsuta L. ♀. D. 1826. etiam
ex h. Crac 1837.

- incana Ait. ♀. D. 1835. 1½
uln. alt.

- officinalis L. ♀. D. 1½ uln. alt.

- orientalis L. ♀. D. 1820;
etiam ex hortis Paris. et
Crac. 1837.

- stricta Ait. ♀. D. 26 pollic.
alt. floret Aug.

Betula (Björk).

- alba L. ♀. D.

- β. pendula 1826.

carpinifolia Ehrh. ♀. D. 1826;
sed hodie non horti incola.
cordata Retz. (non *Alnus cor-*
difolia Tenor.) ♀. D. 1806,
nec hodie.

davurica Pall. ♀. D. 10 uln.
alt.

excelsa Ait. ♀. D. olim in
horto culta, ibi diutius
non exstat.

nana L. ♀. D. 2 uln. alt.

nigra L. ♀. D. 27 uln. alt.
profert quotannis flores,
sed nunquam semina ma-
tura.

papyracea Ait. ♀. D. arbor ster-
ilis, 2½ uln. alt.

pumila L. ♀. D. 1½ uln. alt.

Bidens

bipinnata L. ♂. C. 1812.

heterophylla Ortega. ♀. T. 1806.
1813.

leucantha W. ♂. D. 1835.

multifida ♂. D. ex h. Paris.
1837.

nivea L. v. *Melanthera hastata*.

sambucifolia Cav. ♂. D. 1835.

Bigelovia

stricta Spreng. (*Spermacoce* L.)

♂. C. ex h. Kil. 1837, a
Cel. Prof. Nolte nob. com-
municata

Bignonia

capensis ♀. C. sub h. nom. ex
h. Kew. 1837. communic.
cfr. *Tecoma*.

Catalpa L. v. *Catalpa syringi-*
folia.

heterophylla W. (β. *Cherere*
Aubl.) ♀. C.

jasminifolia Kunth. ♀. T. ex
h. Kew. 1837.

radicans L. v. *Tecoma*.

Billardiera (Billardiera).

scandens Sm. Spr. ♀. T. 1 uln.
alt. intr. ex h. Hamburg.
1835.

Biscutella

apula L. ♂. D. ex h. h. Pa-
ris. Hafn. et Crac. 1837.

auriculata L. ♂. D. 1812 et
ex h. Crac. 1837.

Columnæ Tenor. ♂. D. 1821.
iterum ex h. Kil. 1836.

depressa W. ♂. D. ex h. Pa-
ris. 1837.

hispida Cand. ♂. D. 1819.

lyrata L. ♂. D. ex h. Hafn.
1836.

marginata Tenor. Hornem. ♂. ?
D. ex h. h. Hafn. et Kil.
1837.

raphanifolia Poir. ♂. D. ex
h. Hafn. 1836.

Biserrula

Pelccinus L. ♂. D. ex hort.
Paris. 1837.

Blechnum

spicant Fr. (*Osmunda* L. *Lo-*
maria Desv.) ♀. D.

Blitum

capitatum L. ♂. D.

chenopodioides Lam. ♂. D.

petiolare ♂. D. ex h. Hafn.
1837.

virgatum L. ♂. D.

Blumenbachia

insignis Schrad. v. *Loasa pal-*
mata.

Bocconia

cordata W. ♀. C. ex h. Paris.
1837.

Boebera

chrysanthemoides W. (*Dysso-*
dia glandulosa. Cav.) ♂. D.

chrysostoma ♂. ? C. ex hort.
Paris. 1837.

radiata (*Dyssodia*) T. 1837.

Boltonia

asteroides Herit. (*Matricaria* L.)
♀. D. 1826.

Bonnaya

brachiata Link (*Ruellia anti-*
poda L. *Gratiola serrata*
Roxb.) T. ex h. Hafnæ
1837.

Bonplandia

gemiflora Cav. v. *Caldasia he-*
terophylla.

Borago

laxiflora Hornem. 2. F. ex h.
Paris. 1837.

officinalis L. (C). D.

β. fl. albe.

Borkhausia Böhm. *Crepis*
Auct.

alpina Link. (C). D. ex hort.
Paris. 1837.

bellidifolia Caud. (C). D. ex
h. Paris. 1837.

Candollei Spr. (C). D. ex h.
Paris. 1837.

hispida Link. (C). D. 1821.

hyemalis Spr. (C). D. ex hort.
Paris. 1837.

leontodontoides Spreng. ♂. D.
1807.

rubra Link. (C). D.

Borya

ligustrina W. (*Adelia* Mx.)
♂. T.

Bosea

Yervamora L. ♂. T. F.

Bowlesia

tenera Spr. (C). C. T. ex hh.
Hafn. 1836, Paris. 1837.

Brassia

maculata R. Br. ♂. T. ex Angl.
1837. cur. J. Ag.

Brassica (Kål).

campestris L. (C). D.

Napus L. (Rapsat) ♂. D.

oleracea L. ♂. D. plures varie-

tates adsunt.

Rapa L. ♂. D.

Briza (*Darrgräs*).

maxima L. (C). D.

media L. 2. D.

virens L. (C). D. ex hh. Hafn.
et Crac. 1837.

Bromelia

Ananas L. 2. C.

Bromus (*Losta*).

arvensis L. (C). D.

commutatus Schrad. (C). ♂. D.
1812.

giganteus L. 2. D. in horto
olim cultus, hodie ex-

stinctus.

laxus Hornem. (*erectus* Huds.)
2. D. 1826.

mollis L. (C). ♂. D.

multiflorus Host. (*patulus* Koch.)
(C). D. 1826.

purgans L. 2. D.

secalinus L. (C). D.

sterilis L. (C). D.

Brotera

Contrayerva Spr. v. *Flaveria*.

Broussonetia

papyrifera Vent. (*Morus* L.)

♂. D. olim in horto arbor

4-5 uln. alt., florifera,

post hiemem duram ex-

stincta, iterata vice 1835

seminibus educata hodie

bene viget.

Brovallia

elongata Kunth. (*elatu* Hortul.)

(C). C. ex h. Bonn. 1836.

Linnæana Spr. (*elata* et *demis-*
sa L. teste Spr.) (C). C.

ex hortis Kil. et Bonn.

1836.

Brunia

superba Don. ♂. T.

Brunsvigia

Radula v. *Amaryllis*.

Bryonia

alba L. 2. D. 3-4 uln. flor. Aug.

dioica L. 2. D. 1826.

Bryophyllum

calycinum Salisb. v. *Verea pin-nata*.

Bubon

buchtormense Fisch. (*Athaman-ta rigida* Horn.) 2. D. ex h. Paris 1837,

macedonicum L. (*Athamanta* Spr.) ♂. F.

Büchnera

cuneifolia Retz. v. *Manulea*.

foetida Andr. v. *Manulea*.

Buddleia

globosa Lam. ♂. T. 2½ uln. alt. flor. coccin. Maj. Jun.

spicata R. et Pav. ♂. T. ex h. Paris. 1837.

Bulbine

aloides W. 2. T. 1 uln. alt. intr. 1833.

annua W. (*Anthericum* L.) ♂. D. ex h. Paris. 1837.

Bulbocodium

vernum L. 2. D.

Bunias

balearica L. v. *Succovia*.

Erucago L. ♂. D. 1818; ex h. Paris. 1837.

β. *aspera* Retz. 1808.

orientalis L. 2. D. in horto jam spontan.

Bunium

Bulbocastanum L. (*Sium* Spr.) 2. D.

Buphthalmum

aquaticum L. ♂. D. 1812; ex h. Paris. 1837.

capense L. v. *Oedera prolifera*. *cordifolium* Kit. 2. D. Flora 1836 p. 8.

helianthoides L. v. *Heliopsis laevis*. *salicifolium* L. (*grandiflorum* L.) 2. D. 1808–1812.

sericeum L. suppl. ♂. D. 1826.

silphioides 2. D. 2½ uln. alt.

spinosum L. ♂. F. 1812.

Bupleurum

angulosum L. 2. D. ex hort. Paris. 1837, per Cl. J. Ag.

fruticosum Hornem. ♂. F.

Gerardi Jacqv. ♂. D. ex h. Kil. 1836.

junceum L. Spr. ♂. D. ex h. Kil. 1836.

longifolium L. 2. D. ex hort. Paris. 1837.

Odontites L. ♂. D. ex h. Paris. 1837.

protractum Link. ♂. D. ex h. Paris. 1837.

rotundifolium L. ♂. D. ex h. Kil. 1836.

semicompositum L. ♂. D. ex h. Paris. 1837.

sibiricum Willd. ♂. D. ex h. Kil. 1836.

species.

Buxus (Buxbom).

halearica Lam. ♂. T.

sempervirens L. ♂. D.

β. fol. varieg.

γ. minor.

Cacalia

articulata Thb. (2.) ♂. T.

cuneifolia L. (2.) ♂. T.

hastata L. 2. D.

repens L. (2.) ♂. T.

sagittata Vahl. ♂. C. 1823.

sonchifolia L. ♂. C.

avaveolens L. 2. D. 3 uln. alt.

Cachrys

cruenta sub hoc nom. ex h. Petr. 1836.

maritima Spr. (*Crithmum* L.) ♂. T.

microcarpa M. B. F. ex hort. Paris. 1837.

Cactus

alatus Sw. ♂. C. intr. 1833.

cæspitosus H. Hamb. ♂. C.

coccineus H. Hamb. ♂. C.

coccionellifer L. ♂. C.

curassavicus L. ♂. C.

elegans Link. (*phylantoides* Cand.) ♂. C.

flagelliformis L. ♂. C.

- foetalis* Salzm. $\frac{1}{2}$ C.
grandiflorus L. $\frac{1}{2}$ C.
humilis Lag. (*foliosus* W.) $\frac{1}{2}$ C.
leucanthus H. Hamb. $\frac{1}{2}$ C.
mesembryanthemoides H. Hamburg. $\frac{1}{2}$ C. intr. 1833 ex h. Gryph.
microdasys Lehm. $\frac{1}{2}$ C.
moniliformis L. $\frac{1}{2}$ C. 3 uln. altus.
Opuntia L. $\frac{1}{2}$ C.
Ottomis H. Hamb. $\frac{1}{2}$ C. ex h. Hafn. 1834.
oxygonus H. Hamb. $\frac{1}{2}$ C.
paradoxus Hornem. $\frac{1}{2}$ C.
pendulus Sw. (*Rhipsalis Cuscuta* Gärtn.) $\frac{1}{2}$ C.
pentagonus L. $\frac{1}{2}$ C.
pusillus Cand. $\frac{1}{2}$ C.
Qvintero H. Hamb. $\frac{1}{2}$ C.
repandus L. $\frac{1}{2}$ C.
roseus H. Hamb. $\frac{1}{2}$ C.
salicornioides Haw. $\frac{1}{2}$ C.
serpentinus W. $\frac{1}{2}$ C.
speciosus W. $\frac{1}{2}$ C. 2 uln. alt.
squamosus H. Hamb. $\frac{1}{2}$ C.
strigosus H. Hamb. $\frac{1}{2}$ C.
tetragonus L. $\frac{1}{2}$ C.
triangularis L. $\frac{1}{2}$ C.
truncatus Link. $\frac{1}{2}$ C. 6-7 poll. altus.
species.

Cæsalpinia
cucullata H. Hamb. $\frac{1}{2}$ C.

Cakile
rugosa Cand. (*Myagrum* L.)
 ♂. D. 1808; et seminibus ex h. Paris. miss. educat. 1837.

Caladium
sagittifolium Vent. (*Arum* L.)
 $\frac{1}{2}$ C.

Calandrinia
compressa Kunth. ♂. C. ex h. Crac. 1837.
discolor Schrad. $\frac{1}{2}$. (an *speciosa*?) ex h. Paris. 1837.

grandiflora Lindl. ♂. $\frac{1}{2}$ C.
 ex h. Crac. 1837.
pilosculula Cand. ♂. ex h. Paris. 1837.
speciosa Lindl. ♂. ♂. C. ex h. Bonn. 1836. 6-10 poll. alta.

Calceolaria
eximia Booth $\frac{1}{2}$ C. T. de hort. Hamb. 1837. emta.
pendula Don. $\frac{1}{2}$ C. T. ex h. Kilens. a Cel. *Nothe nobis* commun. 1836.
picta $\frac{1}{2}$ C. T. ex h. Hamb. 1837.
pinnata L. ♂. C.
plantaginea Sm. $\frac{1}{2}$ T. C. ex h. Kil. 1836,
salicifolia B. et Pav. $\frac{1}{2}$ T. $\frac{1}{2}$ uln. alt. intr. 1833.
species.

Caldasia
heterophylla W. (*Bonplandia geminiflora* Cav.) $\frac{1}{2}$ C. ex h. Paris. 1837.

Calendula
 (Ringelblomma).
arvensis L. ♂. D.
hybrida L. ♂. D.
incana W. ♂. $\frac{1}{2}$? T. ex h. Crac. 1837.
maritima Guss.; ex h. Crac. 1837.
officinalis L. ♂. D.
pluvialis L. ♂. D.
sicula W. ♂. D. ex h. Crac. 1837.
speciosa; ♂. D. ex h. Crac. 1837.
stellata Cav. ♂. D.
suffruticosa Vahl. $\frac{1}{2}$ F. 1835.
 $\frac{1}{2}$ uln. alt.
tomentosa Desf. (an *incana* W. ut voluit Sprengel?) ♂.
 $\frac{1}{2}$? T. ex h. Crac. 1837.

Calla
aethiopia L. (*Zantedeschia* Spr.)
 $\frac{1}{2}$ T.

Callicarpa
macrophylla Vahl. $\frac{1}{2}$ T. ex Angl. 1837. J. Ag.

Callicoma

serratifolia R. Br. (*Hermesia Banksiaefolia* Spr.) ♀. T. ex h. Kew. 1837.

Calligonum

Pallasii Ait. (*Pallasia caspica* L.) ♀. F. ex h. Petrop. 1836.

polygonoides L. ♀. T. ex h. Petrop. 1836.

Calliopsis

bicolor Reichb. (*Coraopsis tinctoria* Nutt.) ♂. D.

tinctoria Cand. ♂. D.

♂. *atrosanguinea* ex h. Kil. 1836.

Callistachys

lanceolata Vent. ♀. C. T. ex h. Crac. 1837.

ovata Sims. ♀. C. T. ex hort. Kew. 1837 Dom. J. Ag.

Callistemon

pinifolius Hortul. v. *Metrosideros*.

rigidus R. Br. (*Metrosider. linearis* W.) ♀. T. ex h. Crac. 1837.

vimineus. ♀. F. sub hoc nom. ex Angl. 1837 miss.

Calocephalus

nitens R. Br. T. ex h. Petrop. 1836. 1 ulq. alt.

Calothamnus

clavata H. Hamb. ♀. T. ex h. Kew. 1837 sub hoc nom. miss.

Calycanthus

floridus L. ♀. F. D. frutex loco umbroso ventisque sub ducto plantat. quotannis flores producit, 1 uln. alt; intr. 1833.

Calythrix

glabra R. Br. ♀. T. ex h. Kew. 1837.

Calythymenia

paniculata. 2? T. ex h. Paris. 1837. forte *oxybaphi* generis.

Camelina

dentata Fr. Fl. sc. (*Myagrum* W.) ♂. D.

laxa ♂. D. ex h. Petr. 1836. sub hoc nom. miss.

paniculata v. *Nestia*.

sativa Crantz (*Myagrum* L. *Alysum* Scop.) ♂. D. 1822—1836.

Camellia

japonica L. ♀. T. ex h. Hamb. intr. 1835.

♂. *alba*.

γ. *anemoniflora*.

δ. *coccinea*.

ε. *longifolia* (floruit 22. Jan. 1838).

Campanula (Klocka).

americana L. (*acuminata* Mx.)

2. D. ex h. Paris. 1837.

cur. Clar. J. Agardh.

azurca Soland. 2. T. ex hort. Gryph. 1837, a Cel. Hornschuch nob. communicata.

betonicæfolia, 2. F. D. sub hoc nom. ex h. Crac. 1837.

bononiensis Spr. (*ruthenica* W.) 2. D. 1822—1826.

carpathica L. 2. D. ex hort. Paris. 1837.

collina M. B. 2. D. ex hort. Crac a Cel. Estreicher nobis commun. 1837.

crenata H. Hamb. 2. D. ex h. Hafn. 1837.

crenulata Fisch. 2. F. D. ex h. Gryph. 1837.

divergens W. (*spathulata* Kit.) ♂. 2. D. ex hh. Hafn. et Paris. 1837.

Erinus L. ♂. D. F. 1827.

falcata R. et Sch. (*Prismatocarpus* Tenor.) ♂. D. ex h. Paris. 1837.

glomerata L. 2. D.

♂. *speciosa* Hornem. 1835 et ex h. Paris. 1837.

gracilis Forst. v. *Wahlenbergia*.

gummifera W. (*sarmatica* Ker.) 2. D. 1835.

latifolia L. (*macrantha* Horn.) 2. D. ex h. Hafn. 1837.

♂. fl. albo ex h. Gryph. 1837.

lamiifolia M. B. 2. D. F. fere
 2 uln. alt.
Lorei Pollin. ☉. D. ex hort.
 Kil. 1836.
 β. fl. albo. 1837.
Lychnitis Hornem. 2. D. ex
 h. Hafn. 1837.
macrostachys W. ♂. 2. F. D.
 ex h. Gryph. 1837.
marcupiiflora Fisch.; D. 1826.
Medium L. ♂. D. bi-tripetal.
 β. fl. albo.
neglecta R. et Sch.; ex h. Crac.
 1837.
nutans Sieb. ♂. D. F. ex h.
 Hafn. 1837.
obliqua Jacq. ♂. D. F. fere
 2 uln. alt.
obliquifolia Tenor. 2. D. 1826.
pannonica W. ♂. 2. D. ex h.
 Gryph. 1837.
pendula M. B. 2. D. 1826 et
 ex h. Gryph. 1837.
pentagona L. ☉. D. 1827.
peregrina L. ♂. F. T. ex h.
 Paris. 1837.
perfoliata L. ☉. D. ex hh. Kil.
 1836, Paris. 1837.
persicifolia L. 2. D.
petrea L. 2. D. ex h. Gryph.
 1837.
planiflora W. (*versicolor* Sibth.)
 2. F. 1826.
pulcherrima Schrank. 2. D.
 ex h. Gryph. 1837.
pulla L. 2. F. 5-6 poll. alt.
 intr. 1833.
punctata Lam. 2. D. 1826.
pyramidalis L. ♂. 2. T. F.
 β. fl. albo.
rapunculoides L. 2. D. spont.
rotundifolia L. 2. D. spont.
sibirica L. ♂. 2. D. F. 1824-
 1826.
speculum L. ☉. D.
stylosa Lam. 2. D. 1½ uln. alt.
svaveolens 2. D. ex h. Hafn.
 1836; sub nom. *C. ucrani-*
ca in h. Lund. 1826 culta.
taurica Kosteletzky. 2. D. ex
 h. Gryph. 1837.
Thaliana Wallr. 2. D. ex h.
 Gryph. 1837.

Trachelium L. 2. D. ex hort.
 Gryph. 1837.
 species.

Campelia
Zanonia Rich. (*Commetina* L.)
 ♀. C.

Canarina
Campanula L. 2. T. olim in
 horto culta, hodie ibi n.
 exstat.

Canna
angustifolia L. 2. C. ex hort.
 Hafn. 1837.
glauca L. 2. C. de h. Hamb.
 1837 emta.
indica Ait. (*rubra* W.) 2. C.
limbata Rosc. 2. C. ex hort.
 Hafn. 1837.
lutea Rosc. 2. C. ex h. Crac.
 1837.
orientalis Rosc. (*chinensis* W.)
 2. C. ex h. Crac. 1837.
nepalensis Wallich. 2. C. ex
 hh. Kil. 1836, Crac. 1837.
patens Rosc. Spr. 2. C. ex hh.
 Kil. 1836, Hafn. 1837.

Cannabis (Hampa).
sativa L. ☉. D.
 β. gigantea.

Cantua
breviflora Juss. v. *Gilia laciniata*.
Hoitzia W. v. *Hoitzia coccinea*.

Capparis

Capraria

biflora L. 2. C. 1835.

Caprifolium

hispidulum ♀. T. sub. hoc no-
 mine ex Anglia commun.
 1837. cur Cl. J. Agardh.

Capsella

Bursa Pastoris Mönch (*Thlaspi*
 L.) ☉. D. spont.

Capsicum

annuum L. (Spansk Pepper).
 ☉. C.

cerasiforme Mill. ♀. C. ex h.
Kil. 1836.
grossum W. ♂. C. ex h. Kil.
1836.

Caragana (Ärteträd).
arborescens Lam. (*Robinia Caragana* L.) ♀. D. 9 uln. alta.
Chamlagu Lam. ♀. D. 2 uln. alta.
frutescens Cand. (*Robinia* L.) ♀. D. 5 uln. alt.
jubata Poir. ♀. F. D. ex hort. Petr. 1836.
pygmaea Cand. ♀. T. ex hort. Paris. 1837.

Cardamine
Lunaria L. v. *Ricotia*.

Cardiospermum
Corindum L. ♂. C. T. 1836.
Halicacabum L. ♂. C. 1836.

Carduus (Tistel).
acanthoides L. ♂. ♂. D. spont.
argentatus L. ♂. D. 1809.
arvensis Curt. Fr. (*Serratula* L. *Cirsium* Spr.) ♂. 2. D. sp.
crispus L. ♂. D. spont.
marianus L. (*Silybum* Spr.) ♂. ♂. D.
pycnocephalus L. ♂. D. 1811.

Carex (Starr).
acuta L. 2. D. 1826.
atrata L. 2. F. D. 1826.
depauperata Gooden. 2. F. ex h. Paris. 1837.
digitata L. ? 2. D. 1826.
distans L. 2. D. ex h. Paris. 1837.
divulsa Gooden. 2. D. 1826.
Drymeia L. Suppl. 2. D. 1826.
lagopodioides W. 2. D. 1826.
muricata L. 2. D. 1828, et ex h. Paris. 1837.
nitida Host. 2. D. ex h. Paris. 1837.
præcox Jacq. 2. D.
recurva Hornem. 2. D. 1808. 1826.
Schreberi W. 2. D. 1827.

stellulata Gooden. 2. D. ex h. Paris. 1837.

Carlina
acanthifolia All. 2. D. intr. 1835.
acaulis L. 2. D. 1820–1826.
vulgaris L. ♂. D.

Carmichaelia
australis R. Br. (*Lotus arboreus* Forst. *Achyronia arborea* Spr.) ♀. T. 2 uln. alt. floret Junio, intr. 1833 ex h. Hafn., et altera vice 1837 ex h. Kew. commun.

Carpesium
cernuum L. 2. D. F. ex h. Paris. 1837.

Carpinus
Betulus L. (Afvenbok) ♀. D.
orientalis Lam. ♀. D. 2½ uln. alta.

Carthamus (Safflor).
creticus L. (*Centaurea* Spr.) ♂. T. ex h. Paris. 1837.
Oxyacantha M. B. ♂. T. ex hort. Petrop. 1836.
tinctorius L. ♂. D.

Cassia
bicapsularis L. ♀. C.
floribunda Cav. (*corymbosa* Ort.) ♀. C.
obtusifolia L. ♀. C.

Castanea
vesca Gärtn. (*Fagus castanea* L.) ♀. D. 16 uln. alta; vetus horti nostri in loco ventis subducto plantat., bene viget et quotannis floret, sed fructus non dedit maturos.

Casuarina

equisetifolia Forst. $\frac{1}{2}$ T. ex h.
Hafnise a D. Mörch 1836.
nobis mis.

Catalpa

Syringæfolia Sims (*Bignonia*
Catalpa L.) $\frac{1}{2}$ T. D., in
nov. nostr. arb. plantat. 3
uln. alt.

Catananche

bicolor H. Hamb. $\frac{1}{2}$ T. ex
Anglia 1837 intr.
cærulea L. $\frac{1}{2}$ T. D.
lutea L. \odot D. ex h. Paris.
1837.

Caucalis

Anthriscus W. v. *Torilis*.
arvensis Huds. v. *Torilis helvetica*.
latifolia L. \odot D. ex h. Paris.
1837.
leptophylla L. (*humilis* Jacq.)
 \odot D. ex h. Paris. 1837.
platycarpus L. \odot D. ex hort.
Hafn. 1837.

Ceanothus

microphyllus Mx. $\frac{1}{2}$ D. ex h.
Paris. 1837.
ovatus Desf. $\frac{1}{2}$ D. ex hort.
Paris. 1837.

Celastrus

buxifolia L. $\frac{1}{2}$ T. $1\frac{1}{2}$ uln. alt.

Celosia

argentea L. \odot C. ex h. Hafn.
1836.
 β . ex h. Hafn. 1836.
castrensis L. \odot C. ex h. Hafn.
1836.
coccinea L. \odot C. ex h. Hafn.
1836.
cristata W. \odot C. var fl. ru-
bro et flavo.
margaritacea L. \odot C. ex h.
Hafn. 1836.
stricta Hornem. \odot C. ex h.
Hafn. 1837.
trigyna L. \odot C. ex h. Kil.
1834.

Celsia

grandiflora $\frac{1}{2}$ T. ex h. Crac.
1837.
orientalis L. \odot D.

persica, sub hoc nom. ex hort.
Petrop. 1836.
urticæfolia Curt. v. *Hemimeris*

Celtis

orientalis L. $\frac{1}{2}$ T. ex Anglia
1837.

Cenchrus

ciliaris L. (*Pennisetum cenchroi-*
das Rich.) \odot T. ex hort.
Paris. 1837.
setosus Sw. (*Pennisetum* Rich.
Panic. violaceum Lam.) \odot .
D. ex h. Crac. 1837.
Spinifex Cav. \odot T. ex hort.
Paris. 1837.
tribuloides L. $\frac{1}{2}$ F. ex hh.
Hafn. et Paris. 1837.

Centaurea (Klint).

egyptiaca Retz. Pugill. 1810
(*pubescens* W.?) $\frac{1}{2}$ D. F.
hodie horti exsul.
alpina L. $\frac{1}{2}$ D. ex h. Bonn.
1836.
amara L. $\frac{1}{2}$ D. 1836. $1\frac{1}{2}$ uln.
alta.
americana; \odot D. ex h. Paris.
1837. cur. Cl. J. Ag.
atropurpurea Kit. $\frac{1}{2}$ D. ex h.
Paris. 1837.
aurea Ait. $\frac{1}{2}$ F. $1\frac{1}{2}$ uln. alt.
1835.

Behen L. v. *Serratula*.

benedicta L. \odot D.
candidissima Lam. (*oineraria*
L.) \odot ? $\frac{1}{2}$ D. ex h. Pa-
ris. 1837.

chilensis Spr. \odot sub h. nom.
ex h. Crac. 1837.

collina L. $\frac{1}{2}$ D. ex h. Paris.
1837.

coriacea Kit. $\frac{1}{2}$ D.
Crocodilium L. \odot D. 1835.
 1 uln. alt.

Cyanus L. (Bläklint) \odot D.
fl. cærul., alb., rubr., aut
violac.

dealbata W. $\frac{1}{2}$ D. 1822. 1835,
et ex h. Paris. 1837.

depressa M. B. \odot D. 1835.
etiam ex h. Petr. 1836.

diluta Ait. (*elongata* Spr.) \odot .
D. ex hort. Paris. 1837.

- ferox* Desf. 2. D. ex. h. Petr. 1836.
Galactites L. (*Galactites tomentosa* Mönch.) ☉. D. ex h. Paris. 1837.
Jacea L. 2. D.
intybacea Lam. 2. D. 2 uln. alta.
macrocephala W. 2. D. ex h. Paris. 1837.
melitensis L. ☉. D. 1809.
montana L. (*mollis* Kit.) 2. D. 1833, etiam ex h. Paris. 1837. fl. caerule. aut albo.
moschata L. (*glauca* W.) ☉. ♂. D. 1 uln. alt.
 ♂. fl. albo.
napifolia L. ☉. D. ex h. Paris. 1837.
nigrescens W. 2. D. ex hort. Paris. 1837.
orientalis L. 2. D. ex h. Paris. 1837.
pectinata L. 2. D. 1826.
persica; ex h. Petrop. sub h. nomin. 1836.
pulchella; sub h. nom. ex h. Petrop. miss. 1836.
reflexa Lam. 2. D. ex h. Petr. 1836 commun.
ruthenica Lam. 2. D. ex h. Paris. 1837.
salicifolia W. 2. D. ex hort. Petrop. 1836.
salmantica L. ☉. D. ex hort. Paris 1837. fl. rubr.
 ♂. fl. flavo-albido.
scabiosa L. 2. D.
sempervirens L. ♂. 2. F. ex h. Paris. 1837.
splendens L. ♂. 2. D. 1½ uln. alt. 1835.
sqvarrosa W. 2. F. D. ex h. Petrop. 1836.
strobilacea Scop. (an *alba* L.?) 2. D. 1835. 2 uln. alt.
svaveolens W. ☉. D.
sulphurea W. ☉. D. ex hort. Paris. 1837.

Centranthus

- Calcitrapa* Dufresne (*Valeriana* L.) ☉. D. ex h. Kil. 1836.
ruber Cand. (*Valeriana* L.) 2. D.
 ♂. angustifolia Hornem.

Cephalaria v. Suocisa.

Cephalophora

- aromatica* Schrö. v. *Grahamia*.

Cerastium

- arvense* L. 2. D.
Biebersteinianum; D. ex hort. Paris. 1837. J. Ag.
collinum Cand. 2. D. ex h. Paris. 1837.
dichotomum L. ☉. D. ex h. Paris. 1837.
pennsylvanicum Hornem. 2. D. 1819–1826.
perfoliatum L. ☉. D. ex h. Paris. 1837.
tauricum Spr. 2. D. 1824.
tomentosum L. (*repens* M. B.) 2. D. 1805; etiam ex h. Paris. 1837.
viscidum; D. sub hoc nom. ex h. Paris. 1837.
vulgatum L. ☉. D. spont.

Cerasus v. Prunus.

Ceratocarpus

- arenarius* L. ☉. D. ex h. Petropol. 1836.

Ceratonia (Johannisbröd).

- Siliqua* L. ♂. T. F.

Ceratospermum

- arenarium*; sub h. nom. ex h. Paris. 1837.

Cercis

- canadensis* L. ♂. D. quotannis abunde floret. 2½ uln. alt. intr. 1830, seminib. educat.
Siliquastrum L. ♂. D. in nov. arbor. quotannis floret, 2½ uln. alt. intr. 1830.

Cérinthe (Vaxblomma).

- major* L. ☉. D.
minor L. 2. D.

Ceropogon

- glabrum*, sub hoc nom. ex h. Paris. 1837.

Cestrum

fastigiatum Jacqv. $\frac{1}{2}$. T. C.
pendulinum Jacqv. $\frac{1}{2}$. T. C.

Chærophyllum

angulatum Kit. $\frac{1}{2}$. D. semina
 ex h. Hafn. miss. 1837.
aromaticum L. v. *Myrrhis*.
aureum L. v. *Myrrhis*.
Cerefolium Fr. (*sativum* Spr.
Scandix Cerefol L.) \odot . D.
coloratum L. v. *Myrrhis*.
hirsutum L. v. *Myrrhis*.
bulbosum L. (*Myrrhis* Spr.) \odot .
 D. 1810. etiam ex hort.
 Crac. 1837.
millefolium; D. ex h. Paris.
 1837.
sylvestre L. $\frac{1}{2}$. D.
temulum L. v. *Myrrhis*.

Chamædorea

Schiedeana Booth. $\frac{1}{2}$. C. fere 2
 uln. alt. fl. d. 6 Jan. 1838;
 intr. 1833. ex h. Hamb.

Chamærops

humilis L. (*Phoenix* Cav.) $\frac{1}{2}$. C.

Chardinia

zeranthemoides Desf. v. *zeran-*
themum orientale.

Cheiranthus (Löfkoja).

alpinus L. $\frac{1}{2}$. D. 1808 - 1819.
annuus L., *fenestralis* L., *in-*
cunus, L. et *tricuspidatus*
 L. v. *Mutthiola*.

Cheiri L. $\frac{1}{2}$. D.

β . fl. pleno.

γ . fruticosus L. $\frac{1}{2}$.

collinus M. B. v. *Erysimum*.

helveticus Jacqv. v. *Erysimum*.

maritimus L. v. *Malcolmia*.

mollis Hornem. v. *Arabis albida*.

litoreus L. v. *Malcolmia*.

Chelidonium (Svalört).

corniculatum L. v. *Glaucium*
phoeniceum.

Glaucium L. v. *Glaucium luteum*.

majus L. $\frac{1}{2}$. D.

β . *luciniatum* Mill.

Chelone

barbata Cav. $\frac{1}{2}$. T. 1812

campanulata Cav. v. *Pentstemon*.

Digitalis Spr. (*Pentstemon* Nutt.)

$\frac{1}{2}$. F. ex h. Paris. 1837.

gracilis Spr. (*Pentstemon* Nutt.)
 C. 1836.

nemorosa Booth. $\frac{1}{2}$. T. ex An-
 lia, etiam ex h. Paris.
 1837.

Pentstemon L. v. *Pentstemon pu-*
bescens.

Chenopodium

(Mell. Molla)

album L. \odot . D.

β . *viride*.

aristatum L. \odot . D.

Bonus Henricus L. $\frac{1}{2}$. D.

Botrys L. \odot . D.

carthagenense Zucc. Hornem,
(opulifolium Schrd. Spr.)

\odot . D. ex h. Hafn. 1837.

chilense \odot . D. sub hoc nom.

ex h. Bonn. 1836.

glaucum L. \odot . D. ; uln. alt.

hircinum \odot . D. sub hoc nom.

ex h. Bonn. 1836.

hybridum L. \odot . D. saepe 1
 uln. alt.

murale L. \odot . D.

olidum Sm. (*Ch. Vulvaria* L.)
 \odot . D.

polyspermum L. \odot . D.

Quinoa W. \odot . T. D. ex h.
 Paris. 1837.

rubrum L. \odot . D.

trifidum, \odot . F. ex h. Paris.
 1837.

Chionanthus

virginica L. $\frac{1}{2}$. olim (1798)

in horto sub *Dio plantata*
 viguit, jam evauuit.

Chironia

Barclayana Booth. $\frac{1}{2}$. T. intr.
 1836 ex. h. Hamb.

Chloris

compressa Cand. (*virgata* Sw.
 Spr.) \odot . C. ex h. Par. 1837.

polyactyla Sw. (*Andropogon*
L.) ♂. G. ex hort. Paris.
1837.

radiata Sw. (*Agrastis* L.) ♂.
C. 1806; et altera vice ex
h. Paris. 1837 cur. Clar.
J. Ag.

Chondrilla

hispida, sub hoc nom. ex h.
Paris. 1837.

junceae L. 2. D. 2 uln. alt.

Chorispora

stricta Cand. (*Raphanus* Fisch.)

♂. ex hort. Petr. 1836.

tenella Cand. (*Hesperis* L. ex
Lam.) ♂. D. ex h. Paris.
1837.

Chrysanthemum

Balsanita L. v. *Pyrethrum*.

carinatum Schpösb. ♂. D.

caucasicum 2. D. ex h. Paris.
1837.

coccineum W. v. *Pyrethrum* ro-
seum.

coronarium L. ♂. D.

β. fl. albo.

Gmelini; sub hoc nom. ex h.
Petr. 1836.

indicum L. (*Anthemis artemi-
siaefolia* W.) 2. β. T. fl.
plenit. et color. var.

inodorum L. v. *Pyrethrum*.

leucanthemum L. 2. D. spont.
maximum Cand. Horn. 2. D.
1826.

multifidum; sub hoc nom. ex
hort. Paris. 1837.

pulverulentum Pers. ♂. D. ex
h. Paris. 1837.

pumilum W. (*glabratum* Thb.)
♂. D. 1820.

Robxburghii H. Hefn. ♂. D.
1835. 1½ uln. alt.

segetum L. ♂. D. spont.
species.

Chrysocoma

(Gullborste).

Linosyris L. 2. D. 1½ uln. alt.

Cichorium

Endivia L. ♂. 3. D. 1½ uln.
alt.

β. crispum.

Intybus L. 2. D.

pumilum Jacq. ♂. D. ex h.
Paris. 1837.

Cimicifuga

foetida 2. D. ex hort. Paris.
1837.

Cineraria

smelloides L. 2. β. T. 1½
uln. alt.

campestris Retz. 2. D. 6-7
poll. alt.

cruenta Herit. 2. β. T. 26
poll. alt. intr. 1833.

geifolia L. β. T. ex h. Paris.
1837.

gibbosa Gusson. β. F. ex h.
Paris. 1837.

maritima L. β. T. in horto
1½ uln. alt.

populifolia Lam. (*aurita* Herit.)
β. T.

Cinna

arundinacea L. (*Agrastis Cinna*
Lam.) 2. D.

Circaea

lutetiana L. 2. D. inter spon-
taneas horti nostri colo-
nias jam numeranda.

Cirsium

lanceolatum Scop. (*Carduus* L.)
3. D.

Cissus

species β. C. ex Ind. occid.

Cistus

albidus L. β. T. 1833, etiam
ex h. Hafn. 1836.

creticus L. (*tauricus* Presl.) β.
T. ex h. Hafn. 1836.

grandiflorus Scop. v. *Hekanihem*
vulgare.

hirsutus Lam. β. T. ex hort.
Paris. 1837.

incanus L. β. T. ex h. Kil.
1836.

laxus Ait. *capensis* L. ? β. T.
ex h. Kil. 1836.

Ledon Lam. β. T. 1833; ex
h. Kil. 1836.

parviflorus Lam. $\frac{1}{2}$. T. ex h. Hafn. 1826.
salviaefolius L. $\frac{1}{2}$. T. ex hort. Paris. 1837.
vaginatus Ait. $\frac{1}{2}$. T. ex h. Kil. 1836; etiam sub nom. *C. symphytifolius* ex h. Paris. 1837.
villosus Lam. (*pilosus* L.) $\frac{1}{2}$. T. ex hort. Paris. 1837.

Citharexylon

quadrangulare Jacq. $\frac{1}{2}$. C.

Citrus

Aurantium L. (Pomeransträd) $\frac{1}{2}$. T.
medica L. (Citronträd), $\frac{1}{2}$. T.

Clarkea

elegans Lindl. \odot . T. D. intr. 1834 ex Angl.
gauroides Dougl. \odot . T. D. ex h. Paris. 1837.
pulchella Pursh. \odot . T. D. fl. rubro.
 β . fl. albo, ex h. Kilens. 1836.

Claytonia

perfoliata Don. Spr. \odot . D. 1805—1837; sub nom. *Cl. cubensis* ex h. Paris. 1837 forte eadem.
sibirica $\frac{1}{2}$. F. ex hort. Paris. 1837.

Clematis

angustifolia Jacq. $\frac{1}{2}$. D. ex h. Gryph. 1837.
aristata R. Br. $\frac{1}{2}$. T. ex h. Kew. 1837, cur J. Ag.
campaniflora Brot. $\frac{1}{2}$. ex hort. Gryph. a Cel. Hornschuch 1837 nob. communic.
corymbosa, in frigid. nostro colitur sub hoc nom. species, quam non nisi *Cl. erecta* videtur.
dioica L. $\frac{1}{2}$. olim in hort. sub *Dio plantata*, hodie ibi non exstat.
erecta (recta) L. $\frac{1}{2}$. D. intr. 1805.

β . *angustifolia* ex h. Gryph. 1837.

integrifolia L. $\frac{1}{2}$. D. 1826.
 \odot etiam ex h. Gryph. 1837.
lasiantha Fisch. $\frac{1}{2}$. ex h. Gryph. 1837.

maritima L. ex h. Gryph. 1837 miss. sed mox emoriebatur.
montana Buchan. $\frac{1}{2}$. T. ex Angl. 1837.

Nepalensis Cand. $\frac{1}{2}$. T. ex Angl. 1837.

virginiana L. $\frac{1}{2}$. in horto olim culta (teste herbar.), hodie abest.

Vitalba L. $\frac{1}{2}$. D.

Viticella L. $\frac{1}{2}$. D. ex h. Gryph. 1837.

Cleome

aculeata L. \odot . C. 1806, teste herbario, hodie horti pro-fuga.

Atkinsoniana. \odot . C. ex hort. Paris. 1837.

canadensis Hort. Betv. \odot . C. intr. 1837 ex hort. Kil., a Cel. Nolte miss.

dodecandra L. \odot . C. 1808, teste herbar.

graveolens Rafin. (*viscosa* L.) \odot . C. 1805; ex h. Paris. etiam 1837.

pentaphylla L. \odot . C.

speciosa Kunth. \odot . C. ex h. Kil. 1836. 1 uln. alt.

viminea Schrad. \odot . C. ex h. Paris. 1837.

violacea L. \odot . T. F. 1806, et iterum intr. ex h. Bonn. 1836 cur Cel. C. A. Ag.

virgata Stev. \odot . T. intr. 1836 ex h. Petrop.

Clerodendron

fragrans Vent. $\frac{1}{2}$. C. 1819, hodie horti exsul.

Clianthus

punicus Booth. $\frac{1}{2}$. T. intr. 1837 ex hh. Hamburg. et Edinb.

Clinopodium

egyptiacum Lam. $\frac{1}{2}$. F. 1811.
carthagineum ex h. Hafn. 1837.
vulgare L. $\frac{1}{2}$. D. 1827.

Clintonia

elegans Dougl. ☉. T. ex hort.
Kil. 1836.

Clitoria

virginiana L. 2. T. ex Angl.
1837.

Cluytia

pulchella L. 2. T.

Clypeola

Jonthlaspi L. ☉. D. 1808.
1837.

maritima L. (*Alyssum* Lam.) ☉.
2. 2. D. F. 1819, et ex
hh. Paris. et Crac. 1837.

Cneorum

tricoceon L. 2. T. 2. uln. alt.

Cnicus

giganteus; D. ex h. Paris. 1837.
monspessulanus W. (*Carduus*
L.) 2. D. 1826.

palustris Hoffm. (*Carduus* L.
Cirsium Spr.) 2. D.

pannonicus. D. ex h. Paris.
1837. sub hoc nom.

stellatus W. (*Carduus* L.) ☉.
D. ex h. Paris. 1837.

syriacus W. (*Carduus* L.) ☉.
D. 1812. et ex h. Paris.
1837.

Cnidium

Silau Spr. (*Peucedanum* L.)
2. D.

Cobæa

scandens Cav. 2. 2. T. F. ex
h. Hamb. 1836.

Cochlearia (Skedört).

Armoracia L. (*Nasturtium* Fr.
Pepparrot), 2. D.

Draba L. v. *Lepidium*.

Coffea (Kaffeträd).

arabica L. 2. C.

Coix

Lacrima L. ☉. C.

Colchicum (Tidlösa).

autumnale L. 2. D.

Collinsia

bicolor Benth. ☉. D. intr.
1837. ex h. Kil.

grandiflora Dougl. ☉. 2. D.
intr. 1835. ex h. Kil.

parviflora Lindl. ☉. 2. F. D.
1835.

Collinsonia

canadensis L. 2. D.

Collomia

coccinea Lehm. v. *Hoitzia*.

grandiflora Dougl. ☉. T. ex
h. Hafn. 1837.

heterophylla Hook ☉. T. ex
h. Kil. 1836.

linearis Nutt. (*Hoitzia* Spr.
Phlox Cav.) ☉. D. ex h.
Kil. 1836.

pinnatifida W. Dougl. ex h.
Gryph. 1837.

pumila H. Hafn. ☉. T. D.

Colutea

arborescens L. 2. D. in Para-
dislyckan.

β. *media* W. 3 uln. alt.

cruenta Ait. (*orientalis* du Roi)
2. D. intr. 1832.

frutescens v. *Sutherlandia*.

Pocockii Ait. 2. D. 1½ uln.
alt. intr. 1832 seminib.
educat.

Comarum

fragarinoides W. v. *Potentilla*
Fragaria.

Commelina

angustifolia Mx. (*truncata* W.)
2. T.

caroliniana Walt. 2. T. F.

clandestina Mart. 2. T. intr.
ex h. Crac. 1837.

coelestis W. 2. T.

communis L. ☉. F. D.

erecta L. 2. T. F.

rubens Red. (*elegans* Humb. *pallida* W.) 2. T. ex hort.
Crac. intr. 1837.

- tuberosa L. 2. T. ex h. Paris. 1837.
- Conioselinum**
Carvisfolia Fr. v. *Selinum*.
- Conium** (Odört).
 croaticum Kit. (*macul.* var. Spr.) ♂. D. 1826. et altera vice ex h. Hafn. 1837.
maculatum L. ♂. D. spont.
- Conobea**
 species.
- Convallaria**
 (Konvalje).
bifolia L. v. *Maianthemum*.
latifolia L. 2. D. 1826, hodie horti exsul.
majalis L. 2. D.
 β. fl. plen. etiam roseo.
multiflora L. 2. D.
Polygonatum L. 2. D.
verticillata L. 2. D.
- Convolvulus**
 (Binda. Vinda).
arvensis L. 2. D.
carneus Spr. (*Ipomoea* Jacqv.)
 ⊙. D. ex h. Crac. 1837.
chilensis Pers. D. ex h. Paris. 1837.
Cneorum L. 2. T. F. 1 uln. alt.
coccineus Spr. (*Ipomoea* L.) ⊙
 C. ex h. Crac. 1837.
Dillenii Desrous. ⊙. D. flor. purp.
 β. fl. albo.
hederaceus L. Spr. (*Ipomoea* Jacqv. *Ipom. barbata* Roth.)
 ⊙. C. ex hh. Crac. et Hafn. 1837.
Icucanthus Spr. (*Ipomoea* Jacqv.)
 ⊙. C. ex h. Hafn. 1837.
luteolus Spr. (*Ipomoea* Jacqv.)
 ⊙. C. ex h. Hafn. 1835.
major Booth. ⊙. T. 1835. 1½ uln. alt. flor. rubro-cærul.
mutabilis Spr. (*Ipomoea* Ker.)
 C. ex h. Crac. 1837.
Nil L. W. ⊙. C. 1820. et nova vice ex h. Hafn. 1837.
pentapetaloides L. ⊙. D. ex h. Hafn. 1835.
pseudosiculus Cav. (*alongatus*

- W.) ⊙. D. ex h. Paris. 1837.
- Pes tigridis* Spr. (*Ipomoea* L.)
 ⊙. C. ex h. Hafn. 1835.
purpureus L. (*Ipomoea* Lam.) ⊙.
 D. 1835.
quinatus Spr. (*Ipom. Caradine* L.) ⊙. D. ex hort. Crac. 1837.
Scammonia L. 2. F. ex hort. Paris. 1837.
sepium L. 2. D. in horto sp.
sibiricus L. ⊙. D. ex h. Hafn. 1837.
siculus L. ⊙. D. ex h. Paris. 1837.
tricolor L. ⊙. D.
 β. fl. albo unicolore.
trilobus Desrous. (an *C. Nil* L.?) ⊙. C. ex h. Kilens. 1837.
- Conyza**
cinerea L. ⊙. C. 1808.
Gouani W. (*Erigeron* L.) 2.
 F. ex h. Paris. 1837.
longifolia; sub hoc nom. ex h. Paris. 1837.
sicula W. (*Erigeron* L.) ⊙.
 D. ex h. Paris. 1837.
squarrosa L. ♂. 2. D. 1813.
- Corchorus**
japonicus Thb. v. *Keria* (Keria Ag.).
siliquosus L. ⊙. 2. C. 1819
 in hort. cult., hodie profugus.
- Coreopsis**
Atkinsonii Booth. 2. F. D. 1-2 uln. alt.
auriculata L. 2. D.
tinctoria Nutt. v. *Calliopsis* bicolor.
Tripteris L. 2. D. 2½ uln. alt.
verticillata L. 2. D. ¾ uln. alt.
- Coriandrum** (Koriander).
sativum L. ⊙. D.

Coriaria

myrtifolia L. ♀. D.

Corispermum

hyssopifolium L. ♂. D.

polygaloides ♂. T. ex h. Petropol. 1836.

Cornucopiæ

cucullatum L. ♂. D. ex hort. Hafn. 1835.

Cornus

alba L. ♀. D. 4-5 uln. alt. Paradiislyckan.

alternifolia L. ♀. D. 1835.

florida L. ♀. D.

mascula L. ♀. D. Vetus horti incola, sub nomine *Olive*. träd ab hortulan. venditur.sanguinea L. ♀. D. 3 uln. alt. sericea Herit. (*C. Amomum* du Roi) ♀. in horto nostro1808 sub *Dio culta*, hodie ibi ignota.

sibirica Booth. ♀. D.

stricta Lam. ♀. D.

Coronillacretica L. (*Ornithopus*) ♂. D. 1835-1837.*Emerica* L. ♀. olim in horto sub *Dio* viguit (teste herbar.), hodie desideratur.glauca L. (*Ornithopus* Horn.). ♀. T.

securidaca L. ♂. D. ex h. Kil. 1835.

valentina L. ♀. T.

varia L. ♀. D.

Coronopusdidymus Spr. (*Lepidium* L.)♂. T., sub nom. *Senebiera pinnatifida* Cand ex hort. Kil. 1837. idem communicatus.**Corrigiola**

littoralis L. ♂. D. herbar.

Corydalis (*Fumaria* spec. L.)

aurea W. ♂. D.

bulbosa Cand. Fr. (*Halleri* W.) ♀. D.

cava Wahl. Fr. ♀. D. fl. rubro aut albo.

capnoides Pers. ♀. D. 1826.

claviculata Pers. ♂. D.

sahæcea Pers. ♀. D.

nobilis Pers. ♀. D.

sempervirens Pers. ♂. D.

Corylus

(Hassel. Nöthuske).

Avellana L. ♀. D.

tubulosa W. (*maxima* Mill.*Lambertsk Nöthuske* Svec.)

♀. D. 17 uln. alt.

Cotoneaster v. *Mespilus*.**Cotula**abrotanifolia W. v. *Lasiospermam*.

anthemoides L. ♂. D. ex h. Paris. 1837 cur. Cl. J. Ag.

aurea L. ♂. D. intr. 1834 ex h. Hafn.

coronopifolia L. ♂. D. 1808, et jam ex h. Paris. 1837.

maderaspata W. (*Artemisia* L. *Grangea* Lam.). ♂. C. 1819. 1820; quoque ex h. Paris. 1837.minima W. (*Grangea* H. Paris. *Artemisia* L.) ♂. C. 1819, et ex h. Paris. 1837.

nudicaulis Thb. ♂. C. ex h. Paris. 1837.

Cotyledon

coccinea Cav. ♀. ♀. T.

orbiculata L. (*C. ramosissima* Haw.) ♀. ♀. T.pinnata Lam. v. *Verea*.**Cousinia**

cynaroides; sub hoc nomine ex h. Petrop. 1836.

hystrix; ex h. Petrop. 1836.

leptophylla; sub hoc nom. ex h. Paris. 1837.

macrocephala Booth. ♀. D. 1836.

tenella; ex h. Petrop. 1836.

Crambe

hispanica L. ♂. D. 1808; et nuper ex h. Paris. 1837.

juncea M. B. ♀. D. ex hb. Petropol. 1836, et Paris. 1837.

maritima L. ♀. D.

strigosa Herit. ♀. T. 1826.

Crassula

- acutifolia* Lam. f. T.
andegaviensis Cand. C. ex h. Hafn. 1837.
arborescens Mill. (Cr. *Cotyledon* Jacq. *portulacacea* W.) f. T.
falcata L. (*Larochea* Cand.) f. T.
Magnolii Cand. T. ex h. Paris. 1837.
obvallata L. f. T. ex h. Paris. 1837.
perfoliata L. f. T.
spathulata Thb. (*flacida* Lam.) f. T.
tetragona L. f. T.

Cratægus v. *Mespilus* et *Sorbus*.

Cræpis (cfr. *Borckhausia*).

- agrestis* Kit. 1812.
biennis L. f. D.
coronopifolia Desf. v. *Tolpis*.
grandiflora W. suppl. Spr. D. 1836.
lappacea W. v. *Picris altissima* Delil.
parviflora Jacq. C. D. 1807. 1812 herb. ex h. Paris. 1837.
pinnatifida W. C. D. 2½ uln. alta.
pulchra L. (*Prenanthes hieracifolia* W.) C. D. 1835. 1½ uln. alt.
rigida Kit. 2. D. ex h. Paris. 1837.
tectorum L. C. D. spont.

Crinum

- africanum* L. v. *Agapanthus umbellatus*.
angustifolium L. suppl. v. *Cyranthus*.
giganteum Andr. v. *Amaryllis*.
lineare Thb. v. *Amaryllis revoluta*.

speciosum Thb. v. *Amaryll. purpurea*.

Cristaria (*Sida*).

Berteri, 2? C. sub hoc nom. ex h. Paris. 1837.

Crithmum

maritimum L. v. *Cuchrys*.

Crocus (*Saffransört*).

- luteus* Lam. (Cr. *masincus* Ker.) 2. D. 3 pollic. alt.
sativus Sm. 2. D.
susianus Curt. 2. D. 2½ poll. alt. intr. 1833. ex hort. Hamb.
vernus W. 2. D. 3 poll. alt.

Crotalaria

incana L. C. C. in hort. 1806 cult. test. herbar.

Crucianella

- angustifolia* L. C. D. ex h. Paris. 1837.
ciliata Lam. (*patula* W.) C. D. ex h. Hafn. 1823.
gilanica Trin. 2. ex h. Petr. 1836.
glomerata M. B. ex h. Petrop. 1836.
latifolia L. C. D. ex h. Paris. 1837.
patula L. C. D. ex h. Paris. 1837.
stylosa Trin. C. D. ex hort. Paris. 1837.
svaveolens Hortul. C. 2. D. ex hh. Petrop. 1836 et Paris. 1837.

Cryptostemma

- calendulaceum* R. B. (*Arctotis* L.) C.
hypochondriacus R. Br. (*Arctotis* L. Syst. *Arctotis tri-stis* L.) C. C. 1809; ex Kil. 1835.

Cucubalus (cfr. etiam *Silene*).

- angustifolius* Tenor. v. *Silene inflata*. var.
bacciferus L. 2. D.
catholicus L. v. *Silene*.
mollissimus L. v. *Silene*.

Cucumis

- deliciosus* Roth. ☉. C. intr.
 1833 ex h. Gryph.
Melo L. (Melon) ☉. C.
sativus L. (Gurka) ☉. D. plures varietat. in h. coluntur.

Cucurbita (Pompa).

- lagenaria* L. ☉. D. 1812.
Pepo L. ☉. T. adsunt plures varietates.

Cuphea

- viscoccissima* Jacq. ☉. C. ex h. Kil. 1836, Paris. 1837.
 Caulis undique insectis minutissimis: Omaliis, Tipulariis cet. adglutinati ob-situs.

Cupressus (Cypress).

- africana* H. Hamb. 2. T.
sempervirens L. 2. F. D.
 β. *pendula* Thb. 2. F. intr.
 1836 ex h. Hamb.
thyoides L. v. *Thuja sphaerocarpa*.

Curculigo

- species*. 2. C. intr. 1833.

Curcuma**Cuscuta****Cyclamen**

- europaeum* L. 2. T. F.
hederifolium Ait. 2. T.
persicum Mill. 2. T.

Cydonia (Qvitten).

- vulgaris* Pers. (*Pyrus Cydonia* L.) 2. D.
 β. *lusitanica*. in nov. arbor.
 γ. *pyriformis*.

Cynanchum

- acutum* L. 2. F.
fuscatum Link. (*Asclepias* W.)
 2. T. intr. 1834 ex hort. Hafn.

- nigrum* Pers. (*Asclepias* L.) 2.
 D. 1½ uln. alt.
rostratum Vahl. v. *Gonolobus*.
Vincetoxicum Pers. Fr. (*Asclepias* L.) 2. D.

Cynara

- horrida* Ait. 2. F. ex h. Paris. 1837.
Scolymus L. (Kronärtskock)
 2. D.

Cynoglossum

(Hundtunga).

- apenninum* L. 2. D.
bicolor W. 2. D. ex h. Paris.
 1837. cur. Cl. J. Ag.
divaricatum Lehm. D. ex h. Petrop. 1836.
elongatum Hornem. 2. D. ex h. Paris. 1837.
linifolium L. ☉. D.
officinale L. 2. D. spont.
Omphalodes L. 2. D.
pictum Ait. 2. D.
virginicum L. 2. F. 1826.

Cynosurus

- cristatus* L. 2. D. spont.
retroflexus Vahl. (*Dineba aegyptiaca* Delil. *arabica* P. B.
 ☉. C. 1811. herbar. ex h. Paris. 1837.
virgatus L. v. *Eleusine*.

Cyperus

- elegans* Vahl. T. 1808 test. herbar. nostro.
esculentus L. 2. D. ex h. Paris. 1837. cur. J. Ag.
strigosus L. 2. F. ex h. Paris. 1837.

Cyrilla

- pulchella* Herit. v. *Trevirania coccinea*.

Cyrtanthus

- angustifolius* L. (*Crinum* L. suppl. *Amaryllis cylindracea* Herit.) 2. C.

Cysticapnos

- africana* Gärt. (*Fumaria vesicaria* L.) ☉. D. ex h. Hafn. 1835.

Cytisus

- alpinus Schm. $\frac{1}{2}$ D. 5 uln.
alt. Paradislyckan.
austriacus L. $\frac{1}{2}$ D. 1826.
biflorus Herit. $\frac{1}{2}$ F. 1 $\frac{1}{2}$ uln.
alt. flor. Jun.
calycinus M. B. $\frac{1}{2}$ F. ex h.
Paris. 1837.
candicans Lam. (*Genista* L.) $\frac{1}{2}$
T. F. ex h. Kil. 1836.
capitatus Jacqv. $\frac{1}{2}$ F. D. 1 $\frac{1}{2}$
uln. alt. flor. Jun.
elongatus Kit. $\frac{1}{2}$ D. 1—2 uln.
alt. seminib. educat. 1835.
hirsutus L. $\frac{1}{2}$ D. 2 uln. alt.
intr. 1826.
Labraum L. (Gullregn) $\frac{1}{2}$
D. per multos annos in
horto cultus a frigore non
laesus.
leucanthus Kit. $\frac{1}{2}$ D. ex h.
Paris. 1837.
nigricans L. $\frac{1}{2}$ D.
purpureus Scop. $\frac{1}{2}$ D. intr.
1836; hodie $\frac{3}{4}$ uln. alt.
sessilifolius L. $\frac{1}{2}$ D. in nov.
arbor. luxurians, 3 uln.
alt., etiam in Paradislyc-
kan.
spinosis Spr. v. *Spartium*.
supinus Jacqv. $\frac{1}{2}$ D. 2-3 uln.
alt.
volgarensis L. suppl. (*Adeno-
carpus volgensis* Spr.) $\frac{1}{2}$ F.
ex h. Paris. 1837.

Dactylis

- glomerata L. $\frac{1}{2}$ D. spont.

Dahlia

- erinita Thb. v. *Trichocladus*.
variabilis Cav. v. *Georgina*.

Dalea

- Lagopus W. (*Psoralea* Cav.) T.
1811. herbar. nostr.

Daphne

- Cneorum L. $\frac{1}{2}$ T. 1819.
Laureola L. $\frac{1}{2}$ T. F. D.

- Mezerium L. (Kallehals. Ti-
bast.). $\frac{1}{2}$ D.

Datisca

- cannabina L. $\frac{1}{2}$ D.

Datura (Spikklubba).

- ceratocaula Jacqv. \odot . C. ex
h. Paris 1837.
fastuosa L. \odot . C. intr. 1833
ex h. Paris.
ferox L. \odot . D. ex h. Paris.
1837.
Metel L. \odot . C. ex h. Hafn. 1836.
Tatula L. \odot . D. ex h. Hafn.
1837.

Daucus

- Carota L. (Morot) $\frac{1}{2}$ D.
 β . radice alba.
gummifer Lam. $\frac{1}{2}$ D. 2 uln. alt.
muricatus L. \odot . $\frac{1}{2}$ F. ex h.
Paris. 1837.
pulcherrimus Koch. \odot . ex h.
Petr. 1836. sub hoc nom.

Davallia

- canariensis Sw. (*Polypodium* L.)
 $\frac{1}{2}$ T. 1836.

Delphinium

(Riddarsporre).

- Ajaxis L. \odot . D.
 β . fl. plenitud. et col. var.
albiflorum Cand. $\frac{1}{2}$ T. ex h.
Paris. 1837.
amoenum Stev. $\frac{1}{2}$ ex h. Gryph.
1837.
cheilanthum Fisch. $\frac{1}{2}$ ex h.
Gryph. 1837.
chinense Fisch. $\frac{1}{2}$ D. ex hh.
Kil. 1836, et Gryph. 1837.
 β . fl. albo.
ciliatum Stev. (*flexuosum* M.
B.) $\frac{1}{2}$ D. 1826, etiam ex
h. Gryph. 1837.
Consolida L. \odot . D.
crassifolium Schr. ex h. Grac.
1837.
dityocarpum Spr. (*laxiflorum*
Caud.) $\frac{1}{2}$ D. 1826; ex
hh. Gryph. et Paris. 1837.
divaricatum; ex hh. Bonn. et
Petr. 1836.

exaltatum Ait. 2. D. ex h. Gryph. 1837.
grandiflorum L. 2. D. ex hh. Crac. et Gryph. 1837.
 β. fl. alb.
 γ. fl. variegat.
 δ. fl. pleno, cærul.
hybridum W. 2. D. ex hort. Gryph. 1837.
intermedium Ait. (*elatum* L.) 2. D. 1826, et iterum ex hh. Crac. et Gryph. 1837.
 β. villesum.
montanum Cand. 2. D. ex h. Gryph. 1837.
palmatifidum Cand. 2. D. intr. 1837 ex h. Gryph. a Cel. Hornsch. nob. miss.
puniceum L. fil. 2. D. ex h. Gryph. 1837.
Requienii Cand. ♂. T. ex h. Paris. 1837.
recurvirostrum H. Gryph. 2. D. ex h. Gryph. 1837.
revolutum Desf. 2. D. ex h. Gryph. 1837.
speciosum M. B. 2. D. ex h. Gryph. 1837.
sulcatum Reichb. 2. D. ex h. Gryph. 1837.
ucrainicum Fisch. 2. D. ex h. Crac. 1837.
urceolatum Ait. 2. D. 1826. species.

Desmanthus

virgatus W. (*Mimosa* L.) ♀. C. intr. 1836. ex h. Kil. a Cel. Nolte nob. commun.

Desmochæta

atropurpurea Cand. (*Achyranthes lappucea* L.) ♀. C. 1819. herbar. nostr.

Deutzia

scabra Thunb. ♀. T. ex Anglia reportat. 1837 a Clar. J. Agardh.

Dianthus (Nejlika).

arenarius L. 2. D.
Armeria L. ☉. D.
 β. fl. alb.
 γ. gigantea; ex h. Hafniæ 1836.
Arrostii Bieb. 2. ex h. Gryph. 1837.
asper W. 2. D. herb. 1822.
barbatus L. 2. D. fl. col. var.
campestris M. B. 2. D. 1820; ex h. Paris. 1837.
Carthusianorum L. 2. D. ex h. Hafn. 1836.
Caryophyllus L. 2. D.
chinensis L. ♂. D.
collinus Kit. 2. D. ex hort. Hafn. 1836.
deltoides L. 2. D. ex h. Hafn. 1835.
 β. glaucus.
 γ. wolgensis Fisch. Horn. ex h. Hafn. 1836.
discolor Sims. (*collinus* var. Spr.) 2. D. ex h. Paris. 1837.
diutinus Kit. 2. D. 1826. ex hh. Hafn. et Paris. 1837.
dubius Hornem. 2. F. ex h. Hafn. 1835.
limbriatus M. B. 2. D. ex h. Paris. 1837.
Fischeri Spr. 2. D. ex h. Gr. 1837.
furcatus Balb. 2. ex h. Gryph. 1837. a Cel. Hornsch. nob. commun.
gracilis Sm. 2. 1822 herbar. nostr., hodie horti exsul.
hortensis Schrd. (*plumarius* var. Spr.) 2. D. 1826.
hungaricus H. Crac. 2. D. 1826, et ex hort. Crac. 1836.
moschatus hortulan. 2. D. ex h. Paris. 1837.
Mussini Hornem. 2. D. ex h. Gryph. 1837.
petraeus Kit. 2. D. ex h. Hafn. 1835.
plumarius L. 2. D.
prolifer L. ♂. 2. D. 1822, et ex h. Hafn. 1835.

punctatus Spr. 2. D. ex h.
Hafn. 1836.
serotinus Kit. 2. D. ex hort.
Hafn. 1835.
squarrosus M. B. Horn. 2. D.
ex h. Hafn. 1835.
superbus L. 2. D. 1826, etiam
ex h. Hafn. 1835.
tataricus H. Crac. 2? D. ex
h. Crac. 1837.
velutinus; ☉? D. ex h. Bonn.
1836 sub hoc nom.
species.

Dichondra

sericea Sw. 2. C.

Dictamnus

albus L. 2. D. intr. 1818.

Didiscus

caeruleus Hook (Umbellif.) ☉.
C. intr. 1836 ex h. Bonn.
cur. Cel. Pr. C. Agardh.

Diervilla

canadensis W. v. *Loxicera Dier-*
villa.

Digitalis

(Fingerborrsblomma).

ambigua L. fil. 2. D. ex hort.
Gr. 1837 a Cel. Hornsch.
nobis missa.

aurea Lindl. 2. D. 1826.

australis Tenor. 2. D. ex h.
Crac. 1837, etiam sub
nom. *D. micrantha* Roth.
ex hh. Kil. Paris. et Gr.
nob. 1837 commun.

canariensis L. 2. T. ex h. Gr.
1837. 1½ uln. alt.

eristachys Bess. 2. F. sub hoc
nom. ex h. Kil. 1835.

fuscescens Kit. 2. F. ex hort.
Kil. 1835.

gigantea H. Gryph. 2. D. intr.
1835. Flora 1836. p. 12;
2½ uln. alt.

laevigata Kit. 2. ♂. D. ex h.
Paris. 1837.

lanata Ehrh. ♂. D. ex h. Crac.
1837.

lutea L. 2. D. 1½ uln. alt. in
horto.

media Roth. 2. D. ex h. Crac.
1837.

mollissima; ex h. Paris. 1837
sub hoc nom.

ochroleuca Jacqv. 2. D. ex h.
Kil. 1836.

orientalis Lam. 2. D. ex h.
Kil. 1835.

purpurea L. ♂. D. saepe 1½
uln. alt.

β. fl. alb.

Thapsi L. 2. D. 1826.

tomentosa Link. ♂. ex hort.
Gryph. 1837.

truncata Estreich. 2. D. ex
h. Kil. 1835.

Digitaria

ciliaris W. (*Panicum* Retz.) ☉.
D. ex h. Hafn. 1837.

criogona Link. ex h. Hafniae
1837.

marginata Link. ex h. Hafniae
1837.

villosa Pers. ex h. Hafn. 1837.

Dillenia

scandens W. v. *Hibbertia volu-*
bilis.

Dineba

arabica P. B. v. *Cynosurus re-*
troflexus.

Dioscorea

Elephantopus Spr. (*Tamus* He-
rit.) 2. C.

Diosma

dioica Ker. 2. T.
serratifolia Curt. (*Barosma* W.)
2. T.

Diospyros

Lotus L. 2. F. D. ex h. Hamb.
1836.

virginiana L. 2. D. coeli no-
stri temperiem bene tole-
rans, intr. 1834 ex hort.
Hamb.; hodie 2 uln. alt.

Dipsacus

ferox Lois. ♂. D. ex h. Hafn. 1835.

Fullonum Mill. ♂. D. 1807. herbar. nostr.

laciniatus L. ♂. D. ex h. Paris 1837.

pilosus L. ♂. D. in nov. arb. jam spontan. et infestum vitium.

sylvestris Mill. L. ♂. D. 1826.

Disandra

prostrata L. Syst. (*Sibthorpia peregrina* Spr.) ♀. F. 1808. herb. nostr.

Dodartia

orientalis L. ♀. ♀. D. ex h. Petrop. 1836.

Dodecatheon

Meadia L. ♀. F. D.

Dolichos

Catjang L. ♂. C.

Columbianus Ag. ♂. C. intr. 1835.

farinosus H. Hafn. ♂. C. intr. 1836.

incanus; ♂. C. 1836.

Lahlab L. ♂. T.

lignosus L. ♀. ♂. T. ex hort. Paris. 1837.

Soja L. ♂. C.

Zebra Gmel. ♂. C. intr. 1834 ex h. Kil.

Dondia

Epipactis Spr. (*Astrantia* L.) ♀. F. D.; recens adhuc horti incola, mox vero civitatem adeptura: 3 hie-mes jam sub Dio bene per-duravit, a d. 1 Maj. ad 1 Jun. florens.

Doronicum

altaicum Pall. ♀. D. 18 poll. alt.

austriacum Jacq. ♀. D. 1819.

Pardalianches L. ♀. D. intr. 1812.

Dorychnium

latifolium W. ♀. F. ex hort. Paris. 1837.

Draba**Dracocephalum**
(Draknos).

canariense L. ♀. T. ex hort. Paris. 1837.

canescens L. ♂. D. ex hort. Kil. 1835.

ibericum Stev. ex h. Petrop. 1836.

nutans L. ♀. D. 1826.

pellatum L. ♂. D. 1808. 1822, et ex h. Petrop. 1836.

peregrinum L. ♀. D. ex hort. Kil. 1835.

Ruyschiana L. ♀. D. 1826.

sibiricum L. ♀. D. 1827.

speciosum Booth. ♀. D. 1½ uln. alt. intr. 1835. cfr.

Flora 1836. p. 8.

thymiflorum L. ♂. D.

virginianum L. ♀. D. 1826.

Drepania

barbata Desf. v. *Tolpis*.

Dryandra

nivea R. Br. ♀. T. ex h. Hamb. 1834; etiam Kew. 1837 cur. J. Ag.

Drymaria

divaricata Kunth; C. intr. 1837 ex h. Paris. cur. Cl. J. Ag.

Drypis

spinosa ♀. F. ex h. Paris. 1837.

Dufresnea

orientalis; T. ex h. Petropol. 1836. sub hoc nomine.

Dysodium

divaricatum Rich. (*Melanopodium puldosum* Spr. Kunth.) ♂. C. ex h. Paris. 1837. Sesquipedale, flor. radio flav.

Dyssodia

glandulosa Cav. v. *Borbera chrysanthemoides*.

radiata v. *Borbera*.

Eccremocarpus

scaber R. et P. 4. 7. T. D.
ex hh. Bonn. et Hamb.
1836.

Echinaria

capitata Desf. v. *Sesleria echinata*.

Echinophora

scabra; 4. C. F. 15 poll. alt.,
flor. Aug.

tenuifolia L. 4. F. ex h. Petrop. 1836.

trichophylla Sm. T. ex h. Petrop. 1836.

Echinops

bannaticus Roch. Booth. 4. D.
ex h. Paris. 1837.

Ritro L. 4. D. 2½ uln. alt.,
flore Aug.

sphaerocephalus L. 4. D. 3
uln. alt.

strigosus L. ☉. F. 1826; et
nova vice ex hort. Paris.
1837.

Echinosperrum

barbatum Lehm. (*Myosotis* M.
B.) ex h. Petrop. 1836.

Lappula Lehm. (*Myosotis* L. *Myosotis squarrosa* Retz.) ☉. D.
1806 herb. nostr.; 1834.

Echium

cynoglossoides; sub hoc nom.
ex h. Paris. 1837.

italicum L. ♂. D.

violaceum L. (*australe* Cand.)
☉. D. 1815 et ex h. Paris.
1837.

vulgare L. ♂. D.

Eclipta

erecta L. (*Verbesina alba* L.)
☉. C. 1807. herbar.

latifolia L. suppl. ☉. C. 1835.
sesquipedalis flor. magn.
alb. mens. Aug. explic.

prostrata L. Spr. (*Verbesina*
L.) ☉. C. 1806. herbar.
nostr.; etiam ex h. Paris.
1822.

Edwardsia

grandiflora Salisb. (*Sophora tetra-
ptera* Ait.) 7. T. F. intr.
1818; hodie 2½ uln. alt.

microphylla Salisb. (*Sophora
tetraptera* Forst.) 7. F.

Ehrharta

panicea Sm. 4. T. 1810, et ex
h. Paris. 1837.

Elæagnus

hortensis M. B. (*angustifolia*
L.) 7. D. in Paradislyc-
kan 4 uln. alt.

macrophylla Thb. (*argentea* Ag.)
7. D. intr. 1830. hodie 3
uln. alt.

Elæocarpus

cyaneus Sims. (*reticulatus* Sm.
Spr.) 7. T. ex Angl. 1837.

Eleusine

coracana Pers. (*Cynosurus* L.)
☉. D. ex h. Hamb. 1824
et Paris. 1837.

virgata Pers. (*Cynosurus* L. *C.
domingensis* Jacq.) ☉. C.
1811. herbar. nostr.

Elichrysium

argenteum W. olim in horto
cultum, hodie exsul.

fulgidum W. (*Xeranthemum* L.)
4. T. 1810 et ex h. Bonn.
1836.

lucidum Hornem. (*bracteatum*
W.) ☉. ♂. F. D. fl. flav.
1½ uln. alt.

β. fl. albo.

proliferum W., olim horti in-
cola, jam profugum.

Ellisia

Nyctelæa L. ☉. D. ex h. Hafn.
1827 et Paris. 1837.

Elsholtzia

cristata W. (*Hyssopus ocimifo-
lius* Lam.) ☉. D.

Elymus

canadensis L. (*glaucofolius* W.)
4. D. sub illo nomine in
hort. cult. 1812 secund.

herbar. nostrum, sub hoc
ex h. Paris. 1837. missus.
geniculatus W. Hornem. 4. D.
1826.

giganteus Vahl. 4. D. 1812;
et nuper ex h. Paris 1837.

sabulosus M. B. 4. D. 1826.

- sibiricus L. 2. D. ex h. Paris. 1837.
 tener L. 2. D. 1811 herbar. nostr.; iterum 1833.
 villosus Mühl. 2. D. ex hort. Paris. 1837.
 virginicus L. 2. D. 1825; etiam ex h. Paris. 1837.
- Embothrium**
sulignum Andr. v. *Hukea*.
- Entelea**
 arborescens R. Br. Spr. 2. T.
- Epacris**
 juniperina L. (*Lissanthe acerosa* Spr.) 2. T. 23 pollic. alt. intr. 1833. ex hort. Hamb.
 purpurascens R. Br. (*pungens* Sims.) 2. T. 28 poll. alt.
- Ephedra**
monostachya L. 2. seminibus educata per 10 annos in horto sub Dio olim bene viguit, tandem vero evanuit. Cfr. Retz. in act. Holm. 1798. p. 55.
- Epilobium**
 alpestre Schm. ex hort. Crac. 1837.
 coloratum Mühl. 2. D. in h. 1 uln. alt.; flor. parv. rubr. Jul. Aug.
 davuricum Fisch. ☉. 2. D. ex hh. Hafniæ et Crac. 1837.
 hirsutum L. 2. D. 1833, etiam ex h. Crac. 1837.
 montanum L. 2. D.
 nutans Schm. (*hypericifolium* Tausch.) 2. D. 1826, etiam ex h. Hafn. 1837.
 trigonum Hortulan. (var. *tetragoni* L.) 2. D.
- Epimedium**
 alpinum L. 2. D. pedale; flor. Maj. Jun.
- Eqvisetum**
 arvense L. 2. D.
 hyemale L. 2. D.
 umbrosum W. 2. D. 1826.
- Eranthemum**
 parviflorum R. Br. (*Justicia* Ort. Vahl. Hornem.) 2. T. 1810. 1824 et ex h. Kil. 1835.
- Erica** (Ljung).
 arborca L. 2. F. 1819.
 gracilis Wendl. 2. T. 16 poll. alt. intr. 1834 ex h. Hamb.
 herbacea L. 2. F. 1812.
 ignescens Andr. 2. T.
 mediterranea L. 2. T. 1808. herbar. nostr.
 Patersonia Andr. (*abietina* L.) 2. T.
 Petiverii L. 2. T.
 vagans L. 2. T. floruit 1835. 12 poll. alt.
 viridipurpurea Horn. 2. T. species.
- Erigeron**
 armeriaefolium; ex h. Paris. 1837.
 caucasicum Stev. 2. D. ex h. Paris 1837.
 delphinifolium W. 2. T. ex h. Paris. 1837.
 elongatum Booth. 2. D. F. ex h. Petrop. 1836.
 heterophyllum W. (*Aster annuus* L.) ☉. D.
- Erinus**
 alpinus L. 2. D. ex h. Hafn. 1827 et Paris. 1837.
- Erodium**
 arborescens W. 2. T. in hort. 1½ uln. alt.
 chamædryoides Herit. Retz. 2. F. 1806 in h. cult. hodie deest.
 obium W. (*Geranium* L.) ☉. D. ex h. Paris. 1837.

- cicutarium* Sm. (*Geran.* L.) ☉.
D. spontan.
geifolium; F. ex h. Par. 1837.
(an *hymenodes*?).
gruinum Ait. (*Geran.* L.) ☉.
D. 1808 herbar.
hymenodes Herit. 2. F.
malacoides W. (*Geran.* L.) ☉.
D. 1810 et ex h. Paris.
1837.
maritimum Sm. 2. D. ex h.
Paris. 1837.
romanum Ait. (*Geran.* L.) 2.
D. ex h. Bonn. 1836.
- Ervum**
dispermum Roxb. ☉. D. 1835.
Ervilia L. (*Vicia* W.) ☉. D.
1821—1837. sesquipedal.
hirsutum L. ☉. D.
- Eryngium**
alpinum L. 2. D. 32 pollic.
alt. intr. 1833 ex h. Gr.
aquaticum L. 2. D. 1 uln. alt.
in hort.
asperifolium Spr. 2. D. ex h.
Paris. 1837.
azureum Hort. 2. D. 1835.
27 poll. alt.
bromelifolium Spr.; ex h. Pa-
ris. 1837.
corniculatum Lam. 2. D. F.
ex h. Paris. 1837.
dichotomum Desf. 2. D. 1835
etiam sub nom. *cæruleum*
ex h. Paris. 1837 miss.
planum L. 2. D. 1826—1837.
pusillum L. 2. F. ex h. Paris.
1837.
tricuspidatum L. 2. D. 1½
uln. alt.
- Erysimum**
Alliaria L. ♂. D.
Barbarea L. v. *Barbarea vut-*
garis.
Cheiranthus Spr. Pers. (*murale*
Desf. Hort. Paris.) ☉. D.
semin. ex h. Paris. 1837.
collinum Andr. (*Cheiranthus*
M. B.) 2. D. 1835, etiam
ex h. Petrop. 1836.
helveticum Cand. (*Cheiranthus*
Jacqv.) ♂. D. 1812. herb.
nostr.
- officinale* L. v. *Sisymbrium*.
virgatum Roth. ♂. D. ex h.
Paris. 1837.
- Escallonia**
floribunda Humb. ♂. T.
rubra Pers. ♂. T. 33 poll. alt.
intr. 1833. ex h. Hamb.
β. ex Angl. 1837.
- Eschscholtzia**
californica Chamiss. ♂. 2. D.
intr. 1832. flor. citrin.
crocea Benth. 2. D. intr. 1836.
- Ethulia**
conyzoides L. ☉. C. intr. 1807;
ex hh. Hafn. et Kil. 1836.
- Eucalyptus**
species ♂. T. ex Angl. 1837.
- Euclea**
racemosa L. 2. T.
- Eucomis**
punctata Herit. (*Ornithogalum*
Thb.) 2. T.
- Eugenia**
australis Wendl. v. *Myrtus*.
trinervia Vahl. v. *Myrtus Vahl.*
- Eupatorium**
ageratoides L. 2. D. 1812.
1826.
altissimum L. 2. D. ex hort.
Paris. 1837.
coelestinum L. (Coelestina cæ-
rulea Cassin.) 2. D. F.
1807. Herb. nostr.
maculatum L. 2. D. ex h. Pa-
ris 1837.
verticillatum W. 2. D. 1812.
1826.
- Euphorbia**
begoniæfolia; sub hoc nom ex
h. Paris. 1837.
Caput Medusæ L. ♂. T.
ceratocarpa Tenor. ♂. 2. T.
F. 1835. 3-pedalis; flor.
Aug.
Cyparissias L. 2. D.
Esula L. 2. D.
exigua L. ☉. D. olim in h.
cult. pedalem attingit lon-

gitudinem; spontanea palmaris.

helioscopia L. ☉. D. spont.

Lathyris L. ♂. D.

orientalis L. ♀. D. ex h. Petropol. 1836.

pallida W. (*lucida* Kit.?) ♀. D. ex h. Paris. 1837.

palustris L. ♀. D.

Peplus L. ☉. D. spont.

procera M. B. ♀. D. 1822. 1826.

prunifolia Jacq. ☉. C. olim (1810) in h. culta, hodie ibi non exstat.

pterococca Brot. ☉.; ex h. Paris. 1837.

salicifolia Host. ♀. D. 1819 in h. sub Dio culta, ibi diutius non obvia.

splendens H. Hamb. ♀. C. de H. Hamb. 1837 cmta, et intr.

Szovitsii; T. ex h. Petr. 1836.

veneta W. ♀. F. 1820 in hort. culta.

verrucosa L. ♂. ♀. D. F. 1½ uln. alt., flor. Jun. species.

Euphrasia

officinalis. L. ☉. D.

Eutoca

wrangeliiana F. et M., h. Lips.

☉. T. ex h. Hafn. 1837.

Evolvulus

alsinoides L. ☉. C.

Evonymus (Benved).

europaeus L. ♀. D. in Paradislyckan.

latifolius Scop. ♀. D. 8½ uln. alt.

nanus M. B. ♀. D. ¾ uln. alt. in nov. arbor.

Tobira Thb. v. *Pittosporum*.

verrucosus Ait. ♀. D. 4. uln. alt.

Exacum

viscosum Sm. ♀. ♀. T. 1810.

Fabricia

laevigata Gärtn. ♀. T.

Fagus (Bok).

castanea L. v. *Castanea vesca*.

sylvatica L. ♀. D. Paradislyckan.

β. atropurpurea ♀. D. 8

uln. alt., in nov. arboretum intr. 1826.

Farsetia

delloidea R. Br. v. *Aubrietia*.

incana R. Br. v. *Alyssum*.

verrucosa; sub hoc nom. ex h. Paris. 1837. J. Ag.

Fedia

angulata; ex h. Crac. 1837.

Auricula Cand. (*Valerianella* H.

Crac.) ex h. Crac. 1837.

congesta (*Valerianella* H. Crac.)

☉. D. ex h. Crac. 1837.

cornucopia Vahl. ☉. D. pedalis, flor. rubicund. Jun.

Aug.

coronata Vahl. (*Valerianella campanulata* Biv.) ☉. D.

discoidea Vahl. (*Valeriana* Brot.)

☉. D.

echinata Vahl. ☉. D. ex hort.

Paris. 1837.

eriocarpa Desv. Hornem. ☉.

D. ex h. Hafn. 1827 et

Crac. 1837.

hamata (*Valerianella* H. Lips.)

☉. D. ex h. Crac. 1837.

pumila Vahl. (*Valerianella* H.

Cr.) ☉. D. ex hort. Crac.

1837.

serioides H. Crac.; intr. 1837.

ex h. Crac. miss.

Felicia

tenella Nees. ☉. (♀. h. Gr.)

C. ex h. Kil. 1837.

Ferula

communis L. ♀. F. ex h. Pa-

ris. 1837.

Ferulago L. ♀. F. ex h. Hafn.

1827 et 1837.

- glauca* L. 2. F. 1826. et nova vice ex h. Crac. 1837.
nodiflora L. 2. D. 1826. et iterum ex h. Hafn. 1837.
Opopanax Spr. (*Pastinaca* L.) 2. D. ex h. Crac. 1837.
orientalis Tournef. 2. F. intr. 1835 ex h. Hafn.
sibirica Spr. (*Peucedanum* W.) 2. D. ex h. Crac. 1837.
tingitana Herm. 2. F. ex h. Paris. 1837.
- Ferraria**
Pavonia L. v. *Tigridia*.
- Festuca** (Svingel).
Alopecurus Schousb. ☉. D. 1812. herb. nostr.
diandra Mx. (*Corynepus arundinaceus* Lea. Spr.) 2. D. 1833; et ex h. Par. 1837. cnr. Cl. J. Ag.
duriuscula L. 2. D. 1826. Mörcck.
glauca Lam. 2. D. 1826.
heterophylla Juss. 2. D. 1810. herbar.
laxa Host, 2. D. 1826.
ovina L. 2. D. spont.
rubra L. 2. D.
- Ficus** (Fikon).
Carica L. 2. F.
cordata Thb. 2. C.
retusa L. 2. C.
stipulata Thb. (*scandens* Lam.) 2. C.
- Filago**
arvensis L. ☉. 2. D.
- Flaveria**
Contrayerva Pers. (*Milleria* Cav.) ☉. C. ex h. Paris. 1837. 12-poll. alt.
repanda Lag. (*Nauenburgia 3 nervata* W. *Brotera Contrayerva* Spr.) ☉. C.
- Fontanesia**
phillyreoides Lahill. 2. T. 1823. herb. nostr.
- Forskahlea**
angustifolia Retz. ☉. C. 1820. herbar. nostr.
- Fragaria** (Smultron).
collina Ehrh. 2. D.
elatio Ehrh. (*Fr. vesca* 2. Wabl.) 2. D.
indica Andr. 2. D. F. 1826, et ex h. Paris. 1837.
vesca L. 2. D.
virginiana Ehrh. 2. D. 1827-1837.
- Franchon** (*Francoa*).
appendiculata Cav. 2. 2. C. ex h. Paris. 1837.
sonchifolia Spr. 2. 2. C. ex h. Hafn. 1835.
- Fraxinus** (Ask).
excelsior L. 2. D. Paradiallykan.
juglandifolia Lam. (*Caroliniana* du Roi.) 2. D. 21 uin. alt.
lentiscifolia Desf. 2. D. 1826 in hort. cult. hodie exsul.
Ornus L. 2. iteratis vicibus, sed sine successu sub dio plantata in hort. nostr.
rotundifolia Ait. 2. in horto aliquando plantata, sed absque successu, frigoris valde impatiens.
sumbucifolia Lam. 2. 1826 in h. sub dio plantata, sed non constans.
- Fritillaria**
imperialis L. 2. D.
latifolia W. 2. D. 1826; hodie horti profuga.
Meleagris L. 2. D.
pyrenaica L. 2. D.
- Fuchsia**
Adolphina H. Hamb. 2. T.
baccillaris Booth. (*globosa*) 2. T. 12 poll. alt. intr. 1836. ex h. Hamb.
coccinea L. 2. T. F.
conica Booth. 2. T. ex hort. Hamb. 1836.

- gracilis* Lindl. f. T. de hort.
Hamb. 1837 emta.
microphylla Booth. Wilbr. f.
T. 10 pollic. usque ad 1
uln. alt. intr. 1833 ex h.
Hafn. (Cfr. etiam *Flora*
1836. p. 4).
mutabilis Booth. f. T. intr.
1837, de h. Hamb. emta..
Smithii grandiflora Booth. f.
T. emta de h. Hamb. 1837.
Thomsoniana Booth. f. T. 12
ad 18 poll. alt. intr. 1833
ex h. Hamb.
thymifolia Humb. Booth. f. T.
14-16 poll. alt. ex hort.
Hamb. 1836. cur. Cel.
Sönnnerberg.
virgata Booth. f. T. 1 uln. alt.
1836.
- Fumaria** (Jordrök).
capreolata L. \odot . D.
officinalis L. \odot . D. spont.
spicata L. \odot . D.
vesicaria L. v. *Cysticapnos afri-*
cana.
- Funkia**
ovata Spr. (*Hemerocallis caru-*
lea Andr. Hem. japonica β .
W.). f. D.
- Galanthus**
nivalis L. f. D. (Vinterlilja).
- Galardia** (*Gaillardia*).
aristata Pursh. f. D. ex hort.
Londin. 1834 et Bonn.
1836. $1\frac{1}{2}$ uln. alt.; flor.
Aug.
bicolor Lam. f. D. ex hort.
Bonn. 1836.
rustica Cass. f. D. ex h. Cr.
1836.
- Galega**
officinalis L. (*persica* Pers.) f.
D. Paradislyckan.
orientalis Lam. f. D. bipeda-
lis et ultra.
- Galeobdolon**
luteum Sm. (*Galeopsis Galeobdol-*
L.) f. D.
- Galeopsis**
Tetrahit L. \odot . D. spont.
versicolor Curt. \odot . D. spont.
- Galinsoga**
parviflora W. (*Siegesbeckia*) \odot .
D. 1808-1820. herb. nosr.
trilobata Cav. \odot . D. 1827.
- Galium**
Aparine L. \odot . D. spont.
austriacum Jacq. Horn. f.
D. 1826.
articulatum Lam. (*rubroides*
Spr.?) \odot . D. ex h. Par.
1837.
Bocconi Hornem. All. f. D.
1815 et ex h. Par. 1837.
boreale L. f. D. spont.
caucasicum Lag. f. D. ex h.
Paris. 1837.
cinereum All. f. D. 1826.
cruciatum Sm. Spr. (*Valantia*
L.) f. D. 6-8-poll. flor.
viridi flav. Maj.
erectum Huds. (*lucidum* All.)
 f. D. 1826.
glaucum. f. D. sub hoc nom.
ex h. Paris. 1837.
humifusum M. B. (*Asperula*
Spr.) f. D. ex h. Hafn.
1837.
læve f. D. sub hoc nom. ex
h. Paris. 1837.
linifolium Lam. f. D. 1826.
parisiense L. \odot . D. 1815 et
ex h. Paris. 1837.
provinciale; D. 1835. 1 uln.
et ultra alt. flor. alba.
sylvaticum L. f. D. 1822.
tyrolense W. f. D. 1815.
1826, et nova vice ex h.
Paris. 1837.
uliginosum L. f. D. spont.
verum L. f. D. spont.
species.
- Garidella**
Nigellastrum L. \odot . D.
- Garrya**
elliptica Booth. f. T. ex Angl.
1837. cur. Cl. J. Ag.

Gaultheria

procumbens L. f. F. intr. 1836.

Gaura

biennis L. f. F. ex h. Paris. 1837.

mutabilis Cav. f. T. 1810 et ex h. Hafn. 1837.

tripetala Cav. f. f. T. ex h. Hafn. 1837.

Genista

athnensis Spr. v. *Spartium*.

canariensis L. f. F. 1810. 1819.

candicans L. v. *Cytisus*.

junceae Spr. v. *Spartium*.

ovata Kit. f. D. 1826.

scoparia Lam. v. *Spartium*.

sibirica L. f. D. flor. flav.

tinctoria L. f. D. 2 uln. alt. flor. flav. Aug. explic.

Gentiana

asclepiadea L. f. D. ex hort. Kil. 1835.

cruciata L. f. D.

lutea L. f. F. 1810.

Georgina

variabilis W. (*Dahlia* Cav.) f. D.

D. flor. plenitud. et color. nimis var.

Geranium

aconitifolium Herit. f. T. 1830—36.

alchemilloides L. v. *Pelargonium*.

anemonifolium Herit. f. T. 1822—37. olim (1807)

sub nom. *G. Palmatum* Cav. in hort. cult.

f. fl. albo.

bohemicum L. f. D. 1807. herb. nostr.

capitatum L. v. *Pelargonium*.

carolinianum L. f. D. 1824. et ex h. Paris. 1837.

collinum Steph. f. D. 1822.

columbinum L. f. D. spont. in Paradislyckan.

curculatum L. v. *Pelargonium*.

dissectum L. f. D. spont.

ibericum Cav. f. D. bipedale; flor. Jun. Jul.

Londesii Fisch. f. D. 1826. 1835.

lucidum L. f. f. D. spont.

macrorrhizum L. f. D. 14-16 poll. alt. flor. purp.

maculatum L. f. D. 1826.

molle L. f. D. spont.

nodosum L. f. D. 1-pedale; flor. Jun. — Aug.

odoratissimum L. v. *Pelargonium*.

petatum L. v. *Pelargon*.

phæum L. (*bivium* Herit.) f. D. 1-pedale.

pratense L. f. D.

f. fl. alb.

pusillum L. Mantiss. 2 (*rotundifolium* L. Svec. Wahl.)

f. D. spont.

pyrenaicum L. f. f. D. in h. jam spont. dissem.

reflexum L. f. D. 12-15 poll. alt., flor. Maj. — Aug.

Robertianum L. f. D. spont.

sanguineum L. f. D.

sibiricum L. f. D.

silvaticum L. f. D.

striatum L. f. D.

viscidulum Fries f. D. 1835 sporadic.

Vlassovianum Fisch. f. D.

zonale L. v. *Pelargonium*. species.

Geropogon

australis Spr. (*G. glaber* et *hirsutus* L.) f. D. ex hort.

Kil. 1835.

Gesnera

Douglasii H. Hamb. f. C. ex Angl. 1837. cur. J. Ag.

Geum

album Gmel. (*canadense* Jacqv.) f. D. 1826, et ex h. Gr.

1837.

atlanticum Desf. f. D. 1826.

coccineum Sm. 2. ex h. Bonn. 1836.

hirsutum; 2. sub hoc nom. ex h. Paris. 1837.

japonicum Thb. 2. D. 1835. 18-22 poll. alt. flor. flav.

intermedium Hornem. 2. D. *macrophyllum* W. 2. D. ex h. Paris. 1837.

nutans Lam. 2. ex h. Paris. 1837.

rivale L. 2. D. flor. simpl. et prolifer. spont.

β. fl. luteo Fr. Fl. Scan. in *Paradislyckan*.

rubifolium H. Kil. 2. ex h. Gryph. 1837. miss.

strictum Ait. 2. D. 1835. 3-pedale; fl. flav. Jun. explic.

urbanum L. 2. D. spont.

virginianum L. 2. D. ex h. Paris. 1837.

Gilia

achilleæfolia Benth. ☉. D. ex h. Petr. 1836. sesquipedal. flor. cærul. Jul. explic.

capitata Hook. ☉. D. 1835. 3-pedal. flor. pallide cærul. Aug. explic.

chilensis, ☉. D. ex h. Bonn. 1836.

eryngioides Trev. ☉. D. ex h. Kil. 1836.

inconspicua Hook. ☉. ♂. D. ex h. Crac. 1837.

laciniata R. et P. (*Cantua breviflora* Juss.) ☉. D. 1834 et ex h. Paris. 1837. 8-10 poll. alt.

tenuifolia ☉. D. ex h. Paris.

tricolor Benth. ☉. D. ex h. Bonn. 1836.

β. fl. albo.

Ginkgo

biloba L. v. *Salisburya*.

Gladiolus

blandus Ait. 2. D. 27 poll. alt. flor. purp., Jul.

byzantinus Mill. 2. T. *Cardinalis* Curt. 2. T.

communis L. 2. D. *psittacinus* Booth. 2. T.

refractus Jacqy. (*Tritonia* Ker.)

2. T. ex hort. Parisiens. 1837.

Glaucium

(Horn-Wallmo Hortul).

flavum Crantz. ♂. D. ex hort. Kil. 1835.

luteum Scop. (*Chelidon. Glaucium* L.) ♂. 2. D.

β. *fulvum* Sm.

phoeniceum Sm. (*Chelid. corniculatum* L.) ♂. ☉. D. ex h. Kil. 1835.

Glechoma (Jordrefva).

hederaceum L. 2. D. spont.

Gleditschia

caspica Desf. ♀. F. ex h. Petropol. 1836.

ferox Desf. (*macrantha* W.) ♀. D. seminib. educata; jam $\frac{1}{2}$ uln. alt.

horrida W. ♀. F.

monosperma Walt. ♀. F.

sinensis ♀. F. ex h. Hamb. 1836.

triacantha L. ♀. D. per 10 et quod excurrit annos in horto sub diu viguit; jam 6 uln. alt.

Globba

nutans W. v. *Alpinia*.

Gloriosa

superba L. 2. C.

Gloxinia

caulescens; 2. C. ex h. Petr. 1836.

maculata Herit. (*Martynia perennis* L.) 2. C. in Cald. nostr. culta fuit; sed nuper emortua.

speciosa Ker. 2. C. 1835 spitham., flor. cærul.

Glycine

coccinea Curt. (*Kennedia prostrata* R. Br.) ♀. C. ex h. Crac. 1837.

tomentosa L. 2. F. ex h. Crac. 1837.

species 2. C.

Glycosmis

citrifolia; C. ex h. Paris. 1837.

Glycyrrhiza

(Lakritsrot).

echinata L. 2. D. 1826.

foetida Desf. 2. F. ex h. Paris. 1837.

glabra L. 2. D. ex hort. Kil. 1835.

glandulifera Kit. 2. D. intr. 1835; deinde ex h. Paris. 1837.

Gnaphalium

apiculatum Labill. 2. T. ex h. Crac. 1837.

arenarium L. 2. D.

chilense Spr. 2. T. ex h. Paris. 1837.

coarctatum W. (*spicatum* Lam.)
⊙. D. ex h. Paris. 1837.

congestum Lam. 2. T.

dioicum L. 2. D.

falcatum Lam. 2. ex h. Crac. 1837.

foetidum L. 3. (2) T. 1834
et ex h. Crac. 1837.

leyserioides Desf. (*Longohampia capillifolia* W. Spr. *Leysera discoidea* Spr. *Pectis discoidea* Horn.) ⊙. D. ex h. Paris. 1837.

margaritaceum L. 2. D. 1-2
pedal. flor. Aug.

maritimum L. 2. T. ex hort. Crac. 1837.

obtusifolium L. in hort. cult. 1806. herb. nostr.

orientale L. 2. F. 1810.

resedæfolium Trev. ⊙. C. ex h. Paris. 1837.

rosmarinifolium Sieb. 2. T. ex h. Crac. 1837.

spathulatum; ⊙. D. sub hoc nom. ex h. Paris. 1837.

Stoechas L. 2. F. 1810 et ex h. Paris. 1837.

lide flav. intr. 1834 ex h. Hamb.

Godetia v. Oenothera.**Gomphocarpus**

arborescens R. Br. v. *Asclepias*.

Gomphrena

decumbens Jacq. ⊙. C. 1810
et ex h. Hafn. 1837.

globosa L. ⊙. C. fl. purpur.,
carneo, albo.

interrupta ⊙. C. 1810 et ex h. Paris. 1837.

Gonolobus

rostratus R. Br. (*Cynanchum*
Vahl.) 2. C.

Gossypium

herbaceum L. ⊙. C. ex hort.
Hafn. 1835.

Grahamia

aromatica Spr. (*Gramia* Hook.
Cephalophora Schrd.) ⊙. T.
intr. 1834. 12 — 20 poll.
alt.

Grangea

maderasputana W. v. *Cotula*.
minima W. v. *Cotula*.

Gratiola (Jordgalla).

lucida Vahl. (*Morgania* Spr.
Capraria crustacea L.) ⊙.
C. ex h. Hafn. 1835.

officinalis L. 2. D. sesquiped.
flor. Jul. — Sept.

serrata Roxb. v. *Bonnaya brachiata*.

Grevillea

buxifolia R. Br. 2. C. T. 20
poll. alt. intr. 1834 ex h.
Hamb.

juniperina R., Br. 2. T. intr.
1834 ex h. Hamb.

linifolia; 2. T.

rosmarinifolia H. Hamb. 2. T.
ex hort. Kew. intr. 1837.
cur. Cl. J. Ag.

Gnidia

sericea Spr. (*Passerina* L.) 2.
T. 33 poll. alt. flor. pal-

Grindelia
incisa Spr. (Adm. Fisch.) 2.
 T. F. ex h. Crac. 1837. 1
 uln. alt.; flor. Aug.; pe-
 tal. albo-coral.

Gundelia
Tournefortii L. 2. F. ex h.
 Petrop. 1836.

Gymnocladus
canadensis Lam. (*Guilandina di-*
caica L.) 2. D. 2½ uln. alt.

Gypsophila
acutifolia Fisch. 2. F. 1826.
arenaria Kit. (*G. fastigiata* var.
 Spr.) 2. D. 1819. 1822.
 herb. nostr.

dichotoma Bess. 2. D. ex hort.
 Paris. 1837.

elegans M. B. (2). D. ex hort.
 Paris. 1837. 1-pedal. flor.
 albidis rubro-striat. Jul.
 explic.

glomerata Pall. 2. F. 1826.
paniculata L. 2. D. ex hort.
 Paris. 1837.

sabulosa Stev. 2. D. ex hort.
 Paris. 1837.

saxifraga L. (*Dianthus* L. Sp.
 1.) 2. D. 1808 herbar.
 nostr.

scorzoneræfolia, sub hoc nom.
 ex h. Paris. 1837.

Vaccaria Sm. (*Süpanaria* L.)
 (2). D. bipedal. flor. parv.
 roseis, Jul. explic.

Hablizia

tamnoides M. B. 2. F. ex h.
 Hafn. 1836.

Hæmanthus

coccineus L. 2. C.
punicus Jacq. 2. C.

Hakea

angustifolia 2. T. ex Anglia
 1837. cur. Cl. J. Ag.

dentata 2. T.

elliptica R. Br. 2. T. 1834 et
 ex Angl. 1837.

florida R. Br. 2. T.

lanigera Booth. 2. T. ex Angl.
 1837.

microcarpa R. Br. 2. T. intr.
 1834.

microphylla Booth. 2. T. 18
 poll. alt. flor. parv. albid.;
 intr. 1834 ex h. Hamb.

ruscifolia Labill. 2. T. ex h.
 Kewens. 1837.

saligna R. Br. (*Embothrium*
 Andr.) 2. T.

Halesia

tetraptera L. 2. D. in hort. olim
 sub Dio, sed sine succes-
 su plantata. Cfr. Retz. in
 Act. Holm. 1798.

Halimodendron

argenteum Cand. v. *Robinia Ha-*
lodendron.

Halleria

lucida L. 2. T. *Flora* 1836.
 pag. 4

Haloragis

Cercodia Ait. 2. T. 1810
 et ex h. Kil. 1835.

Hamamelis

virginica L. 2. D.

Harrachia

speciosa Jacq. (*Justicia infun-*
dibuliformis L.) 2. C. —
 etiam ex h. Crac. 1837.

Hasselqvistia

cordata L. (2). D. ex h. Paris.
 1837.

Haworthia

concinna Haw. v. *Aloe viscosa*.

Hebenstreitia

tenuifolia 2. T. 1833 et sub
 hoc nom. ex h. Paris. 1837.

Hedera (Murgrön).

Helix L. 2. D.

hibernica H. Hamb. $\frac{1}{2}$ D. 1836.
quinquefolia L. v. *Vitis hederacea*.

Hedychium

coronarum Kön. $\frac{1}{4}$ C.
Gardnerianum? Wall. $\frac{1}{4}$ C.

Hedyotis

biflora H. Hafn. v. *Oldenlandia*.
diffusa W. (*Oldenlandia* Roxb.
 Horn.) $\frac{1}{4}$ C. ex h. Hafn.
 1837.

Hedypnois

rhagadioloides L. (*Hyoseris cre-*
tica Cav.) \odot . D. 1810.
 1815 et ex h. Paris 1836.

Hedysarum

alpinum L. $\frac{1}{4}$ D. F. 1826.
canadense L. $\frac{1}{4}$ D. 1809. herb.
 nostr. et nova vice ex h.
 Paris. 1837.

canescens L. $\frac{1}{4}$ F. 1811 herb.
 nostr.

Caput Galli L. v. *Onobrychis*.
carpathicum W. v. *Onobrychis*
petraea.

coronarum L. $\frac{1}{4}$ D. ex hort.
 Bonn. 1836.

Crista Galli L. v. *Onobrychis*.
elongatum; $\frac{1}{4}$ sub hoc nom. ex
 h. Petrop. 1836.

flexuosum L. \odot . T.
marilandicum L. $\frac{1}{4}$ F. ex h.
 Hafn. 1836.

muricatum Jacq. \odot . D. 1815
 herb. nostr.

obscurum L. $\frac{1}{4}$ D. ex h. Petr.
 1834 et 1836.

Onobrychis L. v. *Onobrychis sa-*
tiva Lam.

petraeum M. B. v. *Onobrychis*.
saxatile L. v. *Onobrychis*.

Helenium

autumnale L. $\frac{1}{4}$ D. 1808.
 herb. nostr.

mexicanum Humb. \odot . D. ex
 h. Paris. 1837.

quadridentatum Labill. (*Rud-*
beckia alata Jacq.) $\frac{1}{4}$ \odot .
 T. 1821.

Helianthemum Tournf.

(*Cisti species* L.).

egyptiacum Mill. \odot . D. ex h.
 Paris. 1837.

ledifolium W. (*Cistus* L.) \odot .
 D. 1834. 1835.

mutabile Pers. $\frac{1}{2}$ F. 1822 et
 ex h. Paris. 1837.

procumbens Dun. $\frac{1}{2}$ F. 1834
 et ex h. Petrop. 1836.

salicifolium Pers. \odot . D. ex h.
 Paris. 1837.

thymifolium Pers. $\frac{1}{2}$ F. ex h.
 Paris. 1837.

vulgare Gärt. (*Cistus Helian-*
them. L. *Cist. grandiflorus*
 Scop.) $\frac{1}{2}$ D.

Helianthus (Solros).

annuus L. \odot . D.

decapetalus L. $\frac{1}{4}$ D. 1826.

indicus L. (*annuus* var. Spr.)
 \odot . D.

lenticularis. Lindl. \odot . D. ex
 h. Hafn. 1836.

multiflorus L. $\frac{1}{4}$ D. 1815 herb.
 nostr.

pubescens Vahl. $\frac{1}{4}$ D. ex h.
 Paris. 1837.

rigidus; sub hoc nom. ex h.
 Paris. 1837.

tuberosus L. $\frac{1}{4}$ D.

Heliophila

amplexicaulis L. \odot . D. ex h.
 Paris. 1837.

pendula W. (*pinnata* Vekt.)
 \odot . D. ex h. Paris. 1837.

Heliopsis

laevis Pers. (*Bupthalm. helian-*
thoides L.) $\frac{1}{4}$ D. ex hort.
 Paris. 1837.

Heliotropium

europaeum L. \odot . D.

indicum L. v. *Tiaridium*.

parviflorum L. \odot . C. 1812.
 herbar. nostr.

peruvianum L. $\frac{1}{2}$ T. 15—16
 poll. alt.

β . fl. alb.

Helleborus (Prustrot).

foetidus L. 2. D. in Paradis-lyckan.

hyemalis L. 2. D. jam horti nostri indigena.

niger L. 2. D. a frigore hie-
mum non læditur.odorus Kit 2. D. F. ex hort.
Paris. 1837.viridis L. 2. D. coelo horti
nostri familiarissimus.**Helminthia**echioides W. (*Pieris* L.) ☉. D.**Hemerocallis**coerulea β. *japonica* W. v. *Fun-*
lia ovata.

flava L. 2. D. 1826.

fulva L. 2. D. 1826.

graminea Horn. 2. D. 1826.

Liliastrum L. (*Czackia Liliastr.*
Andrz.) 2. D. 1826.**Hemimeris**urticefolia W. (*Celsia* Curt.)p. T. 1810 et ex h. Crac.
1837.

species, T,

Heracleumamplifolium Lapeyr. *pyrenai-*
cum Cuss. *speciosum* Hoffm.)

2. D. 1826; et nov. vice

ex h. Paris. 1837; 5 uln.
alt.gummiferum W. ♂. 2. F. ex
h. Paris. 1837.persicum; sub hoc nom. ex h.
Paris. 1837.

sibiricum L. ♂. D. 1826.

Sphondylium L. ♂. 2. D.

β. sub nom. *Wendia Cho-*
rodanum Hoffm. ex h. Pe-
tropol. miss. 1836.

species.

Hermanniaangularis Jacq. p. T. 1810
et ex h. Paris. 1836.denudata L. p. T. 1810 et ex
h. Crac. 1837.

hirsuta Schrad. p. T. 1804

herbar. nostr.

hyssopifolia L. p. T. 1810.
1814—1837.**Hermesia***Banksiaefolia* v. *Callicoma*.**Herniaria**

glabra L. 2. D.

hirsuta L. 2. D.

Herpestis (*Herpestes*).chilensis H. Crac. T. sub hoc
nom. ex h. Crac. 1837.crysanthia H. Hafn. T. ex h.
Hafn. 1836.**Hesperis** (Nattviol).*africana* L. v. *Malcolmia**chia* Lam. v. *Malcolmia*.*luxa* Lam. v. *Chorisporea tenella*.

elata Hornem. ♂. D. 1826.

matronalis L. ♂. D.

tristis L. ♂. D.

verna L. (*Arabis* R. Br.) ♂. ☉.
D. 1820.**Heterospermum**pinnatum Cav. ☉. D. 1811,
et iterum ex h. Par. 1837.**Heterotheca**inuloides Cass. H. Lips.; ex h.
Paris. 1837.**Heuchera**americana L. 2. D. 2-3-ped.
flor. Jun. — Aug.lucida H. Hamb. 2. F. ex h.
Paris. 1837.macrophylla H. Hamb. 2. F.
ex h. Paris. 1837.villosa Mx. 2. F. ex h. Paris.
1837.

species.

Hibbertiadentata R. Br. p. T. 3-pedal.
flor. pallid. flav.velubilis Andr. (*Dillenia volubi-*
lis Vent. *scandens* W.) p.
T. 1836.**Hibiscus**

Abelmoschus L. p. C.

cannabinus L. ☉. C. 1835.

- e seminibus maturatis
 1834.
esculentus L. ☉. C. 1834 et
 ex h. Hafn. 1835.
gossypinus Thb. ☿. C. 1834.
 1835.
Humboldtii Mart. ☉. D. ex h.
 Gryph. 1835. 6-8-pollic.
 flor. albesc.
liliiiflorus Cav. ☿. C.
Muluviuscus L. v. *Achunia*.
Manihot L. ☿. 2. C. 1810 et
 ex h. Hafn. 1836.
β. palmatus Cav. Horn.
mutabilis L. ☿. C.
pedunculatus Thb. ☿. C. 18
 poll. alt. flor. ros
populneus L. (*Thespesia* Corr.)
 ☿. C.
Richardsonii Sweet. ☿. intr.
 1834. ex h. Kil.
Rosa sinensis L. ☿. C.
speciosus Ait. 2. C. ex h. Kil.
 1836.
splendens H. Hamb. C. intr.
 1834.
surattensis L. ☿. ☉. C. 1834-
 1837.
syriacus L. ☿. F. D. 1819-
 1837.
tortuosus H. Hamb. ☿. C.
Trionum L. ☉. D.

Hieracium

- aurantiacum* L. 2. D.
aureum Scop. (*Leontodon* L.) 2.
 F. 1822.
Auricula L. 2. D.
boreale Fries. (*sabaudum* L.
 Retz.) D. 2., 1837 ex h.
 Crac.
collinum Gochn. (*stoloniferum*
 Reichb.) 2. ex h. Gryph.
 1837.
cydonifolium Vill. 2. D. 1822.
cymosum L. 2. D. 1812-24.
 herbar. nostr.
echioides Lumnitz. 2. D. 1810
 et ex h. Paris. 1837.
foliosum Kit. 2. D. ex hort.
 Paris. 1837.

- fruticosum* W. ☿. F. 1810 et
 ex h. Paris. 1837.
glaucum All. 2. D. 1810 et
 ex h. Gryph. 1837.
grandiflorum Horn. 2. D. 1810
 et ex h. Paris. 1837.
Jacqvini Vill. 2. D. F. ex h.
 Paris. 1837.
lavigatum W. 2. D. 1811 her-
 bar. nostr.
lanatum Kit. 2. F. ex h. Pa-
 ris. 1837.
lapsanoides Gouan. 2. D. ex
 h. Crac. 1837.
Lawsonii Vill. 2. F. 1823;
 etiam ex h. Paris. 1837.
Ledebourianum Tausch. 2. ex
 h. Gryph. 1837.
nigrescens W. 2. D. 1810.
 1815 et ex h. Paris. 1837.
nudicaule Mx. 2. D. ex hort.
 Gryph. 1837.
Pilosella L. 2. D. spont.
polymorphum H. Crac.; com-
 munic. 1837.
præmorsum L. 2. D.
pratense Tausch. 2. D. ex h.
 Gryph. 1837.
ramosum Kit. 2. D. ex hort.
 Paris. 1837 (an *glaucum*
 All.?)
rupestre All. 2. D. 1822.
 1833.
speciosissimum Hornem. 2. D.
 ex h. Gryph. 1837.
speciosum Hornem. Wilbr. 2.
 D. ex h. Crac. 1835,
Sternbergii Hopp. 2.; ex h.
 Gryph. 1837.
umbellatum L. 2. D. etiam ex
 h. Gryph. 1837.
β. coronopifolium Bernh.
villosum L. 2. D. 1 uln. alt.

Hierochloa

- borealis* Schrad. et Fr. (*Hol-
 cus odoratus* L.) 2. D.

Glycosmis

citrifolia; C. ex h. Paris. 1837.

Glycyrrhiza

(Lakritsrot).

echinata L. 2. D. 1826.

foetida Desf. 2. F. ex h. Paris. 1837.

glabra L. 2. D. ex hort. Kil. 1835.

glandulifera Kit. 2. D. intr. 1835; deinde ex h. Paris. 1837.

Gnaphalium

apiculatum Labill. 2. T. ex h. Crac. 1837.

arenarium L. 2. D.

chilense Spr. 2. T. ex h. Paris. 1837.

coarctatum W. (*spitatum* Lam.) 2. D. ex h. Paris. 1837.

congestum Lam. 2. 2. T.

dioicum L. 2. D.

falcatum Lam. 2. ex h. Crac. 1837.

foetidum L. 3. (2) T. 1834 et ex h. Crac. 1837.

leyserioides Desf. (*Longohampia capillifolia* W. Spr. *Leysera discoidea* Spr. *Pectis discoidea* Horn.) 2. D. ex h. Paris. 1837.

margaritaceum L. 2. D. 1-2 pedal. flor. Aug.

maritimum L. 2. T. ex hort. Crac. 1837.

obtusifolium L. in hort. cult. 1806. herb. nostr.

orientale L. 2. F. 1810.

resedæfolium Trev. 2. C. ex h. Paris. 1837.

rosmarinifolium Sieb. 2. T. ex h. Crac. 1837.

spathulatum; 2. D. sub hoc nom. ex h. Paris. 1837.

Stocchas L. 2. F. 1810 et ex h. Paris. 1837.

lide flav. intr. 1834 ex h. Hamb.

Godetia v. Oenothera.**Gomphocarpus**

arborescens R. Br. v. *Asclepias*.

Gomphrena

decumbens Jacq. 2. C. 1810 et ex h. Hafn. 1837.

globosa L. 2. C. fl. purpur., carneo, albo.

interrupta 2. C. 1810 et ex h. Paris. 1837.

Gonolobus

rostratus R. Br. (*Cynanchum* Vahl.) 2. C.

Gossypium

herbaceum L. 2. C. ex hort. Hafn. 1835.

Grahamia

aromatica Spr. (*Gramia* Hook. *Cephalophora* Schrd.) 2. T. intr. 1834. 12—20 poll. alt.

Grangea

maderaspatana W. v. *Cotula*.
minima W. v. *Cotula*.

Gratiola (Jordgalla).

lucida Vahl. (*Morgania* Spr. *Capraria crustacea* L.) 2. C. ex h. Hafn. 1835.

officinalis L. 2. D. sesquiped. flor. Jul. — Sept.

serrata Roxb. v. *Bonnaya brachiata*.

Grevillea

buxifolia R. Br. 2. C. T. 20 poll. alt. intr. 1834 ex h. Hamb.

juniperina R., Br. 2. T. intr. 1834 ex h. Hamb.

linifolia; 2. T.

rosmarinifolia H. Hamb. 2. T. ex hort. Kew. intr. 1837. cur. Cl. J. Ag.

Gnidia

sericea Spr. (*Passerina* L.) 2. T. 33 poll. alt. flor. pal-

Grindelia

incisa Spr. (*Aster* Fisch.) 2.
T. F. ex h. Crac. 1837. 1
uln. alt.; flor. Aug., petal.
albo-cærul.

Gundelia

Tournefortii L. 2. F. ex h.
Petrop. 1836.

Gymnocladus

canadensis Lam. (*Gilundina di-*
oica L.) 2. D. 2½ uln. alt.

Gypsophila

acutifolia Fisch. 2. F. 1826.
arenaria Kit. (*G. fastigiata* var.
Spr.) 2. D. 1810. 1822.
herb. nostr.

dichotoma Bess. 2. D. ex hort.
Paris, 1837.

elegans M. B. 2. D. ex hort.
Paris, 1837. 1-pedal. flor.
albidis rubro-striat. Jul.
explic.

glomerata Pall. 2. F. 1826.
paniculata L. 2. D. ex hort.
Paris, 1837.

sabulosa Stev. 2. D. ex hort.
Paris, 1837.

saxifraga L. (*Dianthus* L. Sp.
1.) 2. D. 1808 herbar.
nostr.

scorzoneraefolia, sub hoc nom.
ex h. Paris, 1837.

Vaccaria Sm. (*Saponaria* L.)
2. D. bipedal. flor. parv.
roseis, Jul. explic.

Hablizia

tamnoides M. B. 2. F. ex h.
Hafn. 1836.

Hæmanthus

coccineus L. 2. C.
puniceus Jacqv. 2. C.

Hakea

angustifolia 2. T. ex Anglia
1837. cur. Cl. J. Ag.

dentata 2. T.
elliptica R. Br. 2. T. 1834 et
ex Angl. 1837.

florida R. Br. 2. T.
lanigera Booth. 2. T. ex Angl.
1837.

microcarpa R. Br. 2. T. intr.
1834.

microphylla Booth. 2. T. 18
poll. alt. flor. parv. albid.;
intr. 1834 ex h. Hamb.

ruscifolia Labill. 2. T. ex h.
Kewens. 1837.

saligna R. Br. (*Embothrium*
Andr.) 2. T.

Halesia

tetraptera L. 2. D. in hort. olim
sub Dio, sed sine succes-
su plantata. Cfr. Retz. in
Act. Holm. 1798.

Halimodendron

argenteum Cand. v. *Robinia Ha-*
lodendron.

Halleria

lucida L. 2. T. *Flora* 1836.
pag. 4

Haloragis

Cercodia Ait. 2. T. 1810
et ex h. Kil. 1835.

Hamamelis

virginica L. 2. D.

Harrachia

speciosa Jacqv. (*Justicia infun-*
dibuliformis L.) 2. C. —
etiam ex h. Crac. 1837.

Hasselqvistia

cordata L. 2. D. ex h. Paris.
1837.

Haworthia

concinna Haw. v. *Aloë viscosa*.

Hebenstreitia

tenuifolia 2. T. 1833 et sub
hoc nom. ex h. Paris. 1837.

Hedera (Murgrön).

Helix L. 2. D.

hibernica H. Hamb. \S . D. 1836.
quinquefolia L. v. *Vitis hederacea*.

Hedychium

coronarum Kön. \mathcal{A} . C.
Gardnerianum? Wall. \mathcal{A} . C.

Hedyotis

biflora H. Hafn. v. *Oldenlandia*.
diffusa W. (*Oldenlandia* Roxb.
 Horn.) \mathcal{A} . C. ex h. Hafn.
 1837.

Hedypnois

rhagadioloides L. (*Hyoseris cre-*
tica Cav.) \odot . D. 1810.
 1815 et ex h. Paris 1836.

Hedysarum

alpinum L. \mathcal{A} . D. F. 1826.
canadense L. \mathcal{A} . D. 1809. herb.
 nostr. et nova vice ex h.
 Paris. 1837.

canescens L. \mathcal{A} . F. 1811 herb.
 nostr.

Caput Galli L. v. *Onobrychis*.
carpathicum W. v. *Onobrychis*
petraea.

coronarum L. \mathcal{A} . D. ex hort.
 Bonn. 1836.

Crista Galli L. v. *Onobrychis*.
elongatum; \mathcal{A} . sub hoc nom. ex
 h. Petrop. 1836.

flexuosum L. \odot . T.
marilandicum L. \mathcal{A} . F. ex h.
 Hafn. 1836.

muricatum Jacqv. \odot . D. 1815
 herb. nostr.

obscurum L. \mathcal{A} . D. ex h. Petr.
 1834 et 1836.

Onobrychis L. v. *Onobrychis sa-*
tiva Lam.

petraeum M. B. v. *Onobrychis*.
saxatile L. v. *Onobrychis*.

Helenium

autumnale L. \mathcal{A} . D. 1808.
 herb. nostr.

mexicanum Humb. \odot . D. ex
 h. Paris. 1837.

quadridentatum Lahill. (*Rud-*
beckia alata Jacqv.) \mathcal{A} . \odot .
 T. 1821.

Helianthemum Tournf.

(*Cisti species* L.).

egyptiacum Mill. \odot . D. ex h.
 Paris. 1837.

ledifolium W. (*Cistus* L.) \odot .
 D. 1834. 1835.

mutabile Pers. \S . F. 1822 et
 ex h. Paris. 1837.

procumbens Dun. \S . F. 1834
 et ex h. Petrop. 1836.

salicifolium Pers. \odot . D. ex h.
 Paris. 1837.

thymifolium Pers. \S . F. ex h.
 Paris. 1837.

vulgare Gärtn. (*Cistus Helian-*
them. L. *Cist. grandiflorus*
 Scop.) \S . D.

Helianthus (Solros).

annuus L. \odot . D.

decapetalus L. \mathcal{A} . D. 1826.

indicus L. (*annuus* var. Spr.)
 \odot . D.

lenticularis. Lindl. \odot . D. ex
 h. Hafn. 1836.

multiflorus L. \mathcal{A} . D. 1815 herb.
 nostr.

pubescens Vahl. \mathcal{A} . D. ex h.
 Paris. 1837.

rigidus; sub hoc nom. ex h.
 Paris. 1837.

tuberosus L. \mathcal{A} . D.

Heliophila

amplexicaulis L. \odot . D. ex h.
 Paris. 1837.

pendula W. (*pinnata* Vcht.)
 \odot . D. ex h. Paris. 1837.

Heliopsis

laevis Pers. (*Bupthalm. helian-*
thoides L.) \mathcal{A} . D. ex hort.
 Paris. 1837.

Heliotropium

europaeum L. \odot . D.

indicum L. v. *Tiaridium*.

parviflorum L. \odot . C. 1812.
 herbar. nostr.

peruvianum L. \S . T. 15—16
 poll. alt.

β . fl. alb.

Helleborus (Prustrot).
 foetidus L. 2. D. in Paradis-
 lyceum.
 hyemalis L. 2. D. jam horti
 nostri indigena.
 niger L. 2. D. a frigore hie-
 mum non læditur.
 odoratus Kit 2. D. F. ex hort.
 Paris. 1837.
 viridis L. 2. D. coelo horti
 nostri familiarissimus.
Helminthia
 echinoides W. (*Pieris* L.) ♂. D.

Hemerocallis
 coerulea β . japonica W. v. *Fun-*
kia ovata.
 flava L. 2. D. 1826.
 fulva L. 2. D. 1826.
 graminea Horn. 2. D. 1826.
 Liliastrum L. (*Czackia Liliastr.*
 Andr.) 2. D. 1826.

Hemimeris
 urticifolia W. (*Celsia* Curt.)
 β . T. 1810 et ex h. Crac.
 1837.
 species, T,

Heracleum
 amplifolium Lapeyr. *pyrenai-*
cum Cuss. *speciosum* Hoffm.)
 2. D. 1826; et nov. vice
 ex h. Paris. 1837; 5 uln.
 alt.
 gummiferum W. ♂. 2. F. ex
 h. Paris. 1837.
 persicum; sub hoc nom. ex h.
 Paris. 1837.
 sibiricum L. ♂. D. 1826.
 Sphondylium L. ♂. 2. D.
 β . sub nom. *Wendta Cho-*
rodanum Hoffm. ex h. Pe-
 tropol. miss. 1836.
 species.

Hermannia
 angularis Jacq. β . T. 1810
 et ex h. Paris. 1836.
 denudata L. β . T. 1810 et ex
 h. Crac. 1837.

hirsuta Schrad. β . T. 1814
 herbar. nostr.
hyssopifolia L. β . T. 1810.
 1814—1837.

Hermesia
Banksiaefolia v. *Callicoma*.

Herniaria
 glabra L. 2. D.
 hirsuta L. 2. D.

Herpestris (*Herpestes*).
 chilensis H. Crac. T. sub hoc
 nom. ex h. Crac. 1837.
 crysantha H. Hafn. T. ex h.
 Hafn. 1836.

Hesperis (Nattviol).
africana L. v. *Malcolmia*
chia Lam. v. *Malcolmia*.
laxa Lam. v. *Chorisporea tenella*.
 elata Hornem. ♂. D. 1826.
 matronalis L. ♂. D.
 tristis L. ♂. D.
 verna L. (*Arabis* R. Br.) ♂. ♂.
 D. 1820.

Heterospermum
 pinnatum Cav. ♂. D. 1811.
 et iterum ex h. Par. 1837.

Heterotheca
 inuloides Cass. H. Lips.; ex h.
 Paris. 1837.

Heuchera
 americana L. 2. D. 2-3-ped.
 flor. Jun. — Aug.
 lucida H. Hamb. 2. F. ex h.
 Paris. 1837.
 macrophylla H. Hamb. 2. F.
 ex h. Paris. 1837.
 villosa Mx. 2. F. ex h. Paris.
 1837.
 species.

Hibbertia
 dentata R. Br. β . T. 3-pedal.
 flor. pallid. flav.
 volubilis Andr. (*Dillenia volubi-*
lis Vent. *scandens* W.) β .
 T. 1836.

Hibiscus
 Abelmoschus L. β . C.
 cannabinus L. ♂. C. 1835.

e seminibus maturatis
1834.
esculentum L. ☉. C. 1834 et
ex h. Hafn. 1835.
gossypinus Thb. ☐. C. 1834.
1835.
Humboldtii Mart. ☉. D. ex h.
Gryph. 1835. 6-8-pollic.
flor. albesc.
liliiflorus Cav. ☐. C.
Muluviviscus L. v. *Achania*.
Manihot L. ☐. 2. C. 1810 et
ex h. Hafn. 1836.
β. palmatus Cav. Horn.
mutabilis L. ☐. C.
pedunculatus Thb. ☐. C. 18
poll. alt. flor. ros
populneus L. (*Thespesia* Corr.)
☐. C.
Richardsonii Sweet. ☐. intr.
1834. ex h. Kil.
Rosa sinensis L. ☐. C.
speciosus Ait. 2. C. ex h. Kil.
1836.
splendens H. Hamb. C. intr.
1834.
surattensis L. ☐. ☉. C. 1834-
1837.
syriacus L. ☐. F. D. 1819-
1837.
tortuosus H. Hamb. ☐. C.
Trionum L. ☉. D.

Hieracium

aurantiacum L. 2. D.
aureum Scop. (*Leontodon* L.) 2.
F. 1822.
Auricula L. 2. D.
boreale Fries. (*sabaudum* L.
Retz.) D. 2., 1837 ex h.
Crac.
collinum Gochn. (*stoloniferum*
Reichb.) 2. ex h. Gryph.
1837.
cydonifolium Vill. 2. D. 1822.
cymosum L. 2. D. 1812-24.
herbar. nostr.
echioides Lumnitz. 2. D. 1810
et ex h. Paris. 1837.
foliosum Kit. 2. D. ex hort.
Paris. 1837.

fruticosum W. ☐. F. 1810 et
ex h. Paris. 1837.
glaucum All. 2. D. 1810 et
ex h. Gryph. 1837.
grandiflorum Horn. 2. D. 1810
et ex h. Paris. 1837.
Jacqvini Vill. 2. D. F. ex h.
Paris. 1837.
lavigatum W. 2. D. 1811 her-
bar. nostr.
lanatum Kit. 2. F. ex h. Pa-
ris. 1837.
lapsanoides Gouan. 2. D. ex
h. Crac. 1837.
Lawsonii Vill. 2. F. 1823;
etiam ex h. Paris. 1837.
Ledebourianum Tausch. 2. ex
h. Gryph. 1837.
nigrescens W. 2. D. 1810.
1815 et ex h. Paris. 1837.
nudicaule Mx. 2. D. ex hort.
Gryph. 1837.
Pilosella L. 2. D. spont.
polymorphum H. Crac.; com-
munic. 1837.
præmorsum L. 2. D.
pratense Tausch. 2. D. ex h.
Gryph. 1837.
ramosum Kit. 2. D. ex hort.
Paris. 1837 (an *glaucum*
All.?)
rupestre All. 2. D. 1822.
1833.
speciosissimum Hornem. 2. D.
ex h. Gryph. 1837.
speciosum Hornem. Wilbr. 2.
D. ex h. Crac. 1835,
Sternbergii Hopp. 2.; ex h.
Gryph. 1837.
umbellatum L. 2. D. etiam ex
h. Gryph. 1837.
β. coronopifolium Bernh.
villosum L. 2. D. 1 uln. alt.

Hierochloa

borealis Schrad. et Fr. (*Hol-
cus odoratus* L.) 2. D.

Hippoerepis

ciliata W. ♂. J. D. 18 poll.
alt. in hort.; flor. flav. Jul.
explic.

comosa L. 2. D. 1809 herb.
nostr.

multisiliqua L. ♂. D. 1835.

syriaca Ehrenb. ♂. D. ex h.
Kil. 1835.

Hippomarathrum Link.

(*Cachrys*).

crispum; sub hoc nom. ex h.
Petrop. 1836.

Hippophaë (Haftorn).

canadensis L. (*Shepherdia* Spr.)

♂. 1826 in hort. nostro
cult., hodie ibi exstirpata.

conferta Booth. ♂. F. ex Angl.
1837. cur. Cl. J. Ag.

rhampnoides L. ♂. D. in nov.
arbor. 5 uln. alt. etiam in
plantatione Paradislyckan.

Hirtella

species.

Hoitzia

coccinea Cav. (*Cantua Hoitzia*
W. *Collomia coccinea* Lehm.)

♂. D. ex hh. Hafniæ et
Bonn. 1836.

linariæ Spr. v. *Collomia*.

Holcus (Tâtel).

avenaceus Scop. *Avena elatior*
L. 2. D. 1826.

lanatus L. 2. D.

mollis L. 2. D. 1826.

odoratus L. v. *Hieroecht. borealis*.

saccharatus Andr. v. *Sorghum*.

Sorghum L. v. *Sorghum vulgare*.

Holostigma

cheiranthifolium; s. hoc nom.
ex h. Paris. 1837.

heterophyllum; ex h. Paris.
1837.

Hordeum (Korn).

bulbosum L. 2. D. 1826, et
iter. ex h. Paris. 1837.

secalinum Schreb. 2. D. 1826.

vulgare L. ♂. D.

Horminum

pyrenaicum L. v. *Melissa*.

Hornemannia

bicolor W. (*Gratiola goodeni-*
folia Horn.) ♂. C. intr.
1807 secund. herb. nostr.;
1834, et iterata vice ex
h. Hafn. 1837.

Hoya

carnosa R. Br. (*Asclepias* L.
Stupelia chinensis Lour.)

♂. C. T.

Humea

elegans Sm. ♂. T. ex h. Kil.
1835.

Humulus (Humle).

Lupulus L. 2. D.

Hura

crepitans L. ♂. C.

Hyacinthus (Hyacint).

ciliatus Cyr. v. *Muscari*.

Muscari L. v. *Muscari Mascha-*
tum.

non scriptus L. v. *Scilla*.

orientalis L. 2. D. fl. alb.,
cærul., ros. et plen.

romanus L. 2. F. ex h. Paris.
1837.

Hydrangea

arborescens L. ♂. F.

hortensis Sm. ♂. F. etiam sub
Dio certis annis sine di-
scrimine hiemis colitur.

nivea Mx. (*radiata* Walt.) ♂.
D. 1 uln. alt. flor. alb.
Aug. explic.

Hydrocotyle

sibthorpioides Lam. 2. C. ex
h. Paris. 1837.

Hydrophyllum

canadense L. 2. D. 1826 et
nova vice ex hort. Paris.
1837.

magellanicum Lam. v. *Phacelia*
circinnata.

- virginicum* L. 2. D. ex hort. Paris. 1837.
- Hymenanchera**
dentata R. Br. 2. T. ex hort. Kew. 1837. cur. Cl. J. Ag.
- Hymenopappus**
pedatus Lag. Spr. (*Stevia* Cav. W.) 2. F. D. 1806. etiam 1837.
- Hyoscyamus**
albus L. 2. 2. D. 1810.
aureus L. 2. F. ex h. Bonn. 1836.
niger. L. (Bolmört) 2. D. sp.
orientalis M. B. 2. D. ex h. Paris. 1837.
physaloides L. 2. D. ex hort Paris. 1837.
pusillus L. 2. D.
Scopolia L. 2. D. usque ad 1½ uln. alt.
- Hyoseris**
cretica Cav. v. *Hedynois-rhagadioloides*.
Hedynois L. (*Hedyn. rhagadioloides* var. Spr.) ex h. Paris. 1837. sub hoc nom.
lucida L. 2. D. 1810. 1822.
scabra L. 2. D. ex h. Paris. 1836.
tubæformis Tenor. 2. sub h. nom. ex h. Paris. 1837.
- Hypericum**
alatum Retz. Pugill. 1810 p. 16. 2. F.; hodie in hort. ignotum.
Androsæmum L. v. *Androsæmum officinale*.
Assyron L. 2. D. F. 1826.
balearicum L. 2. F. T. ex h. Kil. 1835.
corymbosum Mühl. 2. F. ex h. Gryph. 1836.
crispatum Retz. l. c. 2. F. 1810 in horto viguit, ibi hodie extirpatum.
dubium Leers 2. D.
- elegans* W. 2. F. ex h. Kil. 1836.
fimbriatum Lam. 2. F. ex h. Gryph. 1836; sub nom. *H. Richeri* ex h. Paris. 1837. forte idem.
floribundum Ait. 2. T. F. ex h. Kil. 1836.
hirsutum L. 2. D. 1826.
humifusum L. 2. D. ex hort. Kil. 1835.
intermedium Retz. l. c. 2. D. et *retusum* Retz. l. c. 2. F. ex America boreali, 1810—1818 et 1821? in hort. nostro culta, hodie profuga.
macrocarpum Mx. (*ascyroides* W.) 2. F. D. ex h. Kil. 1836.
perforatum L. 2. D.
montanum L. 2. D. ex hort. Hafn. 1827 et Gryph. 1837.
pulchellum W. 2.; ex h. Gr. 1836.
pyramidatum Ait. 2. D. 1809. herb. nostr.
tomentosum L. 2. F. 1810 et ex h. Paris. 1837.
undulatum Schousb. 2. F. 1810 et ex h. Paris. 1837.
- Hypochæris**
hispida W. (*Achyrophorus* Horn.) 2. D. 1810—1822.
pinnatifida Cyr. Tenor. (*Robertia* Spr.) 2. D. ex hort. Paris. 1837.
radicata L. 2. D.
- Hypoxis**
gracilis H. Hamb. T. ex hort. Paris. 1837.
- Hyptis**
capitata Jacq. (*Clinopodium* Sv.) T.
- Hyssopus** (Isop).
coymifolius Lam. v. *Elsholtzia cristata*.
officinalis L. 2. D.
scrophulariaefolius W. 2. D. olim h. incola, hodie exsul.

Jasminum

- azoricum L. ♀. T.
 fruticans L. ♀. T.
 odoratissimum L. ♀. T. 2½
 uln. alt.
 Sambac L. ♀. C. T.
 Wallichianum ♀. T. intr. 1837
 ex Anglia sub hoc nom.
 cur. Clar. J. Agurdk.

Iberis

- amara L. ☉. D. 8-10 pollic.
 alt. flor. alb., Aug.
 β. intermedia Guers.
 arabica L. v. *Æthionema cappu-*
docicum.
 contracta Pers. ♀. ☉. D. intr.
 1837. ex h. Kil.
 gibraltarica L. ♀. F. flor. albo-
 incarnat.
 integerrima Moris. ☉. D. ex h.
 Kil. 1837.
 Lagascana Cand. (*pubescens* W.)
 ☉. ♀. D. F. 12 poll. alt.
 flor. alb.
 odorata L. ☉. D. 12 pollic.
 alt. fl. alb. Aug.
 pinnata L. ☉. D. 1812. 1837.
 flor. alb.
 sempervirens L. ♀. D. 6--8
 pollic. alt. flor. alb. Maj.
 — Decemb. in horto.
 Tenorii Cand. ♀. T. 9-11 pol-
 lic. alt.
 umbellata L. ☉. D. fl. roseo
 aut rubro-violaceo.
 β. superba ex hh. Kil. et
 Hamb. 1836.

Ilex (Jern-ek).

- æstivalis Lam. (*prinoides* Ait.)
 ♀. F. ex h. Paris. 1837.
 Aquifolium L. ♀. F. D., per
 2 hiemes sub dio duravit.

- balearica Desf. ♀. F. ex hort.
 Paris. 1837.

Ilecebrum

- lanatum L. W. (*Celosia* L.
Ærva Juss.) ☉. C. 1810
 et ex h. Hafniae 1836.
 sessile W. (*Gomphrena* L. *Pa-*
ronychia sess. H. Par.) ☉.
 C. 1810. 1837.
 spinosum Spr. (*Achyranthes* Hor-
 nem.) ☉. ♀. C. ex hort.
 Hafn. 1837.

Impatiens

- Balsamina L. ☉. C. fl. color.
 et plenit. var.
 β. capensis Thb. ex hort.
 Hafn. 1837.
 nolitangere L. ☉. D. 1834 in
 umbros. sesquipedal.; flor.
 flav.
 parviflora Cand. ☉. C.

Imperatoria

- caucasica Spr. (*Selinum* M. B.)
 2. ex h. Petr. 1836.
 pimpinelloides; sub hoc nom.
 ex h. Bonn. 1836.
 ruthenica; sub hoc nom. ex h.
 Paris. 1837.

Incarvillea

- chinensis Lam. ♀. C. ex hort.
 Petr. 1836.

Inula

- arabica L. 2. T. ex h. Paris.
 1837.
 Bubonium Jacq. 2. D. 1826-
 1835. 26 poll. alt. flor.
 flav. Septemb. explic.
 crithmoides (*crithmifolia* W.)
 ♀. F. ex h. Par. 1837.
 dysenterica L. 2. D. 1807 herb.
 nostr.
 gnaphaloides; sub hoc nom. in
 horto 1826.
 Helenium L. 2. D. 4-pedal. et
 ultra.
 hirta L. 2. D. 1821--1835;
 etiam ex h. Paris. 1837.
 sesquipedal.
 Oculus (Christi) L. 2. D. ex
 h. Paris. 1837.
 Pulicaria L. ☉. D. vix pe-
 dalis.

salicina L. 2. D. 1826. 1835.
18-24 poll. alt.
squarrosa L. 2. D. fere 1 uln.
alt., flor. flav. Septemb.
verbascifolia Poir. 2. D. 1826.

Ipomoea v. *Convolvuli* species.

Iris (Svärdslilja).

florentina L. (Fiolrot). 2. D.
1½ uln. alt. flor. lact. Jul.
germanica L. 2. D. sesquipedal.
flor. cyaneis.
graminea L. 2. D. scap. 18,
folia 30 poll. alt. flor. vio-
laceo-cærul. Jul. explic.
Monnerii Cand. 2. D. ex h.
Paris. 1837.
nepalensis Wallich. 2. T. ex
Angl. 1837. cur. Cl. J. Ag.
ochroleuca M. B. 2. D. simili-
lis *spuria*; intr. 1805.
pallida Lam. 2. D. 2 uln. alt.
flor. odor. albo-cærul., bar-
ba alba.
pavonia L. (*Moræa* Ker. *Vicus-
seuxia* Cand. Spr.) 2. T.
persica L. 2. T. F.
Pseudacorus L. 2. D. (in sicco).
pumila L. 2. D. 1818 intr.
3-6-pollicaris.
sambucina L. (*J. squalentis* var.
Spr.) 2. D. 1826.
sibirica W. 2. D. intr. 1812.
1½ uln. alt.
β. fl. alb.
spuria L. (*spatulata* Lam.) 2.
D. 20-22 poll. alt.
squalens L. (*sordida* Retz.) 2.
D. 1½ uln. alt.
Swertii Lam. ? 2. D. 1836.
variegata L. 2. D.
versicolor L. 2. D. 1833 et
ex h. Hafn. 1835.
virginica L. 2. F. D. ex h.
Hafn. 1837.
Xiphium L. 2. F. D. 1½ uln. alt.

Isatis (Wejde).

canescens Cand. (*iberica* Stev.)
♂. D. ex hh. Petr. 1836,
et Paris. 1837.
latisiliqua Stev.; F. ex hort.
Petr. 1836.
littoralis Cand.; F. ex h. Pa-
ris. 1837.
orientalis W. ♂. D. 1835.
præcox Kit. ♂. D. 1826.
tinctoria L. ♂. D.

Isopogon

anemonifolius R. Br. (*Protea*
Salisb.) ♀. T. ex Angl. 1837.

Juglans (Valnöt).

amara Mx. (*Carya* Spr.) ♀. D.
1826 in hort. plant., sed
mox emortua.
cinerea L. ♀. 15 uln. alt.;
trunci diametr. juxta ter-
ram circit. 12-poll.; in
Plantatione Paradislyckan.
nigra L. ♀. D. 27 uln. alt.,
trunci diametr. 20-poll. in
Paradislyckan.
regia L. ♀. D. 30 uln. alt. trunc.
diametr. 1½ uln.; in Para-
dislyckan.
β. nucæ majori; in Para-
dislyckan 29 uln. alt.

Juncus

helodes Link. 2. D. ex hort.
Paris. 1837.

Juniperus (En).

Oxycedrus L. ♀. D. 11 uln. alt.
Sabina L. ♀. D. (Säfvenbom)
3 uln. alt., floret April.
virginiana L. ♀. D. 12 uln. alt.

Jurinea

mollis; s. h. nom. ex h. Petr. 1836.

Jussieua

erecta L. ♂. C. 1805 in hort.
cultæ, recentiori tempore
ibi ignota.
scabra W. C. ex hort. Paris.
1837. cur. Cl. J. Ag.

Justicia

- Adhatoda* L. ♀. T.
infundibuliformis L. v. *Harra-*
chia speciosa.
paniculata Burm. 2. T. ex h.
 Hafn. 1837.
parviflora Horn. v. *Eranthemum*.

Ixia

- crocata* Ker. v. *Tritonia squalida*.

Ixodia

- Achilleoides* R. Rr. ♀. T. C.
Ammobium Spr. (*Ammobium*
alatum R. Br.) 2. T. F. 8
 poll. alt. flor. med. Sept.

Kalmia

- glaucia* Ait. ♀. F. 22 poll. alt.
 intr. 1833 ex h. Hamb.
 floruit mens. Oct.

Kaulfussia

- amelloides* Nees. ☉. C. 1834.
 1835. hodie horto abest.

Kennedia

- prostrata* R. Br. v. *Glycine coc-*
cinea.

Keria

- japonica* Cand. (*Rubus* L. *Cor-*
chorus Thb.) ♀. D. 2½ uln.
 alt. intr. 1818. — Para-
 dislyckan.

Kitaibelia

- vitifolia* W. 2. D. 2 uln. alt.
 flor alb. Sept. Oct.

Knautia

- hybrida* v. *Scabiosa*.
orientalis L. ☉. D. 3-pedal.
 et ultra; flor. incarn.
plumosa L. (*Scabiosa* Sm. *Ptero-*
cephalus Coult. Spr.) ☉. D.
 1810 et ex h. Paris. 1837.

Koeleria

- rigida*; ex h. Hafn. 1837.

Koelpinia

- linearis* Pall. (*Rhagadiolus Köl-*
pinia W. Spr. *Lapsana Köl-*
pinia L.) C. ex h. Petrop.
 1836.

Koelreutera

- paniculata* Lam. ♀. D. intr.
 1832 in nov. nostr. arbor-
 retum, ubi jam 8 uln. alt.

Kydia

- calycina* Roxb. ♀. C.

Kyllingia

- monosephala* W. 2. C. 1822.
triceps Spr. (*Scirp. glomeratus*
 L.) 2. C. 1810.

Lachenalia

- tricolor* Jacq. v. (*Phormium aloi-*
des L. suppl.) 2. T.

Lactuca (Laktuk).

- angustana* All. ☉. D. ex h.
 Paris. 1837.
elongata Mühl. ♂. D. ex hort.
 Paris. 1837.
perennis L. 2. D.
sagittata Kit. ♂. D. ex hort.
 Paris. 1837.
sativa L. ☉. D.
 β. *capitata*.
 γ. *crispa*.
Scariola L. Fr. ♂. ☉. D.
stricta Kit. ♂. D. 1815 et ex
 h. Paris. 1837..
virgata; ☉. D. ex h. Paris.
 1837.

Lasagea

- mollis* Cav. ☉. C. intr. 1806,
 et ex h. Paris. 1837.

Lagoecia

- cuminoides* L. ☉. D. ex hort.
 Paris. 1837.

- Lagunæa**
sinuata Horn. (*lobata* W.) ☉.
 C. 1810 et ex h. Hafniæ
 1835.
- Lagurus**
ovatus L. ☉. D.
- Lamarckia**
aurea Mönch. Cand. (*Cynosurus* L. *Chrysurus* P. B.)
 ☉. D. ex h. Paris. 1837.
- Lambertia**
formosa Sm. ♀. T. ex h. Kew.
 1837.
- Lamium** (Plister).
album L. ♀. D. spont.
amplexicaule L. ☉. D. spont.
garganicum L. ♀. D. 8--10
 poll. alt.
incisum W. Fr. ☉. D. spont.
intermedium Fr. ☉. D. spont.
Orvala L. ♀. D. 10-12 poll.
 alt. flor. magn. rubr., ma-
 culat.
purpureum L. ☉. D. spont.
- Lantana**
acanthifolia; ♀. C. intr. 1837
 ex h. Paris.
scabrida Ait. ♀. C. 1822 herb.
 nostr.
trifolia L. ♀. C. intr. 1837
 ex h. Paris.
- Lappago**
racemosa Schreb. (*Cenchrus* L.)
 ☉. D. 1808. herb. nostr.
- Lapsana**
communis L. ☉. D. spont.
crispa W. ☉. D. 1½ uln. alt.
 flor. flav. Aug. explic.
Kölpinia L. v. *Koelpinia linariis*.
lyrata W. ♀. D. 1½ uln. alt.
 flor. pallide flav. Jun.
stellata L. v. *Rhagadiolus*.
- Larochea**
falcata Spr. Cand. v. *Crassula*.
- Laserpitium**
angustifolium L. (*peucedanoides*
 Spr.) ♀. D. 1833, et ex
 h. Paris. 1837.
- aquilegifolium* Jacq. (*Siler aqi-
 legifol.* Gärtn.) ♀. D. 1833,
 et ex h. Paris. 1837.
gallicum Tournef. ♀. D. ex h.
 Paris. 1837.
latifolium L. ♀. D. etiam ex
 h. Paris. 1837.
- Lasiopetalum**
solanaceum? ex h. Crac. 1837.
- Lasiospermum**
pedunculare Lag. (*Santolina al-
 pina* L. *Cotula abrotanifolia*
 W. *Santol. pinnata* Donn.
 et H. Lund.) ♀. ♂. F.
 1806 herbar. nostr.
- Lasiospora**
lanata; sub hoc nom. ex hort.
 Petropol. 1836
latifolia; sub hoc nom. ex h.
 Petropol. 1836.
- Lasthenia**
glabrata Lindl., sub hoc nom.
 ex h. Paris. 1837.
- Lathyrus**
amphicarpos L. ☉. D. F. 18
 poll. alt. flor. rubr.
annuus L. ☉. D. 1809. herb.
 nostr.
Aphaca L. ☉. D. flor. flav.
articulatus L. ☉. D. 1812; et
 ex h. Gryph. 1837.
auriculatus Bertol. ☉. D. 1835.
 1 uln. alt. flor. purpur.
californicus Dougl. ♀. F. 1835
 florentem non vidi.
Cicera L. ☉. D. 1810. 1812.
 1837.
Clymenum L. ☉. D. 1812-
 1837.
grandiflorus Sibth. ♀. F. ex
 h. Gryph. 1837.
heterophyllus L. ♀. D. 1814.
hirsutus L. ☉. D. 1812-1821.
incurvus L. ♀. D. ex h. Par.
 1837.
latifolius L. ♀. D. 3 uln. alt.
lusitanicus; ☉. D. sub hoc nom.
 ex h. Paris. 1837.
Nissolia L. ☉. D. 1809. 1815.
Ochrus Cand. (*Prism* L.) ☉.
 D. flor. sordide flav.
odoratus L. ☉. D.

- β. fl. violaceo- et rubro-
 striato.
 pratensis L. 2. D. spont.
 purpureus Desf. (*spurius* W. an
 etiam *auriculatus* Berth.?)
 ⊙. D. 1812, et ex h. Gr.
 1837.
 rotundifolius W. ⊙. D. 1835.
 hodie abest.
 sativus L. ⊙. D.
 β. fl. cæruleo.
 sphaericus Retz. ⊙. D. 1835.
 sylvestris L. 2. D.
 tenuifolius Desf. 2. ex h. Gr.
 1837.
 tuberosus L. 2. D.
- Laurus**
 camphora L. (*Persea* Spr. Sv.
 Kamferträd) ½ T. 1 uln. alt.
 nobilis L. (Lager) ½. T. F.
- Lavandula** (Lavendel).
 multifida L. ⊙. ♂. F.
 pinnata L. ½. F.
 spica W. ½. D.
 β. *latifolia* W. 1808. herb.
 nostr.
- Lavatera**
 arborea L. ♂. ½. F. 2½ uln.
 alt. 1824 et iterum intr.
 1832 ex h. Gryph.
 cretica L. ⊙. D. 1808. herb.
 nostr.
 maritima Gouan. ½. T. F. 1824
 et ex h. Paris. 1837.
 micans L. (*lusitunica* L.) ½. F.
 1810 et ex h. Paris. 1837.
 olbia L. ½. T. F. 1810, et ite-
 rum intr. 1832 ex h. Hafn.
 hodie 3½ uln. alt.
 pseudolbia Poir. ex h. Paris.
 1837.
 silvestris Brot. ⊙. F. olim (1810)
 horti incola, hodie exsul.
 thuringiaca L. 2. D.
 trimestris L. ⊙. D. 1½ uln.
 alt. flor. rubr. l. roseo.
 β. fl. albo.
- Lecanocarpus**
 nepalensis; sub hoc nom. ex h.
 Paris. 1837.
- Lemna**
 minor. L. 2. D. spont.
- Leontodon**
 hostile L. v. *Apargia*.
 hirtum L. v. *Thrinicia*.
 hispidum L. v. *Apargia*.
 Taraxacum L. 2. D. spont.
 tuberosum L. v. *Thrinicia*.
- Leonurus**
 altaicus Spr. 2. D. ex h. Pa-
 ris. 1837.
 Cardiaca L. 2. D.
 crispus Murr. 2. D. 1826.
 Marrubiastrum L. 2. ♂. D.
 ex h. Paris. 1837.
 sibiricus L. ⊙. ♂. 2. D. 1805;
 etiam ex h. Paris. 1837.
 tataricus L. ♂. D.
 villosus Desf. 2. D. ex h. Pa-
 ris. 1837.
- Lepigonum** F. et M.
 species ex h. Petropol. 1836.
- Lepidium** (Krasse).
 campestre R. Br. (*Thlaspi* L.)
 ♂. D.
 chilense. H. Par. ⊙. D. ex h.
 Paris. 1837.
 cordatum W. herb. 2. ex h.
 Paris. 1837.
 didymum L. v. *Coronopus*.
 Draba L. (*Cochlearia* Draba L.
 sp. 2.) 2. D. 1810. 1834.
 Ecclonii Schrd. ⊙. D. ex h.
 Paris. 1837.
 graminifolium L. (*L. Iberis*
 Spr.) 2. D. ex h. Crac.
 1837.
 latifolium L. 2. D. 1810-1826.
 ruderales L. ⊙. D.
 sativum L. (*Thlaspi* Crantz.)
 ⊙. D.
 β. *crispatum* Hornem.
 virginicum L. ⊙. D. 1815 et
 ex h. Paris. 1837.
- Leptospermum**
 virgatum Forst. v. *Bäckeua*.

Leucadendron

Levisianus Berg. (*Protea* L. Prot.
hirsuta W. herb.) ½. T.

Leucas

martinicensis R. Br. (*Phlomis*
W.) ☉. C. ex h. Paris.
1837.

Leucoium

aestivum L. ¼. D.
vernum L. ¼. D.

Leucopogon

Richet R. Br. v. *Styphelia*.

Leuzea

carthamoides Cand. (*Cnicus* W.)
¼. D. ex h. Petr. 1836.

Leycesteria

formosa Wall. ½. T. ex Angl.
1837.

Leyssera

discoidea Spr. v. *Gnaphal. ley-*
serioides.

gnaphaloides L. ½. C. intr.
1837 ex h. Paris.

Libanotis (Athamanta).

villosa; ex h. Petrop. 1836 s.
hoc nomiu.

Ligusticum

apiifolium; sub hoc nom. ex
h. Paris. 1837.

Fischeri H. Hamb. ¼. D. ex
h. Hafn. 1837.

Levisticum L. (*Libbsticka*)
¼. D.

peloponense Horn. ¼. D. ex
h. Hafn. 1835.

pyrenaicum Gouan. *Cnidium*
Spr. *Seseli aristatum* Ait.
1826 et ex h. Par. 1837.

Ligustrum (Liguster).

sinense Lour. v. *Syringa villosa*.
valgare L. ½. D.

β. *foliosum*; ex h. Hamb.
1836.

γ. *italicum* MiH. Retz. in
Act. Holm. 1798. p. 59;
hodie in hort. ignot.

Libium (Lilja).

hathiferum L. ¼. D. sere 3-ped.
candidum L. ¼. D. 3-pedal.

chalconicum L. ¼. D. 1½
uln. alt. flor. flav.

Martagon L. ¼. D.
β. fl. plen.

pouponicum L. ¼. D. ¾ uln.
alt. fl. cinnabarin.

tigrinum Ker. ¼. D. 1½ uln.
alt. fl. Aug. explicat. mi-
nist. nigro-punctat.

**Linaria (Antirrhini species
Linw.)**

acutangula H. Crac.; T. ex h.
Crac. 1837, an *Lin. Cym-*
balaria?

alpina Cand. ¾. ¼. F. ex h.
Gryph. 1837.

bipartita W. (*speciosa* Jacqv.)
♂. ☉. D. intr. 1806; re-
centius ex hh. Kil. et Gr.
1836. 1837. fl. flavo aut
rubente; herba ¾ uln. alt.

chalepensis Mill. ☉. D. 1812
herbar. nostr., etiam ex
h. Paris. 1837 miss.

cirrhusa W. ☉. D. ex hort.
Paris 1837.

Cymbalaria W. ¼. D. intr.
1808.

Elatine Desf. ☉. D. flor. lut.
calcarea albid.

genistaeifolia MiH. ¼. D. 1810.
herb. nostr.

micrantha Spr. (*Antirrhin. Cav.*)
☉. D. 1808 horti incola,
recentiori ævo profuga.

minor Cand. ☉. D. flor. parv.
dilute violac.

monsessulana (*Antirrhin. L. Li-*
maria striata Spreng. *Ant-*
tirrhin. repens L.) ¼. T.
1808—1818.

odora Retz. (*Antirrhin. M. B.*
Limuria Lœselii Spr.) 1807
in horto culta; in præsen-
ti abest.

parviflora Desf. ex hort. Crac.
1837.

polycerata H. Crac. ex hort.
Crac. 1837.

- purpurea* Mill. 2. D. 1808-1835.
pyrenaica Cand. ☉. 2. F. ex h. Kil. 1837.
reticulata Smith. 2. T. ex h. Gryph. 1836.
spuria W. ☉. D. ex h. Paris. 1837.
stricta Hornem. 2. D. ex h. Gryph. 1836.
supina Desf. 2. D. 1808-1823. flor. flav.
triphylla W. ☉. D.; olim in hort. culta, jam. deest.
versicolor Mönch. ☉. D. ex h. Gryph. 1836.
vulgaris Bauh. (*Antirrh. Linaria* L.) 2. D. spont. specics.

Linum (Lin. Hör),

- africanum* L. (*æthiopicum* Thb.) 2. T. 1835-1837, 20-pollicare, fl. flav. mens. Sept. explic.
anglicum Mill. (*perenne* Sm.) 2. D. ex h. Crac. 1837.
angustifolium Hornem 2. D. ex h. Hafn. 1836, fl. cærul.
asiaticum H. Crac. D. ex h. Crac. 1837.
austriacum L. 2. ♂. D. ex h. Hafn. 1836, fl. cærul.
catharticum L. ☉. D.
croaticum H. Crac. ☉. D. ex h. Crac. 1837.
decumbens Desf. 2. F. ex h. Crac. 1836.
grandiflorum Desf. ☉. 2. D. ex h. Crac. 1836. 20 pollic. alt.
maritimum L. 2. F. ex hort. Paris. 1837.
monogynum Sm. 2. F. 1½ uln. alt. flor. alb. intr. 1834 ex h. Londin.
nervosum Kit. ☉. 2. ex h. Crac. 1837.
perenne L. 2. D. flor. cærul.
Radiola L. v. *Radiola Millegrana*.

- usitatissimum* L. ☉. D.
 β. fl. alb. ex h. Kil. 1836.
viscosum L.? ☉. D. ex h. Kil. 1837.

Lippia

- triphylla* Kunth. (*Verbena Herit. Lippia citriodora* Spr.) 2. T. 26 poll. alt.; fl. parv. alb. odor. citr. spirant.

Liriodendron

(Tulpanträd).

- Tulipifera* L. 2. D. 4 uln. alt.; intr. 1835. folia jam mens. Maj. explicantur; bene a dio viget, sed flores non protulit.

Listera

- ovata* R. Br. (*Ophrys* L. *Neottia latifolia* Spr.) 2. D. olim in hort. nostr. plantata, ultra 20 annos ibi viguit et floruit, sed jam extirpata. Cfr. Retz. Flor. Oecon. p. 473.

Lithospermum

- arvense* L. ☉. D. spont.
officinale L. 2. D. in umbros. bipedale, flor. albo-flavescente.
orientale Vahl. (*Anchusa* L.) 2. F. ex h. Paris. 1837.

Loasa

- palmata* Spr. Trev. (*Blumenbackia insignis* Schrad.) T. ex h. Paris. 1837.
triloba Juss. ex h. Kil. 1835. flor. flav.

Lobelia

- alata* R. Br. 2. T. ex h. Hafn. 1837.
bicolor Curt. 2. T. ex h. Kil. 1836.
canariensis Juss. ex hh. Kil. 1836, Paris. 1837.
candida Hort. ☉. ex h. Gr. 1836.
Cardinalis L. 2. F. fl. coccin.
debilis L. ☉. C. ex h. Gryph. 1837.

Erinus L. ☉. T. flor. cærul.
lab. infer. albo-bimaculato.
fulgens Humb. 2. T. F. 1½
uln. alt. fl. coccin.
inflata L. ☉. ♂, F. T. 1810
et ex h. Kil. 1836, pedal.
flor. cærnul.
longiflora L. 2. C. 1834 et ex
h. Paris. 1837.
Nuttalliana R. et Sch. ☉. 2.
ex h. Gryph. 1836.
rhizophyta Spr. (*decumbens* Bot.
Mag.) 2. T. ex h. Hafn.
1837.
Salzmanniana Presl. ex h. Kil.
1835.
secunda L. ☉. T. ex h. Crac.
1836.
siphilitica L. 2. F. D. 21 poll,
alt. flor. cæruleis.
triquetra L. 2. 1810 et ex h.
Paris. 1837.
Tupa L. ♀. T. 1810 et ex h.
Paris. 1837 cur. Cl. J. Ag.
urens L. ☉. 2. F. ex h. Crac.
1836.

Lolium (Repe)-

arvense Wither. (*canadense*
Mx.?) ☉. D.
perenne L. 2. D. spont.
temulentum L. ☉. D.

Lomandra

longifolia Labill. (*Xerotes* R.
Br.) 2. ? T. ex h. Paris.
1837.

Longchampia

capillifolia W. Spr. v. *Gnaphal.*
leyserioides.

Lonicera

(Try. Benved. **Caprifolium**).

alpigena L. ♀. D. intr. 1826.
in nov. arbor., ubi jam
2½ uln. alt. flor. brunn.
antheris flav. Jun. explic.
Caprifolium L. ♀. D.
cærulea L. ♀. D. fl. sordide lut.

Diervilla L. (*Diervilla canadensis* W.) ♀. D. Paradisl.

dioica L. ♀. D. 3 uln. alt. fl.
miniat. Jun.

iberica M. B. ♀. D. 1 uln.
alt.; intr. 1834, seminib.
educata.

orientalis Lam. ♀. F. D. ex h.
Petrop. 1836.

Periclymenum L. ♀. D.

sempervirens L. ♀. D. 2-3 uln.
alt. flor. Jun. explicat.
miniat.

symphoricarpos L. v. *Symphoria*
glomerata.

tatarica L. ♀. D. in Planta-
tione Paradislychan.

Xylosteum L. ♀. D.

Lopezia

mexicana Jacqv. ☉. D. sesqui-
ped. flor. rubr. Aug.

β. *racemosa* Cav. ex hort.
Paris. 1837. miss. a var.
α non differt.

Lophospermum

scandens (erubescens) Don. H.
Hamb. 2. T. intr. 1835
et 1836 ex h. Bonn. se-
mina in hort. nostro bene
maturat.

Lotus

ambiguus Bess.

arboreus Forst. v. *Carmichaëlia*
australis.

conjugatus L. (*Tetragonolobus*
Link.) ☉. D. ex h. Par.
1836. flor. flav.

corniculatus L. 2. D. spont.

diffusus Smith. ☉. D. ex h.
Petr. 1836.

edulis L. ☉. D. 1835 fl. flav.
Gebelia Vent. 2. T. ex hort.
Petr. 1836.

Jacobæus L. ☉. 2. T. 1812,
etiam ex h. Paris. 1837.
flor. badiis.

peregrinus L. ☉. D. ex hort.
Paris. 1836. fl. flav.

siliquosus L. (*Tetragonolob.* Roth.)
2. D. 1826.

tenuifolius Poll. 2. D. 16 poll.
alt. flor. flav. Aug.

tenuis Kit. 2. D. 1821. 1822.

tetragonolobus L. (*Tetragonolobus purpureus* Mönch.) ☉.
D. 7 poll. alt. flor. umbrin. Jul.

Lunaria

annua L. (*biennis* Mönck.) ♂.
D. pedal. flor. rubro-violac. mens. Maj. explic.
rediviva L. ♀. planta pedalis, flor. albo-rubro mens. Maj.
— Novemb.

Lupinus

albus L. ☉. D. 1810. 1812, et ex h. Kil. 1836. 1½ uln. alt. petal. alb. carina apice cærulescente.
angustifolius L. ☉. D. 1812-1836. tripedalis. fl. cærul.
bicolor Lindl. Ag. ☉. D. ex h. Hafn. 1834. Kil. 1836.
canaliculatus Sweet. ☉. D. ex h. Kil. 1836.
Cruikshankii Lindl. ☉. D. ex h. Kil. 1835. est var. *L. mutabilis* sec. Ag. flor. cærul.
flexuosus Dougl. ♀. ex h. Gr. 1837.
hirsutus L. ☉. D. fl. cærul. 1810. 1834. tripedalis.
linifolius Roth. Ag. ☉. D. 1834. 1836.
littoralis Dougl. Ag. ♀. D. F. ex h. Kil. 1836, etiam ex h. Chiswick Angliæ 1837 cur. Cl. J. Ag.
luteus L. ☉. D. sesquipedalis; flor. flav. Aug.
Marshallianus Sweet. Ag. ♀. D. ex h. Gryph. 1836. fl. cærul.
micranthus Dougl. Ag. ☉. intr. 1834, etiam ex h. Paris. 1837 missa. 12 poll. alt. fl. parv. cærul.
mutabilis Sweet. Ag. ☉. ex h. Gryph. 1836. flor. albis.
nanus Dougl. Benth. Ag. ☉. D. intr. 1834 ex h. Lon-

din.; *Flora* 1836. pag. 7. flor. cærul.

nootkatensis Donn Sims. (*variegatus* Poir. test. Spr.) ♀. D. ex h. Gryph. 1837. Cfr. Agardh. Synops. gen. Lupini Lund. 1835. p. 21.
perennis L. ♀. D. 18 poll. alt. flor. purpurascens. Jun.
pilosus L. ☉. D. 1810 Retz. et ex h. Gryph. 1837. tripedalis fl. rubr.
polyphyllus Lindl. Ag. ♀. D. 2 uln. alt. intr. 1833. *Flora* 1835 p. 5; flor. cærul. mens. Jun. explic.
rivularis Lindl. Ag. ☉. ♀. D. 1834 et ex h. Par. 1837. bipedalis. flor. pallide cærul. vexilli medio albido. Jun.

succulentus Lindl. ☉. D. ex h. Gryph. et Paris. 1837.
Termis Forsk. ☉. D. ex hort. Kil. 1836. flor. alb.
varius L. ☉. 1807-1834. 1835. villosus W. ♂. ☉. ex h. Gr. 1836.

Luzula Cand. (*Juncus* Linn.).
albida Cand. ♀. D. 18 poll. alt.
campestris Cand. ♀. D.
nivea Cand. ♀. D. ex h. Par. 1837.
pediformis Cand. ♀. F. ex h. Paris. 1837.
vernalis Cand. (*Juncus pilosus* L.) ♀. D. spont.

Lychnis

chalconica L. ♀. D.
β. fl. plen.
Fulgens Fisch. ♀. D. bipedalis, fl. magn. coccin. Jun. explic.
læta Ait. ☉. D. 1815 et ex h. Kil. 1836.
mutabilis H. Gryph. ♀. ex h. Hafn. 1836.
pratensis Rafn. Fries (*dioica* L.) ♀. D.
silvestris Rafn. Fr. (*dioica* L.) ♀. D.

Lycium

afrum L. ♀. F.

- barbarum L. $\text{\textcircled{D}}$.
 carolinianum $\text{\textcircled{D}}$. D. tripedale
 et ultra; intr. 1833 ex h.
 Hamb.
 Treviranum $\text{\textcircled{D}}$. D. intr. 1834
 ex h. Hamb.
- Lycopersicon**
 cerasiforme $\text{\textcircled{C}}$. C. ex h. Pa-
 ris. 1837.
 esculentum Dunal. (*Solan. Ly-*
copersicum L.) $\text{\textcircled{C}}$. D.
 β . Humboldtii W.
 pyriforme Dun. Spr. (*Solanum*
Witbr.) $\text{\textcircled{C}}$. T. ex h. Pa-
 ris. 1837.
- Lycopodium**
 denticulatum L. $\text{\textcircled{D}}$. T.
- Lycopsis**
 arvensis L. (*Anchusa* Spr. Fr.)
 $\text{\textcircled{C}}$. D. spont.
 lutea Lam. (*Nonea* Cand. *An-*
chusa M. B. *Anchusa Ly-*
copsidis Retz. pugill. eadem
 videtur) $\text{\textcircled{C}}$. $\text{\textcircled{D}}$. D. 1810-
 1835; etiam ex h. Petrop.
 1836.
 rosea Lehm. $\text{\textcircled{C}}$. D.
 variegata L. (*Anchusa* Lehm.)
 $\text{\textcircled{C}}$. F. ex h. Hafn. 1837.
- Lycopus**
 exaltatus L. $\text{\textcircled{D}}$. D. 1813. herb.
 nostr.
 virginicus L. $\text{\textcircled{D}}$. D. 1826.
- Lyonia**
 paniculata Nutt. *Andromeda*
 L.) $\text{\textcircled{D}}$. D. in nov. arboret.
 1835; hodie 16 poll. alt.
- Lyonsia**
 straminea R. Br. $\text{\textcircled{D}}$. T. ex h.
 Kew. 1837. Ag.
- Lysimachia**
 ciliata L. $\text{\textcircled{D}}$. D. $1\frac{1}{2}$ uln. alt.
 flor. pallide flav. Aug.
 dubia Ait. $\text{\textcircled{D}}$. D. 1835 et ex
 h. Petrop. 1836.
 Ephemerum L. $\text{\textcircled{D}}$. $\text{\textcircled{D}}$. D. 20
 poll. alt. flor. alb. venis
 rubris, mens. Aug. explic.
 nemorum L. $\text{\textcircled{D}}$. D. 1826.
- Nummularia L. $\text{\textcircled{D}}$. D.
 punctata L. $\text{\textcircled{D}}$. D. flor. flav.
 quadrifolia L. $\text{\textcircled{D}}$. D. fl. flav.
 β . hirsuta Mx.
 verticillata Pall. $\text{\textcircled{D}}$. D. intr.
 1835 et ex h. Petr. 1836.
- Lythrum** (Fackelros).
 Hyssopifolia L. (*tenellum* Thb.)
 $\text{\textcircled{C}}$. D. flor. rubr.
 Salicaria L. $\text{\textcircled{D}}$. D.
 tomentosum Mill. $\text{\textcircled{D}}$. F. 1835.
 bipedal. flor. rubr. mens.
 Octob. explic.
 virgatum L. $\text{\textcircled{D}}$. D. flor. rubr.
- Madia**
 sativa Molin. Spr. (*mellosa* Jacqv.
viscosa Cav.) $\text{\textcircled{C}}$. D. 1808-
 1835 et ex h. Kil. 1836.
 27 pollic. alt. flor. flav.,
 Aug. explic.
 speciosa; $\text{\textcircled{C}}$. D.
- Magnolia**
 obovata Thb. (purpurea) $\text{\textcircled{D}}$. T.
 Umbrella Lam. (*tripetala* L.
frondosa Salisb.) $\text{\textcircled{D}}$. T.
 species, quæ in hort. nondum
 floruit. $\text{\textcircled{D}}$. T.
- Mahernia**
 pinnata L. (*Hermannia* L.) $\text{\textcircled{D}}$.
 T. 1814 herb. nostr.
- Maianthemum**
 bifolium Cand. (*Convallaria* L.)
 $\text{\textcircled{D}}$. D.
- Malachra**
 alceaefolia Jacqv. (*heptaphylla*
 Fisch.) $\text{\textcircled{C}}$. C. 15 poll. alt.
 fl. parvo, rubescent. inod.
 mens. Aug. explic.
- Malcolmia** (*Malcomia*).
 africana R. Br. (*Hesperis* L.)
 $\text{\textcircled{C}}$. D. 1810 et ex h. Pa-
 ris. 1837. flor. purpur.
 chia Cand. (*Cheiranth.* L. *He-*
speris Lam.) $\text{\textcircled{C}}$. D. ex hh.
 Kil. et Crac. 1837.

- littorea* R. Br. (*Ceiranth.* L.)
 ♂. D.
maritima R. Br. (*Cheiranth.* L.)
 ♂. D. 4-5-pollicaris, flor.
 rubris l. pallide violaceo-
 rubris; Jul.

Malope

- grandiflora* Hook. ♂. D. 1835.
 sesquipedal. flor. purpur.
 mens. Sept. explic.
trifida Cav. ♂. D. bipedalis,
 flor. purpur., Aug.

Malva (Kattost).

- Alcea* L. ♀. D. 1824 et iterum e convalli Fogelsång 1836 in hortum introd. 1 uln. alt. flor. Aug.
asperrima Jacqv. ♀. T. 1823.
brasiliensis Desr.; 1822, et ex h. Paris. 1837.
Capensis L. ♀. T. ex h. Crac. 1837.
caroliniana L. ♂. D. 1822. herbar.
crispa L. ♂. D. 2 uln. alt.
fragrans Jacqv. (*capensis* γ. L.) ♀. T. 1810 et ex h. Crac. 1837.
grossulariaefolia Cav. ♂. ♀. T. 1808. herbar. 1824.
maritima H. Gryph. ♀. T.
mauritanica Spr. (*mauritanica* L.) ♂. D.
miniata Cav. ♀. T. F. triped. flor. Jul.
morifolia Pollin.; ex h. Crac. 1836.
moschata L. ♀. D.
 β. fl. alb.
niceensis All. ♂. D. 1822—1826, etiam ex h. Paris. 1837.
parviflora L. ♂. D. ex hort. Paris. 1837. cur. Cl. Ag.
pulchella Bernh. ♂. T. ex h. Hafn. 1836.
purpurata; ex h. Paris. 1837.
Sherardiana L. Spr. ♀. T. sub nomin. *M. cymbalariaefolia* etiam ex h. Petr. 1836.
silvestris L. ♂. ♂. ♀. T.
Tournefortiana L. ♀. F. D. 1822. herbar.

- tricolor* H. Crac.; ex h. Crac. 1836.
verticillata L. ♂. D. 1808 herb. nostr.
 species.

Manulea

- alternifolia* Sternbg. ♂. C. ex h. Paris. 1837
cuneifolia Spr. ♀. C. sub nom. *Buchnera cuneifolia* Retz. cum syn. *B. oppositifolia*, in hort. Lund. 1806 culta, sed deinde profuga.
foetida W. Palmstruchia Retz. Pugill. *Buchnera* Andr.) ♂. F. intr. 1810; etiam ex h. Gryph. 1837.
violacea Link. ♂. C. intr. 1834 ex h. Kil.

Marica

- californica* Ker. v. *Sisyrinchium*.
coelestis Lehm. ♂. ♀. T. ex h. Kil. 1836.
striata Ker. v. *Sisyrinchium*.

Mariscus

- ovularis* Vahl. (*Schoenus umbellatus* Jacqv.) T. ex h. Paris. 1837.
umbellatus Vahl. (*Kyllingia* L.) ♀. C. 1810.

Marrubium

- astracanicum* Jacqv. (*affine* Hornem. *leonuroides* Desr.) ♀. F. 1826, et ex h. Petrop. 1836.
creticum Mill. Spr. ♀. D. 1826.
crispum L. ♀. F. 1810. 1826.
hispanicum L. (*rupestre* Bivon.) ♀. F. 1826 et ex h. Paris. 1837.
Jussieum; sub hoc nom. ex h. Petrop. 1836.
paniculatum Desr. Spr. ♀. D. ex h. Paris. 1837.
parviflorum; sub hoc nom. ex h. Petrop. 1836.

peregrinum L. 2. D. 1810 et
ex h. Paris. 1837.
propinquum H. Gryph. 2. ex
h. Petrop. 1836.
rugosum; sub hoc nom. ex h.
Paris. 1837.
supinum L. 2. F. 1826.
vulgare L. 2. D.

Martynia

perennis L. v. *Gloxinia maculata*.
proboscidea Glox. (annua L.)
⊙. C. 1808. herb. et ite-
rum ex h. Kil. 1836.

Matricaria

(Kamillblomma).

asteroides L. v. *Boltonia*.
Chamomilla L. ⊙. D. spont.
Parthenium L. v. *Pyrethrum*.

Matthiola (Cheiranthus L.).

annua L. ⊙. D.
incana R. Br. 2. F. D. flor.
simpl. var. color.
β. fl. plen. var. color.
fenestralis L. fil. 2. ♂. F.
tricuspidata R. Br. ⊙. D. ex
h. Paris. 1837.

Maurandia

antirrhiniflora Humb. 2. T.
1822. herb. nostr.
Barclayana Lindl. 2. C.
intr. 1836 ex h. Bonn.
semperflorens Jacq. (*Usteria*
scandens Cav.) 2. 2. T.
flor dilute violac.

Medicago

arborea L. 2. F. 2 uln. alt.
flor. flav.
caspica Fisch. ⊙. ex h. Hafn.
1827 et Kil. 1835.
ciliaris W. ⊙. D. 1815 et ex
h. Kil. 1836.
corymbosa Schmidt. ⊙. ex h.
Kil. 1836.
elegans Jacq. ⊙. D. intr. 1810;
etiam ex h. Paris. 1837.
falcata L. 2. D.
glutmosa M. B. 2. D. 1826.
laciniata Gouan. ⊙. D. 1812
et ex h. Kil. 1837.

lappacea Desr. ⊙. D. ex hort.
Kil. 1836.

lupulina L. ♂. D.

maculata W. ⊙. D. 1810.
1812 et ex h. Kil. 1835.

minima Desr. ⊙. D. 1810 et
ex h. Kil. 1836.

nigra W. ⊙. D. 1810. 1814
et ex h. Kil. 1836.

orbicularis Desr. (*polymorpha* α.
L.) ⊙. D. 1815 et ex h.
Kil. 1836.

pubescens Cand. ⊙. D. 1813,
et ex h. Hafn. 1827. 1836.

sativa L. 2. D.

β. fl. alb.

scutellata Lam. ⊙. D. ex h.
Kil. 1836.

Terebellum W. ⊙. D. ex h.
Hafn. 1827 et Kil. 1836.

tornata Desr. (*polymorpha* γ. L.)
⊙. D. 1811. 1815.

tuberculata W. ⊙. D. 1815
et ex h. Paris. 1837.

turbinata W. (*polymorph* δ. L.)
⊙. D. 1812.

uncinata W. ⊙. D. 1812.

Melaleuca

densa R. Br. 2. T.
diosmæfolia Andr. 2. T.
hypericifolia Sm. 2. T. ex h.
Crac. 1837.
linariæfolia Sm. (*Metrosideros*
hyssopifolia Cav.) 2. T.
species 2 ignot. e Nova Hol-
landia, haud determinan-
dæ, quia in horto nondum
floruerunt.

Melampodium

(*Wedelia* W.).

paludosum Kunth. v. *Dysodium*
divaricatum.

perfoliatum Kunth. ⊙. C. 1810
et 1837 bipedale et ultra.

Melananthera

hastata Mx. (*Bidens nivea* L.)

⊙. C. ex h. Paris. 1837
cur. Cl. J. Ag.

Melia

Azedarach L. ♀. T.
sempervirens Sw. ♀. T.

Melianthus

major L. ♀. T. F. 4 uln. alt.;
bene viget, sed rarissime
in horto floruit ex gr.
1827 mens. Maj.

Melica

altissima L. 2. D. fere tripe-
dalis; flor. Jun.

Bambini All. 2. D. ex h. Hafn.
1837.

ciliata L. 2. D. 1826.

pyramidalis Link. 2. D. 1819.
1826. et ex h. Par. 1837.

Melicope

ternata Forst. ♀. T. ex h. Kew.
1837, cur. Cl. J. Agardhi
intr.

Melilotus (Trifolii species Linn).

Besseriana; ⊙. D. ex h. Paris.
1837.

cærulea Desv. ⊙. D. bipedal.
et ultra.

cretica Desr. ⊙. D. ex h. Pa-
ris. 1837.

dentata Pers. ♂. 2. T. 1806
herb. etiam 1837.

diffusa Tren., ex h. Crac. 1837.

gracilis Cand. ⊙. D. T. ex h.
Crac. 1836 unipedalis.

hamosa M. B. ⊙. D. ex hort.
Paris. 1837.

indica Desv. ⊙. D. 1815 et
ex h. Paris. 1837.

italica Lam. ⊙. D. tripeda-
lis, flor. flav. Aug. explic.

officinalis Lam. ⊙. ♂. D. vix
tripedal.

palustris Spr. ♂. ex h. Crac.
1836.

parviflora Desf. H. Hafn. 1822.

rugulosa W. ♂. ⊙. D. ex h.
Crac. 1836.

sulcata Desf. H. Lips. ⊙. D.
ex h. Paris. 1837.

vulgaris W. ♂. D. fl. alb. sp.

Melissa

grandiflora L. 2. F. 1808 herb.
nostr.

officinalis L. (Citronmelissa.)
2. D.

β hirsuta Horn. 1826.

pyrenaica Jacqv. (Horminum L.)

2. D. 8 poll. alt. flor. vio-
laceo-rubr. mens. Jul. expl.

Menispermum

canadense L. ♀. D. vetus hort.
incola.

Mentha (Mynta).

arvensis L. 2. D.

cervina L. 2. D. F. ex hort.
Paris. 1837.

crispa L. (Krusmynta) 2. D.

crispata Schrö. 2. D. 1826.

macrostachya Tenor. 2. D. s.
hoc nomine commun., sed
dubia.

piperita L. (Pepparmynta) 2.
D. 1812.

sylvestris L. 2. D. 1826.

Mercurialis

annua L. ⊙. D.

ovata Sternb. 2. D. ex h. Pa-
ris. 1837 an distinct. a

M. perenni.

perennis L. 2. D.

tomentosa L. ♀. F. ex h. Pa-
ris. 1837.

Mesembryanthemum

aureum Ait. ♀. T.

barbatum L. ♀. T. flor. rubr.

coccineum Haw. ♀. T.

cordifolium L. 2. ♀. C. T. bi-
pedale. flor. rubr.

crassifolium L. ♀. T.

crystallinum L. (Is-plantan).

⊙. T. flor. albid. & roseis.

deltoides Haw. ♀. T. fl. rubr.

densum Haw. (barbatum γ. W.)
♀. T.

heteropetalum W. $\frac{1}{2}$ T.
 hispidum L. $\frac{1}{2}$ T.
 longum Haw. $\frac{1}{2}$ T.
 maximum Haw. $\frac{1}{2}$ T.
 pinnatifidum Horn. \odot T. 1822.
 flor. flav.
 pulverulentum Haw. $\frac{1}{2}$ T.
 rubricaulis Haw. $\frac{1}{2}$ T.
 tortuosum L. $\frac{1}{2}$ T.
 tricolor W. \odot T. ex h. Kil.
 1836.
 tuberosum L. $\frac{1}{2}$ T.
 verruculatum L. $\frac{1}{2}$ T.

Mespilus

acuminata Wall. $\frac{1}{2}$ D. 1835.
 1 uln. alt.
Amelanchier v. *Pyrus*.
 apiifolia Spr. $\frac{1}{2}$ D. in arbor.
 8 uln. alt.
 arbutifolia L. v. *Pyrus*.
 Azarolus Poir. (*Crataegus* L.)
 $\frac{1}{2}$ D. in nov. arbor. 6 uln.
 alt.
 coccinea Ehrh. (*Crataegus* L.) $\frac{1}{2}$ D.
 8 uln. alt.
 Cotoneaster L. (*Cotoneaster vul-*
garis Lindl. Fr.) $\frac{1}{2}$ D.
 cuneifolia Ehrh. (*Crataegus pun-*
ctata Ait. W.) $\frac{1}{2}$ D. 13
 uln. alt.
 frigida; $\frac{1}{2}$ F., sub nom. *Coto-*
neast. frigida ex h. Chis-
 wich Angliæ, cur. Clar.
 Agardh 1837, miss.
 germanica L. $\frac{1}{2}$ D. intr. 1798;
 hodie 10 uln. alt. fl. alb.
 β . sine nucleo.
 grandiflora Sm. $\frac{1}{2}$ D. in nov.
 arboret. $3\frac{1}{2}$ uln. alt. se-
 min. nigr.
 grossulariæfolia. $\frac{1}{2}$ D. in nov.
 arbor. $3\frac{1}{2}$ uln. alt.
 lucida Ehrh. (*Crataegus Crus Gallæ*
 Ait.) $\frac{1}{2}$ D.
 microphylla H. Hamb. $\frac{1}{2}$ F.
 monogyna Ehrh. (*Crat.* Jacq.)
 $\frac{1}{2}$ D. 17 uln. alt. fl. alb.
 Jun.
 nigra W. (*Crataegus* Kit.) $\frac{1}{2}$ D.
 4 uln. alt.

Oxyacantha Gärtm. (*Crataeg.*
 L.) $\frac{1}{2}$ D.
 β . fl. rubicundo.
Pyracantha L. $\frac{1}{2}$ D. (Cfr. *Flo-*
ra 1836 p. 12.) 3 uln. alt.
 pyrifolia W. $\frac{1}{2}$ D. 12 uln. alt.
 rotundifolia Ehrh. (*glandulosa*
 W.) $\frac{1}{2}$ D. ex h. Chisw.
 1837.
 species.

Messerschmidia

fruticosa L. v. *Tournefortia*.

Metrosideros

citrinus $\frac{1}{2}$ T. ex h. Crae. 1837.
hyssopifolius Cav. v. *Molaleuca*
linariifolia.
 pinifolius Wendl. Spr. (*Calli-*
stemon Hortulan.) $\frac{1}{2}$ T.

Meum

athamanticum Jacq. (*Ethusa*
Meum Ait. *Athamantis* *Me-*
um L.) $\frac{1}{2}$ D.
 fœniculum Spr. (*Anethum* L.)
 $\frac{1}{2}$ D.
 sibiricum Spr. $\frac{1}{2}$ D. ex hort.
 Crae. 1837.

Michauxia

lævigata Vent.; T. ex h. Petr.
 1836.

Microloma

lineare R. Br. (*Ceropegia tenui-*
flora Thb. *Periploca tenui-*
folia L.) $\frac{1}{2}$ C.

Micropus

supinus L. \odot D. intr. 1810
 deinde etiam ex h. Kil.
 1835. miss.

Microtea

debilis Sw. \odot T. 1834, et ex
 h. Hafn. 1836.

Milium

latifolium Cav. v. *Paspalum sto-*
loniferum.
 lendigerum Schreb. (*Gastridium*
australe Spr.) ex h. Hafn.
 1837.
 paradoxum L. $\frac{1}{2}$ F. D. in h.
 1806 herb. nostr.

racemosum Jacq. v. *Paspal. stolonifer*.

Milleria

Contrayerva Cav. v. *Flaveria*.

Mimosa

distachya Vent. v. *Acacia lo-phanta*.

floribunda Vent. v. *Acacia*.

pubica L. ♀. C. 1810. 1835.

hodie unipedalis.

sensitiva L. ♀. C. 1806-1810.

Mimulus

guttatus Fisch. ♂. 4.? F. 1826; et ex h. Gr. 1837.

luteus L. 4. F. 1822-1835; et ex h. Gryph. 1836. 12 poll. alt.

moschatus Dougl. Wilbr. ♂. 4. T. ex h. Londin. 1834 et Gryph. 1836. 6 pollic.

alt. flor. flav.

propinquus Lindl. ♂.; ex h. Gryph. 1836.

quinquevulnerus Link. H. Lips. 4. intr. ex h. Bonn. 1836.

ringens L. 4. F. D. ex hort. Paris. 1837.

roseus H. Hamb. 4. F. ex h. Londin. 1834 et Bonn. 1836

variegatus H. Paris. 4. intr. 1834 ex h. Kil.

species.

Minuartia

dichotoma L. ♂. D. ex hort. Hafn. 1834 et iterum 1837.

montana L. ♂. D. ex h. Hafn. 1834. 1837.

Mirabilis

Jalappa L. 4. ♂. D. fl. ruhr., luteo, variegat. cet.

longiflora L. 4. ♂. F. D. 1834 et ex h. Hafn. 1836.

Mitella

diphylla L. 4. D.

grandiflora Pursh. (*Tellima* Dougl.) 4. D. 26 pollic. alt. flor. lutescent., Maj. β. fl. rubicund.

Mollugo

dichotoma Schrk. v. *Pharnaceum verticillatum*.

pentaphylla L. v. *Pharnaceum*.

Moluccella

frutescens L. ♀. T. ex h. Paris. 1837 cur. Cl. Ag.

Momordica

Balsamina L. ♂. C. unipedalis; flor. sulphur. Aug. explicat.

Charantia L. ♂. C. intr. 1837 ex h. Paris.

Elaterium L. ♂. D. in hort. jam spont. dissemin.

Monarda

altissima W. 4. D. bipedalis flor. Jun.

clinopodia Horn. (*glabra* Lam.) 4. D. bipedalis et ultra.

ciliata L. 4. F. ex h. Hafn. 1836.

Kalmiana Pursh. 4. F. 1833 et ex h. Hafn. 1836.

fistulosa L. (*media* W.) 4. D. 1826.

lilacina Wendlrth. 4. ex hort. Gryph. 1837.

mollis W. 4. D. unipedalis; flor. lilacinis. Aug. expl.

oblongata Ait. 4. D. bipedalis. flor. Aug.

purpurea Pursh. 4. F. ex h. Gryph. 1837.

Monsonia

ovata Cav. (*Geran. emarginatum* L.) ♂. T. ex h. Par. 1837.

pilosa W. 4. T. intr. 1837 ex h. Paris.

Moræa

pavonia Ker. v. *Iris*.

Morgania

lucida Spr. v. *Gratiola*.

Morus (Mulbärsträd).

alba L. ♀. D. 9 uln. alt.; in Paradislyckan.

Calcar Galli $\frac{1}{2}$. T. sub hoc
nom. ex h. Kew. 1837.
Mbrettiana H. Hamb. $\frac{1}{2}$. D.;
variet. *M. albæ* videtur. 2
uln. alt.
nigra L. $\frac{1}{2}$. D.
papyrifera L. v. *Broussonetia*.
rubra L. $\frac{1}{2}$. D. 14 uln. alt.

Musa (Bananas).
paradisiaca L. 2. C. in hort.
floruit 1823.

Muscari (*Hyacinthus* L.)
botryoides W. 2. D.
ciliatum Ker. 2. F. ex h. Petr.
1836.
comosum W. 2. D. 8, demum
20, poll. alt.; intr. 1808.
 β . *monstrosum* Hornem.
moschatum W. 2. D. 8 pol-
lic. alt.
racemosum W. 2. D.

Mutisia
ilicifolia Cav. $\frac{1}{2}$. T. ex hort.
Petr. 1836.

Myagrurn
dentatum W. v. *Camelina*.
paniculatum L. v. *Neslia*.
rugosum L. v. *Cakile*.
sativum L. v. *Camelina*.

Myosotis
arvensis L. Fr. \odot D.
collina Reich. Fr. \odot D.
Lappula L. v. *Echinosperrnum*.
lithospermifolia Horn. 2. D.
6 poll. alt.; intr. 1832 ex
h. Gryph.
stricta Link. Fr. \odot D. 5 pol-
lic. alt.
versicolor Sm. Fr. \odot D. 4-5
poll. alt.

Myosurus
minimus L. \odot D.

Myrrhis (*Cherophylli* spe-
cies L.).
aromatica Spr. 2. D. ex hort.
Paris. 1837.
aurea Spr. 2. D. ex h. Paris.
1837.
canadensis Moris. 2. D. 1826,
et ex h. Crac. 1837.
colorata Spr. 2. D. ex h. Hafn.
1837.
hirsuta Spr. 2. D.
humilis R. et Sch.; 2. ex h.
Crac. 1837.
odorata Scop. (*Scandix* L. *Ce-
refolium hispanicum* Phar-
macop.) 2. D.
temula Gärtn. et Spr. 2. D.
Myrsine
africana L. $\frac{1}{2}$. T. pedalis; fl.
parvis, rubescenti albis.
variabilis R. Br. $\frac{1}{2}$. T. ex h.
Kew. 1837.

Myrtus (Mirten).
australis Spr. (*Eugenia* Wendl.)
 $\frac{1}{2}$. C. T. 1 $\frac{1}{2}$ uln. alt.
communis L. $\frac{1}{2}$. F.
 β . *angustifolia*.
 γ . *macrophylla*.
 δ . fl. pleno.
Vahlia Spr. $\frac{1}{2}$. T. sub nomin.
Eugenia trinervia Vahl ex
h. Kew. 1837.

Napæa.
dioica L. v. *Sida*.
levis L. v. *Sida Napæa*.

Narcissus (Narciss).
bicolor L. 2. D.
biflorus Curt. 2. D.
incomparabilis Curt. 2. D.
unipedalis.
 β . fl. plen.
Jonquilla L. 2. T.
poëticus L. 2. D.
 β . fl. plen.
Pseudonarcissus L. 2. D.
 β . fl. plen.
Tazetta L. 2. T. F. fl. flav.
 β . fl. alb.

Nasturtium
Armoracia Fr. v. *Cochlearia*.

- atrovirens* v. *Sisymbrium*.
clandestinum Spr. ☉. C.
globosum Trin. ☉.; ex hort.
 Petr. 1836.
microspermum Cand. C. ex h.
 Petrop. 1836.

Nauenburgia

- trinervata* W. v. *Flaveria repanda*.

Navaretia

- involuta* R. et P. (*Collomia eryngioides* ex h. Hafniæ)
 C. ☉. ? ex h. Par. 1837.
 cur. *Ag.* jam flor. in Cald.
 20 Oct. 1837; flor. sub-
 capit. rubr.

Nemesia

- chamædryfolia* Vent. (*Antirrh. macrocarpum* Ait.) ♀. T.
 ex h. Paris. 1837.

Nemophila

- atomaria* F et M. ☉. D. ex
 h. Kil. 1836.
aurita H. Hafn. ☉. D. 1835.
 hodie hort. deest.
insignis Dougl. ☉. D. ex h.
 Hafn. 1836.
phacelioides Bart. ☉. 2. D.
 ex h. Hafn. 1834 et Par.
 1837.
pedunculata Dougl. ☉. D. ex
 h. Hafn. 1836.

Neottia vide *Listera*.

Nepenthes

- destillatoria* L. 2. C. intr.
 1834; et iterum 1837 cur.
 Cl. J. *Ag.*

Nepeta

- Cataria* L. 2. D.
cærulea Ait. 2. D. 1826.
grandiflora M. B. (*colorata* W.)
 2. D. 1826. 1835. flor.
 cærul.
incana Ait. 2. ♂. D. ex h.
 Gryph. 1836.
italica L. 2. D. flor. alb.
 Meyeri; sub hoc nom. ex h.
 Petr. 1836.

- multifida* L. 2. D. ex h. Par.
 1837.

- Mussini* Henck. 2. F. 1826,
 et ex h. Paris. 1837.

- Nepetella* L. (*amethystina* Poir.?)
 2. D. 1826, et ex h. Par.
 1837.

- nuda* L. 2. D. ex h. Gryph.
 1836 a Cel. *Hornschuch*. fl.
 albis.

- pannonica* L. 2. D. 1810.
 1826.

- reticulata* Desf. 2. D. F. ex
 h. Paris. 1837.

- sibirica* M. B. (*Teucrium* L.)
 2. F. D. bipedal. fl. vio-
 laceo-cærul. Jun.

- teucrioides* Lam. s. hoc nom.
 ex h. Paris. 1837.

- tuberosa* L. 2. F. in h. 1810.
 herb. nostr.

- ucranica* L. 2. D. 1826, et ex
 h. Gryph. 1837.

- violacea* L. 2. D. 1810. 1826,
 et ex h. Gr. 1837. herb.
 cyanescens, flor. rubicund.

Nerium

- odorum* Ait. ♀. C. nuper. intr.
 cur. Cl. *Stenkula*.
 β. fl. plen.

- Oleander* L. ♀. T. F.
 β. album novum H. Hamb.
 γ. cardinale.
 δ. carneum.

Nesæa

- salicifolia* Kunth. (*Heimia* Link.)
 ♀. C. ex h. Paris. 1837.
 cur. Cl. *Ag.*

Neslia

- paniculata* Desv. (*Myagrum* L.
Vogelia Horn.) ☉. D. tri-
 pedalis.

Nevrada

- procumbens* L. ☉. T.

Nicandra

- physaloides* Gärt. (*Atropa* L.)
 ☉. D. 3-pedalis; flor. di-
 lute cærul. inod.; Aug.

Nicotiana (Tobak).

- acuminata* Hook. H. Lips. ☉.
 D. 1834. quadripedal.
alata Link. et Otto. ☉. ex h.
 Bonn. et Crac. 1836.

coccinea H. Crac. ☉, ex h. Crac. 1837.

fruticosa L. ☉. *h.* C.

glauca Grah. *h.* C. ex h. Bonn. 1834 et Paris. 1837.

gigantea H. Crac.; ☉. F. 1822 et ex h. Crac. 1837.

humilis H. Crac. Link. ☉. ex h. Crac. 1837.

Langsdorffii Nces. ☉. F.; 1822 ex h. Gryph. 1836.

Lehmanni H. Kil. Ag. ☉.; ex h. Hamb. 1824 et Hafn. 1837.

macrophylla Spr. Cand. ☉. D. 1819 et ex h. Hafn. 1836.

multivalvis Lindl. ☉. D. ex h. Kil. 1836. flor. alb.

micrantha H. Kil. Dougl. ☉. D. ex hh. Kil. et Paris. 1837. pedalis, florib. sordide albis.

nyctaginiiflora Lehm. v. *Petunia*.

noctiflora Hook. ☉. ex hort. Gr. 1833 et Paris. 1837.

odorata; ☉. D. 1837.

petiolata Ag.; ex hort. Hafn. 1837.

quadrivalvis Pursh. ☉.; 1822 et ex h. Hafn. 1836. flor. albis.

rustica L. ☉. D. flor. vire-scenti albis.

sanguinea H. Gryph. ☉.; ex h. Hafn. 1836.

svavecolens Lehm. ☉. D. ex h. Kil. 1836. floribus alb., nocte svavecolentibus.

Tabacum L. ☉. D.

versicolor H. Gryph. ☉.; ex h. Kil. 1836.

vinciflora Lag. ☉.; ex h. Kil. 1836.

Nigella

damascena L. ☉. D.

h. *coarctata* Hortulan.

foeniculacea Cand. (*segetalis* Spr.) ☉. D.

Nolana

atriplicifolia Don. H. Crac. ☉. D. ex h. Hafn. 1837.

grandiflora Hook. Hortulan. ☉. ex h. Hafn. 1834, Kil. 1836.

prostrata L. ☉. T. flor. magn. cærul.

Nonea

arvensis Cand. v. *Lycopsis*.

flavescens. ☉. D. sub hoc nom. ex hh. Petr. 1836, et Paris. 1837.

lutea Cand. v. *Lycopsis*.

Notelæa

punctata R. Br. *h.* T. ex h. Chiswich Augl. 1837, our.

Cl. Ag.

Nymphæa (Näckros).

cærulea Savign. *h.* F. in hort. olim culta.

Obeliscaria

columnaris Cand. v. *Rudbeckia*.

pinnata Cass. v. *Rudbeckia*.

Ocimum

americanum L. ☉. C. 1810.

Basilicum L. ☉. T.

minimum L. ☉. C. 1810.

Oedera

prolifera Thb. Spr. (*Buphthalmum capense* L.) *h.* T. intr.

1833 ex h. Hafn.

Oenanthe

australis H. Hafn. *h.* D.

crocata L. *h.* D. 1826.

pencedanifolia Poll. *h.* D. 1810. 1826.

silaisfolia M. B. *h.* D. ex h. Paris. 1837.

pimpinelloides L. *h.* D. ex h. Paris. 1837.

Oenothera

acaulis Cav. ☉.; ex h. Kil. 1836.

amoena Lehm. ☉. D. ex h. Paris 1837.
biennis L. ♂. ☉. in graminos. horti jam spont. dissemin. capensis H. Lips.; ex h. Hafn. 1837.
chinensis (chilensis) H. Crac.; ex hh. Crac. et Parisiens. 1837.
concinna Don. ex h. Petrop. 1836.
cruciata Nutt. ♂.; ex h. Par. 1837.
decumbens Dougl. ☉. ♂. D. 1834 et ex h. Par. 1837. caule sesquipedal. fl. parv. purp.
densiflora Lindl. ☉. ♂. D. 1834 et ex h. Hafn. 1837. caul. unipedal. flor. pallide rubr.
dentata Cav.; 1824 et ex h. Crac. 1836.
erosa Hornem. Spr.; ex hort. Paris. 1837.
Fraseri Pursh ♀. D. ex hort. Kil. 1836.
fruticosa L. ♀. D. fere bipedalis, flor. flav.
gauroides Horn. (*biennis* var. Spr.) ex h. Hafn. 1836.
grandiflora Ait. ♂. F. D. 1819 et ex h. Hafn. 1835. caul. 3-pedali et ultra in hort.
Lindleyana (Lindleyi) Hook. ☉. ♂. D. 1834--1837. caul. bipedal. petal. ros. macul. centrali purpur.
longiflora Jacqv. ☉. ♂. D. F. ex h. Hafn. 1837.
media Link.; D. ex h. Hafn. 1837.
micrantha Hornem. (*hirta* Link.) ♀. 1822.
mollissima L. ☉. D. ex hort. Paris. 1837.
nervosa Horn.; ex h. Hafniae 1837.
nocturna Jacqv. (*striata* Ledeb.) ♂. ☉. D. F. ex h. Kil. 1836.
odorata Jacqv. ♀. (☉.?) F. 1810 et ex h. Hafn. 1837.

purpurea Cart. ☉. D. caul. fere bipedali.
quadrivulnera Dougl. ☉. D. ex h. Paris. 1837.
rhizocarpa Spr. ☉. ♀. D. 1834 et ex h. Kil. 1836 pedunc. demum pedal. flor. flav.
Romanzovii Ledeb. ☉. ♂. D. 1822 et ex h. Par. 1837. bipedalis, fl. cœrulescenti violac.
rosea Ait. ♀. ♂. F. 1808. 1834; nec hodie.
roseo-alba Bernh. ☉. D. ex h. Hafn. 1837.
rubicunda; ☉. D. ex h. Par. 1837.
salicifolia Desf.; ex h. Berol. 1824 et Paris. 1837.
serotina H. Hafn.; ex h. Hafn. 1836.
spectabilis Horn. ♂. D. ex h. Hafn. 1837.
stricta Ledeb. ☉. D. ex hort. Berol. 1824 et Hafn. 1837.
taraxacifolia Reichb. ♀. F. D. ex h. Paris. 1837.
tenuifolia Cav. T. ex h. Par. 1837.
tetraptera Cav. ♀. (☉. Willbr.) F. 1810 et ex h. Hafn. 1837.
viminea H. Hafn.; D. ex h. Hafn. 1837.

Oldenlandia

biflora L. (*Hedyotis* Horn.) ☉. D. ex h. Hafn. 1836.
diffusa Roxb. v. *Hedyotis*.

Olea (Oljetråd).

europæa L. ♀. T. F.

Oncidium

Papilio Lindl. ♀. C. ex hort. Edinb. 1837. cur. Clar. J. Ag.

Onobrychis

Buxbaumiana Desv. ♀. F. ex h. Paris. 1837.

- Caput Galli** Lam. (*Hedysarum* L.) 2. (Spr.) ⊙. D. 1812. 1835, et ex h. Par. 1837.
- crinita** Spr. 2. T. ex h. Petr. 1836.
- Crista Galli** Lam. (*Hedysarum* L.) ⊙. ex h. Hafn. 1834 Paris. 1837.
- petraea** Spr. (*Hedysarum* M. B. *Hed. carpathicum* W.) 2. F. ex h. Paris. 1837.
- sativa** Lam. (*Hedysar. Onobrychis* L.) 2. D. 1809.
- saxatilis** Lam. *Hedysar. L.*) 2. F. ex h. Paris. 1837.
- Onoclea**
sensibilis L. 2. D.
- Ononis**
alopecurioides L. ⊙. D. intr. 1810; etiam ex h. Paris. 1837.
- arvensis** Retz. (*hircina* Jacq.) 2. D. 1826.
- mitissima** L. ⊙. ? F. intr. 1837 ex h. Paris.
- parviflora**; ⊙. F. ex h. Paris. 1837.
- pingvis** Spr. (*On. Nutrix* L.) 2. F. intr. 1837 ex hort. Paris.
- repens** L. 2. D. sub hoc nom. in hort. diu culta, bipedalis.
- spinosa** L. 2. D.
- variegata** L.; ex h. Par. 1837.
- Onopordum** (Sempertin).
Acanthum L. ♂. D.
- macracanthum** Schousb. ♂. D. 1835.
- tauricum** W. ♂. 1835.
- Onosma**
sericeum M. B. 2. ex h. Petropol. 1836.
- Ophrys**
ovata L. v. *Listera*.
- Orchis**
mascula L. 2. D.
- militaris* L. 2. D. 1826. 1829.
- Morio* L. 2. D. 1826.
- sambucina* L. 2. D. fl. rubr. aut flav. intr. 1836.
- ustulata* L. 2. D. (Obs. *Orchidis* species in hort. plantatæ tantum per unum alterumve annum viguerunt).
- Origanum**
creticum L. 2. D. 1826.
- heracleoticum* L. 2. D. 1811. herb. nostr.
- Majorana** L. (Mejram) ⊙. D.
- smyrnaeum** L. 2. T. 1815 hort. incolæ, jam exsul.
- vulgare** L. 2. D.
- Ornithogalum**
exscapum Tenor. 2.
- luteum** L. 2. D.
- minimum** L. 2. D. in Lunda-gård spont.
- narbonense** L. 2. D. 33 pell. alt. flor. alb.
- nutans** L. 2. D. in hort. jam plurib. locis spontan. et copiosum, ægre pellendum.
- pyramidale* L. 2. D.
- pyrenaicum* L. 2. D. 1822. 1826.
- spathaceum* Hayn. 2. D. 1826.
- Ornithopus**
compressus L. ⊙. D. 1810 herb. nostr.
- creticus* v. *Coronilla*.
- glauca* v. *Coronilla*.
- scorpioides* L. ⊙. D. 1822.
- Orobus**
angustifolius L. 2. D. 8 pollic. alt.
- lathyroides* L. 2. D. 24-26 pollic. alt.
- niger** L. 2. D. culta planta bi-tripedalis.
- tuberosus** L. 2. D.
- variegatus** Tenor. 2. F. intr. 1824 ex hort. Berol.; vix nisi variet. *Orob. verni*. bipedalis et ultra.
- vernus* L. 2. D. bipedalis.
- Oryza** (Ris).
sativa L. ⊙. C.
- Osmunda**
regalis L. 2. D.
- Spicant.* L. v. *Blechnum*.
- Struthiopteris* v. *Struthiopt. europæa*.

Osteospermum

moniliferum; $\frac{1}{2}$. C. 1819; etiam
ex h. Paris. 1837.

pinnatifidum; $\frac{1}{2}$. T. sub h. nom.
in h. olim cult.

Ostrya

vulgaris W. (*Carpinus Ostrya* L.)

$\frac{1}{2}$. D. olim in hort. cult.,
jam profuga.

Oxalis

cernua Thb. $\frac{1}{4}$. T. olim horti
incola, hodie exsul.

crassicaulis Zuccar. $\frac{1}{4}$. F. in
hort. botanicum introd.
1833 — Sub nom. *Oxal.*
arakatscha l. *aracacha* in
hortis quoque colitur. Cfr.
Flora 1835. p. 4-5.

incarnata L. $\frac{1}{4}$. T. ante plures
annos in horto culta, ibi
diutius non exstat.

rosea Jacqv. \odot . C. ex hh. Bonn.
et Hafn. 1836.

stricta L. \odot . $\frac{1}{4}$. D. in horto
neglecta sponte propaga-
tur; fere vitium.

tetraphylla Cav. $\frac{1}{4}$. F. intr.
1834 ex h. Kil.

violacea L. $\frac{1}{4}$. F. olim hort., in-
cola, jam exstincta.

Pæonia (Pion).

albiflora Pall. $\frac{1}{4}$. D. caul. bi-
pedali et ultra.

anomala L. $\frac{1}{4}$. D. 1826 flor.
roseis.

arietina Anders. $\frac{1}{4}$. D. ex h.
Paris. 1837.

corallina Retz. $\frac{1}{4}$. D. tripeda-
lis, flor. rubr.

humilis Retz. $\frac{1}{4}$. D. bipedalis.
flor. rubr.

β . fl. simpl. pallid.

hybrida Horn. $\frac{1}{4}$. D. (vix dif-
fert a *P. tenuifolia*); bipe-
dalis.

β . flor. major. alb.

Mutan Sm. (*officinalis* Thb. non
L.) $\frac{1}{2}$. F.

officinalis L. $\frac{1}{4}$. D. adsunt pl.
variet. flor. plen., color.
rubr., alb., carneo.

tenuifolia L. $\frac{1}{4}$. D. bipedalis
et ultra, flor. purpur.

Palavia

malvæfolia Cav. (*Malope parvi-
flora* Herit.) \odot . C. 1808
herb. nostr., etiam ex h.
Kil. 1836.

Paliurus

australis Gärtn. Spr. (*aculeatus*
Lam. *Rhamnus Paliurus* L.
Zizyphus Paliur. W. Horn.)
 $\frac{1}{2}$. F. ex h. Hamb. 1834.

Palmstruckia

foetida Retz. v. *Manulea*.

Pancratium

littorale $\frac{1}{4}$. C. ex h. Crac. 1837
miss.

Panicum (Hirs).

ægyptiacum Retz. (*Digitaria*
W.) olim in hort. cult.
secund. specim. in herb.
nostr.

attenuatum W. \odot . D. ex h.
Crac. 1837.

capillare L. \odot . D. 1810 et ex
h. Paris. 1837.

colonum L. \odot . C. 1808 herb.
nostr.

coloratum L. $\frac{1}{4}$. C. 1806, et
ex h. Paris. 1837.

Crus Galli Spr. (*Pan. Crus Corvi*
L.) \odot . D. sesquipedal.

echinatum W. (*Orthopogon* Spr.)
 \odot . D.

erythrospermum Horn. (*Setaria*
Spr.) \odot . D.

filiforme L. (*Digitaria* Spr.);
1809 herb. nostr.

italicum L. (*Setaria* Spr. *Pan.*
germunicum Baub.) \odot . D.
3-pedal. et ultra.

miliaceum L. \odot . D. var. se-
mine nigr., luteo cet. bi-
pedal. et ultra.

- proliferum Lam. ☉. D. ex h. Paris. 1837.
 verticillatum L. (*Setaria* P. B.) ☉. D.
violaceum Lam. v. *Cenchrus setosus*.
 virgatum L. 2. C. ex h. Paris. 1837.
- Papaver (Vallmo).**
Argemone L. ☉. D. ped. fl. rubr.
cambricum L. (*Stylophorum* Spr. *Meconopsis* Vig.) 2. D. ped. et ultra. flor. flav.
caucasicum M. B. ☉. 2. D. 1824, et ex h. Gr. 1837.
croceum Ledeb. ☉. ♂. D. ex h. Paris. 1837.
dubium L. ☉. D. flor. rubr.
floribundum Desf. ☉. ex hort. Gryph. 1837.
hispidum Desf. ☉. ♂. D. intr. 1833 ex h. Bonn. biped. et ultra., florib. rubicund.
laevigatum M. B. ☉. D. 1834. 1835.
nudicaule L. ♂. 2. D. sesquipedal., flor. aurantiacis.
orientale L. (*bracteatum* Lindl.) 2. D. bipedal. flor. magn. cinnabarinis.
Rhoeas L. ☉. D. fl. color. var. β. fl. plen. et color. var.
somniferum L. ☉. D. fl. col. var.
- Parietaria**
judaica L. 2. D. 1822. 1826.
lusitanica L.; ex h. Hafniae 1837.
microphylla L. (*Urtica* Sw.); ex h. Hafn. 1837.
officinalis L. 2. D.
- Paronychia**
verticillata; ex h. Paris. 1837.
- Paspalum**
stoloniferum Bosc. (*Milium latifolium* Cav. *racemosum* Jacq.) ☉. D. ex h. Paris. 1837.
- tenellum* W. (*elegans* Spr.) 2. D. ex h. Hamb. 1824 et Paris. 1837.
- Passerina**
fliformis L. ♂. T.
sericea L. v. *Gnidia*.
- Passiflora**
 (Passionsblomme).
alata Ait. ♂. C. ex h. Paris. 1837.
albida Ker. ♂. C.
caerulea L. ♂. T.
capsularis L. ♂. C. ex h. Kil. 1835.
gracilis Jacq. ♂. C. 1834. 1835.
Londoniana; ♂. C. etiam sub hoc nom. ex hort. Edinb. 1837. cur. Cl. Ag.
princeps H. Hamb. ♂. C.
Vespertilio L. (*discolor* Link.) ♂. C.
- Pastinaca**
 (Palsternacka).
Anethum Spr. v. *Anethum graveolens*.
sativa L. ♂. D.
- Patrinia**
rupestris Juss. (*Valeriana sibirica* W.) 2. D. ex h. Paris. 1837.
- Paulinia**
species. ♂. C. in hort. nondum floruit.
- Pavia**
rubra Lam. v. *Aesculus Pavia*.
- Pectis**
discoidea Horn. v. *Gnaphal. leyserioides*.
- Peganum**
Harmala L. 2. D. ante plures annos in horto cult. ex h. Paris. 1822 intr.
- Pelargonium** (*Geranii* sp. Linn.).
alchemilloides Ait. 2. T.
australe W. ♂. T. 1810.
capitatum Ait. ♂. T.
coriandrifolium Ait. 2. T. 1810, et ex h. Par. 1837.
cucullatum Ait. ♂. T. biped. intr. 1822 ex h. Paris.

glutinosa Ait. $\frac{1}{2}$. T. 1835.
tripedale.

graveolens Herit. $\frac{1}{2}$. T. fere
3-pedale.

grossularioides Ait. $\frac{1}{4}$. T. 1807
in h. cult.

inodorum W. $\frac{1}{4}$. \odot . T. 1806.
1810.

inquinans Ait. $\frac{1}{2}$. T. 1812.

lacerum Jacqv. $\frac{1}{4}$. F. 1809.

malvæfolium Jacqv. (*laterisium*
W.) $\frac{1}{2}$. T. bipedal.

macranthum Sweet. (*speciosum*
Spr.) $\frac{1}{2}$. T. tripedal.

multicaule Jacqv. $\frac{1}{4}$. T. ex h.
Paris. 1837.

odoratissimum Ait. $\frac{1}{4}$. T.

peltatum Ait. $\frac{1}{2}$. T. 1810, et
iterum intr. 1835. biped.

Radula Herit. $\frac{1}{2}$. T. fere 2
uln. alt.

viscosum H. Hamb. $\frac{1}{2}$. T.

zonale Ait. $\frac{1}{2}$. T. bipedale et
ultra.

β . marginatum.
species.

Peltaria

alliacea L. $\frac{1}{4}$. D. sesquipedal.
fl. Jun. flor. alb.

Pennisetum

cenchrroides Rich. v. *Cenchrus*
ciliaris.

setosum Rich. v. *Cenchrus*.

Penthorum

sedoides L. $\frac{1}{4}$. T. flor. albis;
intr. 1835.

Pentstemon

(*Pentastemon*).

atropurpureum Dougl. $\frac{1}{4}$. T. F.
Flora 1836. p. 12.

campanulatum W. (*Chelone cam-*
panulata Cav.) $\frac{1}{4}$. T. F.

confertum Dougl. $\frac{1}{4}$. F. T. ex
h. Paris. 1837.

diffusum H. Hamb. $\frac{1}{4}$. T. F.
ex hort. Londin. 1834 et

Paris. 1837. sesquipedal.
flor. rubr.

Digitale Nutt. v. *Chelone*.

gracile Spr. v. *Chelone*.

ovatum Dougl. $\frac{1}{4}$. F. 1834, et
ex h. Paris. 1837.

pubescens Ait. (*Chelone Pent-*
stemon L.) $\frac{1}{4}$. D. 1827.

pulchellum H. Hamb. $\frac{1}{4}$. F.

roseum H. Hamb. $\frac{1}{4}$. T. F.
Flora 1836. p. 12; intr.

1834. caul. bipedal. flor.
rubro-viofac.

Scouleri H. Hamb. $\frac{1}{4}$. T. ex
h. Chiswich 1837.

speciosum H. Hamb. $\frac{1}{4}$. T. ex
h. Chiswich 1837.

venustum H. Hamb. $\frac{1}{4}$. T. intr.
1834 ex h. Londin.; dein-
de 1836. ex h. Bonn.

Peperomia

inæqualifolia R. et P. v. *Piper*.

Peplis

Portula L. \odot . F.

Perilla

ocymoides L. \odot . C.

Periploca

græca L. $\frac{1}{2}$. T. F. etiam sub
Dio leniores hiemes per-
ferre videtur.

Petiveria

alliacea L. $\frac{1}{2}$. C. 1819 secund.
herb. nostr.

Petunia

nyctaginiflora Juss. Spr. (*Ni-*
cotiana Lehm.) \odot . $\frac{1}{4}$. C.
ex h. Berol. 1824 et Gr.
1837.

Peucedanum

album Horn. $\frac{1}{4}$. D. ex h. Pa-
ris. 1837.

Silva L. v. *Cnidium*.

tenuifolium Thb. $\frac{1}{4}$. T. ex h.
Paris. 1837.

species.

Phaca

alpina Jacqv. $\frac{1}{4}$. D. bipedalis.
frigida L. Spr. $\frac{1}{4}$. F. ex hort.

Petr. 1836.

membranacea Fisch. 2. D. ex h. Hafn. 1826.

Phacelia

circinata Jacq. (*Heliotrop. pinnatum* Vahl. *Hydrophyllum magellanicum* Lam.) 2. ♂.

F. ex h. Paris. 1837.

tanacetifolia Benth. T. ex h. Paris. 1837.

Phalangium

Pendulum Red. v. *Arthropodium*. species.

Phalaris (Flen).

arundinacea β. *picta* L. 2. D.

bulbosa Cav. 2. D. 1812. 1826.

canariensis L. ♂. D. bi-quadripedal.

paradoxa L. ♂. D. bipedal. et ultra.

utriculata L. v. *Alopecurus*.

Pharnaceum

Cerviana L. ♂. D. ex h. Par. 1837.

pentaphyllum Spr. (*Mollugo* L.)

♂. C. 1819 herb. nostr.

verticillatum Spr. (*Mollugo* L.

Moll. dichotoma Schrauk).

♂. D. 1837.

Phaseolus (Bönor).

bicolor. ♂. D. ex hort. Crac. 1837.

lilacinus ♂. D. ex hort. Crac. 1837.

multiflorus Spr. (*vulgaris* β. *coccineus* L.) ♂. D.

nanus L. ♂. D. varietates plures coluntur in h.

vulgaris L. ♂. D. varietat. seminib. varie pictis adsunt.

Philadelphus

(Schersmin).

coronarius L. β. D.

grandiflorus W. β. D.

Phillyrea

angustifolia L. β. T. 1 uln. alt.

latifolia L. β. T. etiam s. Dio

2 hiemes perduravit. 3½ uln. alt.

Philodendron

crassifolium. 2. C. sub h. nom. ex h. Edinb. 1837.

Phleum (Kampe).

arenarium L. (*Phalaris* L.) ♂.

D. ex h. Crac. 1837.

asperum Vill. (*Phalaris* Retz.)

♂. D. 1815, et ex h. Hafn. 1836.

Böhmeri Wibel. 2. D. 1826, et ex h. Hafn. 1837.

Michellii All. 2. D. 1812. 1826.

pratense L. 2. D. spont.

Phlomis

armeniaca W. β. F. ex h. Petropol. 1836.

fruticosa L. β. F. 1810, et ex h. Paris. 1837.

laciniosa L. 2. F. ex h. Par. 1837,

pungens W. 2. D.

tuberosa L. 2. D. 2½ uln. alt.

Phlox

acuminata Pursh. (decussata)

2. D. tripedal. flor. violaceo-rub. medio Aug. explic.

divaricata L. 2. D. bipedal. flor. violaceo-caerul. Jun.

paniculata L. 2. D. 4-pedalis, flor. pallide rubr. Aug.

β. fl alb. intr. 1835. cur.

Cel. Engeström.

pyramidalis Sm. Spr. (*lutifolia* Mx.) 2. D. 1835.

svaveolens Ait. 2. D. sesquipedalis, flor. alb., intr. 1833.

Phoenix (Palmträd).

dactylifera L. β. C.

farinifera Roxb. (*pusilla* Lour.)

β. C. intr. 1834.

humilis Cav. v. *Chamaerops*.

Phormium

aloides L. v. *Lachenalia tricolor*.

tenax Forst. 2. T. fol. 1--2

uln. long.; flores non protulit in horto.

Phyllanthus

calycinus Labill. β. C. ex h.

Kew. 1837. 4g.

cantonensis Horn. ♂. C. 1812. in h. cult.

juglandifolius W. & C.
speciosus Jacq. v. *Xylophylla*
Arbuscula.

Phyllis

Nobla L. & T.

Phyllobium

chinense Fisch. T.

Physalis

Alkekengi L. & D. 1808 herb. nostr.

angulata L. ☉. C. ex h. Par. 1837.

curassavica L. & T.

parviflora R. Br. ☉. T. ex h. Hafn. 1837.

peruviana L. Horn. (*pubescens* Spr., *edulis* Sims.) ☉. & T.

Rothiana H. Hafn. communic. 1837.

Phyteuma*

canescens Kit. & D. 2 uln. alt.

Scheuchzeri All. & F.

virgatum W. (*Campanula virgat.* Labill.) ♂. F. ex h. Par. 1837.

Phytolacca

decandra L. & F. olim (1810) in hort. cult.

icosandra L. & C.

octandra L. & T. 1810.

Picris

altissima Delil. (*Crepis lappacea* W.) ☉. D. ex h. Paris. 1837.

globifera H. Par. ☉. D. 1815, et ex h. Paris. 1837.

hieracioides L. (*davurica* Fisch.) ♂. & D. 1826.

Sprengeriana Chaix. Spr. ex h. Paris. 1837.

Pimelea*)

decussata R. Br. & T. ex h. Kew. 1837. *J. Ag.*

glauca R. Br. & T.

hypericina; & T. sub h. nom. ex h. Kew. 1837.

*) *nimum consonans* cum *vetter. Coleopterorum genere Pimelia.*

Pimpinella

Anisum L. (*Sison* Spr. Svetic. Anis) ☉. D.

peregrina L. (*Tragium* Spr.) & F. 1835.

rotundifolia M. B. (*Sison* Spr.) & D. ex h. Hafn. 1837.

Saxifraga L. & D. spont.

Pinus (Tall. Gran).

Abies L. & D. in *Paradislyc-* kan 29 uln. alt., trunci diamet. 14-pollicar.

alba Ait. & D. 33 uln. alt., trunci diamet. mox supra terram 1 uln.

Balsamea L. & D. 6 uln., in nov. Arboret.

canadensis L. & D. 1½ uln., in nov. Arboret.

Cedrus L. & F. 1 uln. alt.; frustra sub Dio in hort. plantata.

Cembra L. & D. 3 uln. alt.

Larix L. (*Lärkträd*) & D. in *Paradislyck.* 26 uln. alt., trunc. diam. 16-poll.

nigra Ait. & D. 20 uln. alt.

Picea L. & D. 16 uln. alt., trunci diamet. 1 ped. supra terram 6-pollic.

Pinca L. & F. 2 uln. alt., in h. etiam sub Dio pluries, sed sine successu plantata.

Strobus L. & D. 2 uln. alt.

sylvestris L. & D. in *Paradislyc-* kan; 27 uln. alt., trunc. diamet. 15-pollic.

Piper (Peppar).

blandum Jacq. & C.

inæqualifolium Vahl. (*Peperomia* R. et P. *Pip. aromati-* cum W.) & C.

magnoliæfolium Jacq. & C.

pellucidum L. ☉. C. 1819.

verticillatum L. & C.

Piqueria

trinervia Cav. & T. 1814 in hort. cult. herb. nostr.

Pisum (Ärter).

arvense L. ☉. D.

- Ochrus* L. v. *Lathyrus*.
sativum L. ☉. D. Plures variet. in hort. adsunt.
- Pittosporum**
Tobira Ait. (*Evonymus* Thunb.)
 ♀. T. sesquipedal.
undulatum Andr. ♀. T. 2 uln.
 alt. florib. alb. fragrantib.
 Mart. April. explicat.
- Plantago**
alpina L. ♀. D. 1822-1826.
asiatica L. ☉. ♀. D. ex hort.
 Paris. 1837.
Lagopus L. ☉. D. ex h. Par.
 1837.
lanceolata L. ♀. D. spont.
major L. (Groblad) ♀. D. sp.
maxima Jacqv. ♀. D. 1826.
 1833.
media L. ♀. D. spont.
microcephala Poir.; ex hort.
 Hamb. 1824 et Parisiens.
 1837.
Psyllium L. ☉. D.
saxatilis M. B. ♀. D. 1824, et
 ex h. Paris. 1837.
stricta Schousb. ☉. D. 1812.
 herb. nostr.
- Platanus**
acerifolia W. ♀. D.
cuneata W. ♀. D. 2½ uln. alt.
 intr. 1833.
occidentalis L. ♀. D. 20 uln. alt.
orientalis L. ♀. D. 21 uln. alt.
- Plectranthus**
fruticosus Herit. ♀. T. 1810.
rugosus H Hamb. ♀. ♀. T.
 intr. 1834.
parviflorus W. ♀. T. 1810.
- Plumbago**
micrantha Ledeb. ☉. ex hort.
 Hafn. 1836.
zeylanica L. ♀. C. flor. alb.
- Poa** (Gröe).
alpina L. ♀. D.
annua L. ☉. D. spont.
compressa L. ♀. D. 1826.
Eragrostis L. ☉. D. 1812. 1815.
laxa Hänk. ♀. D.
 species.
- Podaliria**
australis W. v. *Baptisia*.
tinctoria W. (*Sophora* L. *Baptisia* R. Br.) ♀. F.
- Podolepis**
gracilis Hook. ☉. T. ex h.
 Bonn. 1836.
- Podophyllum**
peltatum L. ♀. D. 8 poll. alt.,
 flor. alb., Jun.
- Poinsettia**
pulcherrima H. Hamb. ♀. C.
 ex h. Edinburg. 1837.
- Polemonium**
cæruleum L. ♀. D. flor. Aug.
 β. fl. alb.
dissectum Reichb. ♀. F. ex h.
 Gryph. 1837 a Ccl. *Horn-*
schuch nob. commun.; ses-
 quipedal.
gracile W. ♀. D. ex h. Lon-
 din. 1834 et Gr. 1837.
lacteam Lehm. ♀. F. ex h. Gr.
 1836.
mexicanum Cerv. ☉. ♂. ♀. T.
 ex h. Paris. 1837.
reptans L. ♀. D. flor. cærul.
 intr. 1824.
- Pollinia**
Gryllus Spr. (*Andropogon* L.)
 ♀. T. ex h. Hafn. 1837.
- Polyanthes** (Tuberoze).
tuberosa L. ♀. C. flor. alb.,
 svaveolent.
- Polycarpon**
tetraphyllum L. ☉. D. 1811.
 herb. nostr.
- Polygala**
- Polygonum**
acidulum W. ♀. D. 14--16
 pollic. alt.; intr. 1830.
aviculare L. (Trampgräs) ☉.
 D. spont.
Bellardi Atl. ☉. D. 1808-1815.
Bistorta L. (Ormrot) ♀. D.;
 in Paradislyckan jam neg-

lectum, sed tamen bene et copiose viget in umbrosis; 1-1½ uln. alt.

Convolvulus L. (Binda) ☉. D. spont.

divaricatum L. 2. D. 1 uln. alt.

elegans Tenor. ☉. D. 12-20 poll. alt. fl. alb. ante explicationem rubr.

emarginatum Roth. ☉. D. 1822.

Fagopyrum L. (Bohveče) ☉. D.

frutescens L. v. *Tragopyrum lanceol.*

Hydropiper L. ☉. D. spont.

orientale L. ☉. D. 1819. 1834.

Laxmanni Lepech. 2. D. 16-18 poll. alt., fl. Jun. uti plurima.

pensylvanicum L. ☉. D. ex h. Paris. 1837.

Persicaria L. ☉. ♂. D. spont.

salignum W. 2. D. 18-20 poll. alt. intr. 1830.

tataricum L. ☉. D.

tinctorium Lour. Ait. ☉. ex h. Bonn. 1836.

undulatum Murr. 2. D. 1826.

Polypodium

aureum L. 2. C. bipedale; intr. 1834 ex h. Hamb.

canariense L. v. *Davallia*.

cristatum L. v. *Aspidium*.

vulgare L. (Stensöta) 2. D.

Polypogon

monspeliensis Desf. (*Agrostis panicea* W.) ☉. D.

Pomaderris

quercifolia H. Hamb. 2. T.

Populus (Poppel. Asp).

alba L. (*nivea* W. Retz.) 2. D. 20 uln. alt.

angulata Ait. 2. D. 8 uln. alt.

balsamifera L. 2. D.

dilatata Ait. (*pyramidata* Münch. Pyramid-Poppel; Italicus Poppel) 2. D. in Paradislyckan; 37 uln. alt.

Lindleyana 2. D. sub h. nom. commun.; in nov. arb. 6 uln. alt.

monilifera Ait. 2. D. in Paradislyck. etc.; 34 uln. alt.

nigra L. 2. D. mas. in Paradislyckan 35 uln. alt.

pendula (*tremula* var. *pendula* H. Hamb.) 2. D. in nov. arbor. intr. 1833; hodie 5 uln. alt.; quotannis floret.

tremula L. 2. D.

Portulaca (Portlak).

oleracea L. ☉. D.

Potentilla

alba L. 2. D. fere pedalis; fl. alb., mens. Maj. explic.

agrimonioides M. B. 2. D. ex h. Paris. 1837.

Anserina L. 2. D. spont.

argentea L. (*neglecta* Baumg.) 2. D.

arguta Pursh. 2. D. ex h. Bonn. 1834 et Gryph. 1837.

atrosanguinea Don. Wilbr. 2. F. 1834, et deinde ex h.

Gryph. 1837. a cel. *Hornschuch* miss.

aurea L. 2. D. 3-4 poll. alt.; flor. Maj.

canescens Bess. (*adscendens* Kit. W.) 2. D. 1826, et ex

hh. Gryph. et Paris. 1837; 3-pedal.

caulescens L. 2. D. 1822 herb. nostr.

collina Wib. (*P. Thomasii* Tenor.) 2. D. ex h. Gryph. 1837.

confertiflora Torr. 2. F. D. 1834. 1 uln. alt.; fl. Aug.

crocea Lehm. (*aurea* Sm.) 2. D. ex h. Crac. 1837.

divaricata Cand. 2. D. ex h. Paris 1837.

formosa Don. Spr. (*nepalensis* Hook. *colorata* Hortul.) 2. D.

F. D. 1834, et ex hh. Crac.
et Paris. 1837. *spithamea*.
Fragaria Poir. (*Comarum fragarioides* W. *Fragaria sterilis* L.) 2. D.
fruticosa L. (*floribunda* Pursh. Svet. Tok) $\frac{1}{2}$. D. $1\frac{1}{2}$ uln. alta.
geoides M. B. 2. D. ex hort. Hamb. 1824 et Par. 1837.
grandiflora L. 2. F. D. ex h. Crac. 1837.
Güntheri Pohl. Spr. (*sordida* Fries. *canescens* Hartm.) 2. D. ex h. Paris. 1837.
heptaphylla Mill. 2. F. ex h. Gryph. 1837.
hungarica H. Crac. 2. D. ex h. Crac. 1837.
laciniosa Kit. 2. D. intr. 1834, et ex h. Gryph. 1836 quoque miss.
mollissima Lehm. 2. F. ex h. Gryph. 1837.
multifida L. 2. D. caul. ped. et ultra; flor. flav. mens. Jun. explic.
nitida L. 2.; ex h. Crac. 1836.
obscura W. 2. D. 1826, et ex h. Gryph. 1837.
opaca L. Fr. 2. D.
pedata W. 2. D. ex h. Paris. 1837.
pensylvanica L. 2. D. ex h. Hafn. 1827 et Paris. 1837.
pimpinelloides L. 2.; ex hort. Crac. 1837.
recta L. 2. D. 1826, et ex h. Gryph. 1836. flor. pallide flav.
reptans L. 2. D. spont.
rupestris L. 2. D. sesquipedal.; flor. alb. mens. Jun. explic.
ruthenica W. 2. D. 9-10 poll. alt.; flor. flav. mens. Maj. explicat.
Salesovii W. 2. F. ex h. Gr. 1837.
taurica W. 2. F. ex h. Paris. 1837.
verna L. Fr. 2. D.
viscosa Donn. 2. D. ex hort. Gryph. 1837.
species.

Poterium

garganicum; sub hoc nom. ex h. Paris. 1837.
hybridum L. 2. D. 1808.
polygamum Kit. 2. D. 3-pedale; flor. Jun.; intr. 1810.
Sangvisorba L. 2. D. 3-ped.; flor. Jun.
Westphalicum; 2. D. 1835. 3-pedal.; flor. Aug.
species 2. D.

Pothos

Harrisii Graham. 2. C. ex h. Edinburg. 1837 cur. Cl. J. Ag.

Prasium (Niccoline).

majus L. $\frac{1}{2}$. T. flor. alb.

Prenanthes

hieracifolia W. v. *Crepis pulchra*.
purpurea L. 2. D. $1\frac{3}{4}$ uln. alt.; flor. violac. mens. Aug. explicantur.
viminea L. ♂. F. ex h. Paris. 1837.

Primula

acaulis Lehm. (*P. veris* var. Fr.) 2. D. 1835; etiam ex h. Hamb. 1836.
Auricula L. 2. D. flor. colore mire ludunt, etiam. plen.
cortusoides L. 2. F. D. flor. rubr.
elatio Jacqv. 2. D.
farinosa L. (Majblomster) 2. D. in horto ægre colitur, vix ultra 2 annos vigens.
inflata Lehm. 2. F. ex hort. Hafn. 1827. 1836.
nitens; 2. D. 3 poll. alt. flor. lilacin. versus finem Apr. explic.; intr. 1834, per 3 hiemes sub Dio duravit.
prænitens Ker. (*chinensis* Lindl. Hortul.) 2. T. flor. rosei, speciosi, abundantes; intr. 1836 ex h. Hafn.
 β . fl. alb.
veris L. (Oxlägg) 2. D.

Prinos

glaber L. $\frac{1}{2}$ F. D. 1 uln. alt.,
intr. 1835.

verticillatus L. (*padifolius* W.?)
 $\frac{1}{2}$ D. 1 $\frac{1}{2}$ uln. alt.; intr.
1834 ex h. Hamb.

Prismatocarpus

falcatus Tenor. v. *Campanula*

Prostanthera

lasiantha Labill. $\frac{1}{2}$ C. T. 20
poll. alt. fl. pallide rubr.
d. 1 Aug. 1836. explicat.
intr. 1835 ex h. Hamb.

Protea

anemonifolia Salish. v. *Isopogon*.
hirsuta W. v. *Leucadendron*.
Levisanus Berg. v. *Leucadendron*.

Prunella

grandiflora L. $\frac{1}{2}$ D. intr. 1837
ex h. Hafn.

hyssopifolia L. $\frac{1}{2}$ D. 1808.

pennsylvanica W. $\frac{1}{2}$ D. ante plu-
res annos in horto cult.
sec. herbar.

vulgaris L. $\frac{1}{2}$ D. spont.

Prunus

Armeniaca L. $\frac{1}{2}$ D.

Avium L. (Fogelbärsträd) $\frac{1}{2}$ D.
 β . asplenifolia H. Hamb.
2 uln. alt.

Cerasus L. (Körsbärsträd) $\frac{1}{2}$ D.
 β . fl. plen.

domestica L. (Plomonträd) $\frac{1}{2}$ D.
 β . fructu apice acuto; in
horto *Aprikos-Plomon* dicta.

grandifolia; $\frac{1}{2}$ D. 3 $\frac{1}{2}$ uln. alt.;
trunci diamet. 2 poll.; in
nov. arbor. quotannis fru-
ctus profert; intr. 1833.

insititia L. (Krikonträd) $\frac{1}{2}$ D.
12 uln. alt.

Laurocerasus L. (Lagerkers)
 $\frac{1}{2}$ F. etiam sub Dio plan-
tata, a ventisdefensa, coeli
nostri temperiem bene to-
lerat.

Mahaleb L. $\frac{1}{2}$ D. in Paradis-
lyckan. 8-18 uln. alt.
 β . fruct. luteo.

nigra Ait. $\frac{1}{2}$ D.

Padus L. (Hägg) $\frac{1}{2}$ D.

serotina Ehrh. $\frac{1}{2}$ D. 4-10 uln.
alt.; in hort. quotannis
fruct. dedit.

sibirica L. $\frac{1}{2}$ D. intr. 1834 e
seminibus enat.

virginiana L. (*rubra* W. Retz.)
 $\frac{1}{2}$ D. Cfr. Retz. Act. Holm.
1798. p. 68.

Psoralea

americana W. $\frac{1}{2}$ T. 1810.

acaulis $\frac{1}{2}$ T., sub hoc nom.
ex h. Petrop. 1836.

bracteata L. (*Ononis trifoliata*
L.) $\frac{1}{2}$ T. 1810, et ex h.
Paris. 1837.

glandulosa L. $\frac{1}{2}$ T.

Lagopus Cav. v. *Dalea*.

palæstina L. $\frac{1}{2}$ T. bipedalis et
ultra.

pinuata L. $\frac{1}{2}$ T. ex h. Kil.
1836.

Ptelea

trifoliata L. $\frac{1}{2}$ D. 5 uln. alt.,
fructum in hort. raro ma-
turat.

Pterocephalus

plumosus Spr. v. *Knutia*.

Pulmonaria (Lungört).

angustifolia L. $\frac{1}{2}$ D.

officinalis L. $\frac{1}{2}$ D.

Pulsatilla

patens Mill. W. (*Anemone* L.)
 $\frac{1}{2}$ D.

pratensis W. (*Anemone* L.) $\frac{1}{2}$ D.

vulgaris W. (*Anemone Pulsatilla*
L.) $\frac{1}{2}$ D.

Pultenæa

daphnoides Sm. $\frac{1}{2}$ T. intr. 1834
de h. Hamb. emta; deinde
ex h. Chiswich Angl. 1837
cur. J. Ag.

linophylla Sm. $\frac{1}{2}$ T. ex hort.
Chiswich 1837.

Punica

Granatum L. (*Guznat-äppel-träd*) $\frac{1}{2}$ T. F.

Pyrethrum

Balsamita W. (*Chrysanthem.* L.)

$\frac{1}{2}$ T. ex h. Paris. 1837.

corymbosum W. (*Chrysanthem.* L.) $\frac{1}{2}$ D. 1808-1826.

frutescens W. (*Chrysanth.* L.) $\frac{1}{2}$ T. 1808.

inodorum Sm. (*Chrysanth.* L.)

⊙ D.

latifolium W. $\frac{1}{2}$ D. 1826.

maritimum Sm. (*Matricaria* L.)

$\frac{1}{2}$ D. 1808.

partheniifolium W. $\frac{1}{2}$ D.

Parthenium Sm. (*Matricaria* L.)

⊙ ♂ D.

β fl. plen.

parviflorum W. (*præcox* Spr.)

⊙ D. 1821.

pulverulentum Horn. $\frac{1}{2}$ D. 2 uln. alt.

roseum M. B. $\frac{1}{2}$ D. 1822-1826.

serotinum W. (*Chrysanthem.* L.)

$\frac{1}{2}$ D.

species.

Pyrus

Amelanchier du Roi (*Mespilus*

L.) $\frac{1}{2}$ D. olim in horto

culta, hodie evanuit. Cfr.

etiam Retz. in Act. Holm.

1798 p. 61.

arbutifolia Ehrh. (*Mespilus* L.)

$\frac{1}{2}$ D. intr. 1810.

Aria Ehrh. v. *Sorbus*.

Aucuparia Sm. v. *Sorbus*.

baccata L. $\frac{1}{2}$ D. 13 uln. alt.

fract. var.

Botryapium L. $\frac{1}{2}$ D. 12 uln. alt.

communis L. (*Päronträd*) $\frac{1}{2}$ D.

coronaria L. $\frac{1}{2}$ D. in arboret.

1 uln. alt.

Cydonia L. v. *Cydonia vulgaris*.

edulis W. $\frac{1}{2}$ D. in nov. arbor.

2 $\frac{1}{2}$ uln. alt.

hybrida Mönch. v. *Sorbus*.

japonica Thb. $\frac{1}{2}$ T. 1 $\frac{1}{2}$ uln.

alt.; quotannis floret.

Malus L. (*Äppel*) $\frac{1}{2}$ D. plu-

res varietat. in horto col-

luntur.

melanocarpa W. $\frac{1}{2}$ D. 1 $\frac{1}{2}$ uln. alt., floret Jun.

nivalis Jacq. $\frac{1}{2}$ D. 6 uln. alt.

prunifolia W. $\frac{1}{2}$ D. 8 uln. alt.;

variat. fructu nigro, stri-

ato etc.

salicifolia L. $\frac{1}{2}$ D. 9 uln. alt.

spectabilis Ait. $\frac{1}{2}$ D. in Ar-

boret.

Quercus (Ek).

australis Link. $\frac{1}{2}$ T. F. ex h. Chiswich 1837.

bicolor W. $\frac{1}{2}$ D. $\frac{1}{2}$ uln. alt.,

intr. 1836, seminib. educat.

Cerris L. $\frac{1}{2}$ D. in arboret. 2 $\frac{1}{2}$

uln. alt.; intr. 1827 ex h.

Hafn.

Ilex L. $\frac{1}{2}$ F.; sub Dio frustra

plantata, in Frigidar. bene

viget.

macrocarpa Mx. $\frac{1}{2}$ D. intr.

1834, e seminib. enatis.

pendunculata W. $\frac{1}{2}$ D. 6 uln.

alta.

Prinos L. $\frac{1}{2}$ D. olim in horto sub

Dio, sed sine successu plan-

tata; Retz. in Act. Holm.

1798. p. 70.

Robur L. $\frac{1}{2}$ D.

trinctoria W. $\frac{1}{2}$ D. in Para-

dislycan.

Suber L. (*Korkträd*) $\frac{1}{2}$ F. 1

uln. alt.

Queria

canadensis L. v. *Amychia dicto-*

loma.

hispanica L. ⊙ D. ex hort.

Hafn. 1836.

Radiola

Millegrana Sm. (*Linum Radiola*

L.) ⊙ D. spont.

Ranunculus (Solbg.

Smörblomster).

- abortivus* L. 2. D. olim in h. cult. flor. flav.
aconitifolius L. 2. D. fl. alb.
acris L. 2. D. spont.
 β. fl. plen.
auricomus L. 2. D. spont.
bracteatus L. H. Crac. 2.; ex h. Gryph. 1837.
bruttius Tenor. 2. D. ex h. Gryph. 1837.
bulbosus L. 2. D.
caucasicus M. B. 2. D.
chærophyllus L. 2. D. ex h. Gryph. 1836.
chius Cand. 2.; ex h. Gryph. 1836.
Ficaria L. 2. D. spont.
fumarifolius Desf. 2.; ex h. Gryph. 1836.
gramineus L. 2. D. 14 pollic. alt. flor. flav. mens. Maj. explic.
illyricus L. 2. D.
muricatus L. 2. D. ex hort. Paris. 1837. flor. flav.
nemorosus Cand. 2. D. ex h. Gryph. 1836.
parviflorus Gouan. 2. D. 1815 et ex h. Paris. 1837.
pedatus Kit. 2. D. ex h. Gr. 1836.
repens L. 2. D. spont.
 β. fl. plen.
trikobus Desf. 2. F. ex h. Crac. 1836.

Raphanus (Rättika).

Landra Morett. 2. ex h. Kil. 1836.

Raphanistrum L. (*Raphanistrum Lampsana* Horn. Sv. Åker-rättika) 2. D.

sativus L. 2. ♂. D.

Rapistrum

ragosum All. v. *Cakile*.

Relbania

pungens Herit. 2. C. ex hort. Paris. 1837.

Renealmia

nutans Andr. v. *Alpinia*.

Reseda (Wau).

alba L. 2. D.

dipetala Ait.; ex h. Kil. 1835.

fruticulosa L. 2. F. 1823 herb. nostr.

Luteola L. 2. ♂. D. sæpe 3-pedalis, flor. viridi. flav.

mediterranea L. 2. D. ex h. Hafn. 1827 et Par. 1837.

odorata L. 2. ♂. D.

subulata (an *R. linifolia* Horn.); sub hoc nom. ex h. Par. 1837.

undata L. 2. D. 14 poll. alt. fl. pallide flav. mens. Jun. explic.

virescens Hornem. 2. D. ex h. Paris. 1837.

Rhagadiolus

edulis Gärtn. (*Lapsan. Rhagadiolus* L.) 2. D. 1835.

Kölpinia W. v. *Kölpinia linearis*. *stellatus* W. (*Lapsana* L.) 2. D.

Rhamnus (Toste).

Alaternus L. 2. T. F. 3 uln. et ultra alt.; etiam sub Dio 3 hiemes in hort. per duravit, sempervirens.

alnifolia Herit. 2. D. 1½ uln. alt.; in nov. arboret. introd. 1834 ex h. Hamb.

Clusii W. 2. F.

Erythroxylon Pall. 2. F. D. nuper horti nostri incola, seminibus educata, per 3 hiemes sub Dio duravit.

Frangula L. (Brackved, Tröske) 2. D. 8 uln. alt.

Paliurus L. v. *Paliurus australis*.

Rhaponticum

species, ex h. Petropol.

Rheum (Rabarber).

australe Don. 2.; ex h. Hafn. 1835.

caspicum W. 2.; ex h. Hafn. 1837.

compactum L. (*ataricum* L. suppl.?) 2. D. 1826, et ex h. Hafn. 1837.

crassicaule; sub hoc nom. ex h. Hafn. 1837.

hybridum Ait. 2. D. 2½ uln. alt. floret mens. Jun. uti sequentia.

palmatum L. 2. D. 3 uln. alt.

Rhaponticum L. 2. D. 1¼ uln. alt., in Paradistücken.

undulatum L. 2. D. 2½ uln. alt.

Rhodiola (Rosenrot).

rosea L. v. *Sedum Rhodiola*.

Rhodochiton

volubile, 2. C. T.; 1837. de h. Hamb. emt., etiam ex hort. Edinb. eodem anno communic.; trienne.

Rhododendron

ponticum L. 2. F. 34 pollic. alt. fl. specios., violaceo-rub. d. 9 Jun. explic.; ex h. Hamb. 1834.

Rhus

Cocodendron Ehrh. v. *Ailanthus glandulosa*.

copallina L. 2. D. intr. 1810.

Coriaria L. (Sumach) 2. D. 1806 horti incola, hodie exsul.

Cotinus L. 2. D.

radicans L. 2. D. 5-8 uln. long. 2. Toxicodendron L.

typhina (*Typhinum*) L. 2. D. 7 uln. alt.; in plantatione Paradistücken.

Ribes

alpinum L. (Måbär) 2. D. 3-4 uln. alt. in Paradistücken. 2. parviflorum.

aureum Pursh. (*jasminiflorum* Ag.) 2. D. 2-3 uln. alt. flor. aureis; variat fruct. magnitud., colore, sapore.

Diacantha L. 2. D.

divaricatum. 2. D. intr. 1834 ex h. Petropol. sub hoc nom.; seminib. educatum; primum floruit 1837.

floridum Herit. 2. D. fl. alb. glaciale Wall. 2. F. D. ex h. Chiswich 1837.

glutinosum 2. F. D. ex hort. Chiswich 1837.

Grossularia L. (Krusbärsbuske) 2. D.

longiflorum; 2. D. 2½ uln. alt. sub hoc nom. communic. vix nisi *R. aureum* est.

malvaceum. 2. F. D. ex h. Chiswich. 1837.

nigrum L. (Svart vinbärsbuske) 2. D.

2. fruct. virid.; in arbor. 1¼ uln. alt.

petraeum L. 2. D.

procumbens Pall.; 2. intr. 1835.

rubrum L. (Röd vinbärsbuske) 2. D.

sanguineum Pursh. 2. D. 2¼ uln. alt. florib. speciosis, purpur., uberrim. mens. Maj. explic., sed baccae haud maturat.

tenuifolium; 2. D. 1½ uln. alt.; intr. 1834 sub nom. *tenuiflorum* ex h. Petrop., e seminibus enatum.

Uva Crispa L. (Stickelbärsbuske) 2. D.

Richardia

scabra L. (2.) C. ex hh. Hafn et Paris. 1837. 7 poll. alt.

Ricinus (Underträd).

communis L. (2.) T. F. 1½ uln. alt. flor. Aug.

inermis Jacq. (*R. communis* var. Spr.) (2.) C. ex hort. Kil. 1836.

viridis W. (2.) C. ex h. Crac. 1836 (var. *R. communis*?)

Ricotia

Lunaria Cand. (*Cardamine* L. *Ricotia aegyptiaca* L.) (2.) F. T. ex h. Hafn. 1837.

Sorbit. in h. nostro Jul.
Aug.; 14-pollicaris.

Riödera

tetraspis Pall. (*Cynoglossum la-*
vigatum L. Spr.) 2. F. D.
ex h. Paris. 1837.

Rivina

humilis L. 2. C. T. ex h. Kil.
1836. flor. parv. alb.
purpurascens Schrad. 2. C. fl.
albis, demum rubris.
spec. 2. C.

Robinia

Caragana L. v. *Caragana arbo-*
rescens.

Chumlagu Herit. v. *Caragana*.

frutescens L. v. *Caragana*.

Halodendron Pall. (*Halimoden-*
dron argenteum Cand.) 2. D.

hispida L. 2. D. intr. 1833.
e seminib. enata; hodie 2
uln. alt.; florib. rubr. Jun.
explicat.

Pseud-acacia L. 2. D. in Pa-
radislyckan 30 uln. alt.;
fructum quotannis matura-
rat; flor. mens. Jun. ex-
plic. albis.

viscosa Vent. 2. D. 2 uln. alt.;
seminibus educata.

Rochea

fulcata Cand. v. *Crasanthus*.

Rödigia

laevigata Spr. (*Seriola alliata*
Biv.) 2. D.

Rosa (Rosor. Törne. Ny- ponbuske).

alba L. 2. D.

2. fl. plen.

Boursaulti H. Hamb. 2. F.

canina L. 2. D.

centifolia L. (Provins-Rosor
Pharmacop.) 2. D.; adsunt
in horto plures varietates,
inter quas eminet *Tuscany-*
black, 1835 introd. benev.
Cel. Prof. *Engeström*.

gallica L. (Ättike-Rosor) 2. D.
in Paradislyckan.

Lawrenceana Sweet. Spr. 2. T.
vix spitham.; sub nom. *R.*
Redoutea ex h. Hafn. 1833.

lutea Dalech. (*Eglataria* L.)

2. D.

2. bicolor Jacqv.

7. fl. plen.

microphylla Roxb. 2. F. intr.
1837, de h. Hamb. cuncta.

Muscosa Ait. 2. D.

odorata var. *Thea* H. Hamb.
2. T. intr. 1836, cur. Cl.

Stenkula.

pimpinellifolia L. 2. D. in
Paradislyckan.

provincialis Ait. 2. D. in Pa-
radislyckan.

repens Gmel. 2. D. in arboret.
et Paradislyckan.

semperflorens Cart. 2. F.

virginiana du Roi 2. F. 1835.

Plures aliae species in horto
quoque coluntur, inter
quas controversa *Königin*
von Dänemark-Rose a Cel.
Engeström nuper commu-
nicata.

Roscoeia

purpurea Sm. 2. C. intr. 1837.
ex h. Edinburg. cur. Cl.
J. Ag.

Rosmarinus

(Rosmarin).

officinalis L. 2. T.

Rothia

runcinata Roth. (*Andryala inte-*
grifolia L.) 2. T. 1807 in
hort. culta.

Rottböllia

biflora Pers. 2. D. 1812.

filiformis Roth. 2. F. ex h.
Hafn. 1837.

incurvata L. 2. D.

Rubia (Krapp).

tinctorum L. 2. D. catid. 3-pe-
dal., tenuis, decumbens,
flor. flavis.

Rubus (Hallon).

cassius L. 2. D.

candicans Weib. (*thyrsiflorus*
Fries Fl. Scan.) 2. D.

idoneus L. f. D.
odoratus L. f. D. 2½ uln. alt.;
 flor. purpur. mens. Jul. ex-
 plic. — Paradislyckan.
spectabilis Pursh. f. D. 1½ uln.
 alt.; flor. incarnat.; spe-
 cios., inodor. mens. Maj.
 explic.; intr. 1835. in ar-
 bore, 1837 in Paradis-
 lyckan.

Rudbeckia

alata Jacq. v. *Helenium 4-den-*
tatum.

amplexifolia Jacq. (*amplexi-*
caulis Vahl.) ♂. D.

columnaris Pursh. (*Obeliscaria*
 Cand. Hornsch.) 2. T. F.
 ex h. Gryph. 1837.

digitata Ait. 2. D. 3 uln. alt.;
 flor. flav. mens. Aug. ex-
 plic.

fulgida Ait. 2. F. 1813, petal.
 flav.

laciniata L. 2. D. 3 uln. alt.,
 flor. flav.

multifida; sub hoc nom. ex h.
 Paris. 1837.

pinnata Vent. Spr. (*Obeliscaria*
 Cass. Hornsch.) 2. D. 1826,
 et ex h. Gryph. 1837.

purpurea L. 2. D. 1½ uln. alt.;
 fl. rubr. mens. Aug. explic.
 1837. — intr. 1810.

triloba L. f. 2. D. 1807, et
 ex h. Paris. 1837. petal.
 flav.

Ruellia

antipoda L. v. *Bonnaya*.

formosa Andr. (*elegans* Poir.)
 2. C. flor. coccin.

strepens L. 2. F. 1806; etiam
 ex hort. Paris. 1837. flor.
 pallide rubr.

Rumex (Syr.)

Acetosa L. 2. D. spont.

β. *hispanica* Lfjehl.

Acetosella L. 2. D.

alpinus L. 2. D. ex h. Paris.
 1837.

cordifolius; sub hoc nom. ex

hh. Crac. 1835, et Paris.
 1837.

crispus L. 2. D. spont.

dentatus Murr. ♂. D. ex h.
 Paris. 1837.

Lunaria L. f. F. 1808. 1826.

maritima L. 2. ♂. D. ex h.
 Crac. 1836.

nepalensis Spr. 2. ex h. Paris.
 1837.

Patientia L. 2. D.

pulcher L. f. 2. D. ex h. Pa-
 ris. 1837.

scutatus L. 2. D. 1 uln. alt.

Rumia

seseloides (Cachrys M. B.) D. ex
 h. Petr. 1836.

Ruscus

aculeatus L. f. F. D. sesqui-
 pedal.; flor. explic. in fri-
 gidar. 8 April.

Hypoglossum L. 2. F. D. ses-
 quipedal.

racemosus L. f. F. D. intr.
 1822 ex h. Paris.; omnes
 hae species in horto sub
 Dio leniores hiemes per-
 duraverunt.

Ruta (Vincuta).

chalepensis L. Hornem. f. F.
 3-pedalis et ultra; flor.
 hepat. intr. 1810.

divaricata Tenor. f. F. D.

graveolens L. 2. D.

sylvestris; sub h. nom. ex h.
 Paris. 1837.

Sagina

procumbens L. ♂. D. spont.

Salicornia

fruticosa L. f. F.

herbacea L. ♂. D.

Salisburia

adanthrifolia Sm. (*Ginkgo bi-*
loba L.) f. F.

Salix (Pil).

- acutifolia* W. ♂. D. 7 uln. alt.
in Plantatione Paradisi.
alba L. ♂. D.
aurita L. ♂. D.
babylonica L. (Tärpil) ♂. D.
in Paradislyckan et nov.
arboresc.; femin. 5 uln. alt.
Caprea L. ♂. D. 8. uln. alt.
Croweana Sm. ♂. D. 1820.
fragilis L. ♂. D.
mollissima Ehrh. ♂. D. Para-
dislyckan.
nigricans L. ♂. D.
pentandra L. ♂. D.
purpurea L. ♂. D. in Paradis-
lyckan 6-8 uln. alt.
rosmarinifolia L. ♂. (femin.)
D. in novo-arboresceto 3 uln.
alt. flor. mens. Maj.
undulata Ehrh. ♂. D.
viminialis L. ♂. D.
viridis Fr. ♂. D.
species.

Salpiglossis

- Barclayana*; ♀. T. ex hh. Be-
rol. et Bonn. 1836. sesqui-
pedalis.
integrifolia Hook. (*Petunia mi-
rabilis*) ♂. ♀. T. 1835 1½
uln. alt.; flor. purpur.
straminea Hook. ♀. ♂. C.
T. ex h. Paris. 1836.
species.

Salsola

- Kali* L. ♂. D.
radiata Desf. (*Kochia dentata*
W.) ♂. ex h. Par. 1837.
Soda L. ♂. D. ex hort. Crac.
1836.
Tragus L. ♂. D. ex h. Paris.
1837.

Salvia

- Æthiopis* L. ♂. D. F. 1810
et ex h. Paris. 1837.
albida Horn. ♀. D. bipedalis.
applanata W. ♀. ex h. Gryph.
1836.
bicolor Lam. ♀. D. 1819, ex
h. Bonn. 1836.

- bullata* Schousb. ♀. F. ex h.
Paris. 1837, an eadem ac
S. lusitanica?
campestris M. B. ♀. F. D.
ceratophylla L. ♂. D. F. 1819
et ex h. Petr. 1836.
coccinea L. ♂. T.
diacermas L. ♀. F. 1810 et ex
h. Paris. 1837.
Forskålei L. ♀. D. F. sesqui-
pedalis.
glutinosa L. ♀. D. 1826. 1833.
Grahami Hort. ♂. T. ex hort.
Gryph. 1837.
grandiflora Etling. ♀. F. ex h.
Kil. 1836, fl. pallid. cærul.
Habliziana W. ♀. D. ex h.
Bonn. 1836.
hispanica L. ♂. D. 1810 et
ex h. Paris. 1837.
Horminum L. ♂. D. bracteis
rubr.; bipedalis.
β., bract. violaceis.
lanceolata Cav. ♂. D. 1810.
lusitanica Poir. ♀. D. 1826 et
ex h. Hafn. 1837.
lyrata L. ♀. D.
micrantha Vahl. ♀. T. ex h.
Hafn. 1837.
napifolia Jacq. ♀. D. (vide-
tur similis *S. verticillata*);
bipedalis.
nepetifolia; ♂. F. 1822 et ex
h. Paris. 1837.
nilotica Murr. ♀. ♂. F. flor.
cærulesc.
nutans Kit. ♀. D. ex h. Kil.
1836.
odorata W. ♂. F. ex h. Paris.
1837.
officinalis L. ♂. D.
patula Desf. ♂. F. D. ex h.
Paris. 1837.
pratensis L. ♀. D.
β. bicolor ex h. Gryph.
1837.
rubra Spr. ♂. D. ex h. Crac.
1836.
runcinata L. ♀. T.
scrotina L. ♂. ♂. F. ex hort.
Paris. 1837.
Sibthorpii Sm. ♂. ♀. D. ex h.
Gryph. 1837.

sylvestris L. 2. D. ex h. Paris. 1837.

Tenorii Spr. 2. T. ex h. Berol. 1824 et Gryph. 1836.
triloba L. (*S. Clusii* Jacqv.) 2. T.

Verbenaca L. 2. D. (*S. oblongata* Vahl. et *Spielmanni* W. huic synon. sunt secund. Sprengel).

verticillata L. 2. D.

viridis L. 2. D. ex h. Paris. 1837. fl. parv. roseis aut violaceis mens. Aug. expl.

viscosa Jacqv. 2. D. 1810 introd.

Sambucus (Fläder. Hyll).

canadensis L. 2. in horto sub Dio olim culta, hodie extincta. Cfr. Retz. Act. Holm. 1798. p. 74.

Ebulus L. 2. D. copiose, in plantatione Paradislyck.; 2-3 uln. alt. Augst. florens.

nigra L. 2. D. 4-8 uln. alt. fl. laciniata Retz. Liljebl. (Persilje-Fläder), in arboret. 2 uln. alt. γ. fructu viridi Retz. l. c.; hodie horti exsul.

racemosa L. 2. D. in arboret. 7 uln. alt., etiam in Paradislyckan.

Sanguisorba

canadensis L. 2. 1826.

officinalis L. 2. D.

β. *carnea* Fisch. ex hort. Gryph. 1836.

media L. 2. D.

tenuifolia Fisch. 2. F. D. intr. 1835.

β. *flor. albo* ex h. Gryph. 1836.

Santolina

Chamaecyparissus L. 2. F.

pinnata Donn. v. *Lastospermum pedunculata*.

viridis W. 2. T.

Sanvitalia

procumbens W. 2. D. 1811, herb. nostr.

Saponaria (Såpört),

cerastoides Fisch. 2. D. ex h. Paris. 1837.

glutinosa M. B.; ex h. Paris. 1837.

ocymoides L. 2. F. ex h. Paris. 1837.

officinalis L. 2. D.

β. *fl. plen.*

orientalis L. 2. D. 1810 et ex h. Paris. 1837.

porrigens L. 2. D. ex h. Paris. 1837.

Vaccaria L. v. *Gypsophila*.

Saracha

contorta R. et P. (*Atropa* Spr. Horn.) 2. C. ex h. Paris. 1837.

umbellata Cand. (*Atropa* Spr.) 2. C. ex h. Paris. 1837.

viscosa R. et P. 2. C. ex h. Paris. 1837.

Satureia (Kyndel).

hortensis L. 2. D.

montana L. 2. D. F. 1812.

Saussurea

species ex h. Petrop. 1835.

Saxifraga (Stenbräcka).

ajugifolia L.; ex hort. Paris. 1837.

caespitosa L. 2. F. 1808, herb. nostr.

cordifolia Haw. 2. D. ex h. Gryph. 1836.

Cotyledon L. 2. D. flor. alb.

crassifolia L. 2. D. 12--18 poll. alt.

cuneifolia L. 2. F. D. ex h. Paris. 1837.

dentata; ex h. Paris. 1837.

furcata; ex h. Paris. 1837.

Geum L. 2. D. ex h. Paris. 1837.

granulata L. 2. D.

hedypnois, ex h. Paris. 1837.

hirsuta L. 2. F. ex h. Paris. 1837.

longifolia Lapeyr. 2. D. ex h.

Gryph. 1836.

palmata Sm. 2. D. 1819. 1826.

punctata L. (*serrata* Ott.) 2.

D. 8-10-pollic. flor. tota
aestate.

sarmentosa L. 2. F. D. 8-10
poll. alt.

Schraderi Sternb. 2. D. ex h.
Paris. 1837.

umbrosa L. 2. D.

Scabiosa (Wädd).

africana L. v. (*Asterocephalus*).

alpina L. v. *Succisa*.

ambigua Tenor. v. *Asterocephalus*.

amoena Jacq. v. *Asterocephalus*.

argentea Murr. v. *Asterocephalus*.

arvensis L. 2. D.

atropurpurea L. v. *Asterocephalus*.

australis Wulf. 2. D. ex h.

Paris. 1837.

bannatica Kit. v. *Asterocephalus*.

carpathica Fisch. 2. D. flor.
mens. Jul.

caucasica M. B. v. *Asterocephalus*.

Columbaria L. v. *Asterocephalus*.

coriacea W. v. *Succisa laevigata*.

corniculata Kit. v. *Succisa* gen.
tauroides.

elata Horn. v. *Succisa tatarica*.

glauca v. *Asterocephalus*.

hybrida All. 2. D. etiam sub

nom. *Knautia hybrida* com-

muniat.; sesquipedal.; flor.

cub. mens. Aug. explic.

integrifolia L. 2. D.

lucida Horn. 2. F. D. 1810.
1819.

ochroleuca L. v. *Asterocephalus*.

palastina L. v. *Asterocephalus*.

prolifera L. v. *Asterocephalus*.

reversa Led. 2. D. 1836. 3-pe-
dalis et ultra.

sicula L. v. *Asterocephalus*.

stellata L. v. *Asterocephalus*.

Succisa L. v. *Succisa pratensis*.

syriaca L. v. *Succisa*.

tatarica L. v. *Succisa*.

transsylvanica L. v. *Succisa*.

species.

Scandix (Körsvel).

Anthriscus L. v. *Anthriscus pul-*
garis.

australis L. (*Wyllia* Hoffm.) 2.

D. ex h. Paris. 1837. in
horto floruit; 1-pedalis.

brachycarpa Guss. 2. F. ex h.
Paris. 1837.

Cerefolium L. v. *Cherophyllum*.

fumarioides Kit. v. *Anthriscus*.

odorata L. v. *Myrrhis*.

Pecten L. 2. D. 1-pedalis;
fruct. fere bipollic.

pinnatifida Vent. 2. D. 1824
ex h. Petr. 1836.

Schinus

mölle L. 2. T. 1836. intr. ex
h. Paris. 1822.

Schizanthus

humilis H. Gryph. 2. D. ex
h. Hamb. 1836.

pinnatus R. et P. 2. D. T. fl.
pulchr. variegat. mens. Jul.

explic.; intr. 1833.

perrigens Hook. 2. D. 1834
et ex h. Hafn. 1837.

retusus Hook. 2. T. ex hort.
Hamb. 1836.

Schkurgia

abrotanoides Roth. 2. T. D.
intr. 1841, sesquiped. flor.

flav.

Schoberia

corniculata Mey. 2. 1835. ho-
die horto deest.

Schoenus (Agh).

Schultzia

crinita Spr. (*Sison* Pers.) 2. D.
ex h. Petr. 1836.

Scilla

amoena L. 2. D. 8-10 pollic.
st. flor. azur. jam d. 10

Maj. in hort. explic.

bifolia L. 2. D. flor. caerulea.
mens. April. explic.

- italica* L. 2. D. 5-7 poll. alt. fl. pall. cærul. 10 Maj. expl.
- nutans* Sm. (*Hyacinthus non scriptus* L.) 2. D. scap. 6-8 poll. alt. in umbrosis pedalis.; flor. Jun.
- peruviana* L. 2. F. 1810, et ex h. Paris. 1837.
- Scirpus** (Säf).
- atrovirens* W. 2. D. ex hort. Hamb. 1824 et Par. 1837.
- Holoschoenus* L. 2. D. 1826.
- Schleranthus**
- annuus* L. ☉. D. spont.
- Scolopendrium**
- officinarium* Sm. (*Asplenium Scolopendrium* L. Fr.) 2. D. in hort. cult. pedale.
- Scolymus**
- grandiflorus* Desf. 2.; ex h. Paris. 1837.
- hispanicus* L. ♂. 2. D. ex h. Paris. 1837.
- maculatus* L. ☉. D. 1806, et ex h. Paris. 1837.
- Scoparia**
- dulcis* L. ☉. T. C. 10 pollic. alt. flor. parv., alb., d. 26 Sept. in Tep. explic. intr. 1834.
- Scorpiurus**
- muricatus* L. ☉. D. intr. 1837, flor. flav.
- sulcatus* L. ☉. D. 1812. 1822. flor. flav.
- vermiculatus* L. ☉. D. 1810. 1815 etc. flor. parv., flav.
- Scorzonera**
- calcitrapæfolia* Vahl. 2. D. ex h. Paris. 1837.
- eriosperma* M. B. ♂. 2. D. ex h. Paris. 1837.
- graminifolia* L. D. 1812.
- hispanica* L. ♂. 2. D.
- humilis* L. 2. D.
- taurica* M. B. 2. D. 1826.
- tomentosa* L. D. 1836,
- Scrophularia** (Flenört).
- appendiculata* Jacq. 2. D. F. 1824. 1835. 1½ uln. alt.; flor. Jun.
- aquatica* L. 2. D. tripedal. fl. Jul.
- auriculata* L. 2. D. 1822. 1826.
- betonicæfolia* L. (*hirsuta* Horn.) 2. D. 1826.
- lyrata* W. 2. F. 1810. 1833.
- nodosa* L. 2. D.
- orientalis* L. 2. F. 1810.
- peregrina* L. ☉. D. 1808. 1812. herb. nostr.
- vernalis* L. ♂. D. in h. neglecta in umbrosis plurib. locis luxuriat. 6-10 poll. alt. fl. flav. versus finem mens. April. explic. species.
- Scutellaria**
- alpina* L. 2. D. 8-10 poll. alt. flor. cæruleo-violac. labio alb. mens. Jun. explic.
- Columnæ* All. 2. D.
- integrifolia* L. 2. D. 1 uln. alt. fl. Jun.
- peregrina* L. 2. D. 1808. herb. nostr.
- Secale**
- Sedum**
- acre* L. (Fetknopp) 2. D. sp.
- Aizoon* L. 2. D. 12 pollic. alt. flore flav. Jul. explic.
- album* L. 2. D. 1826.
- altissimum* Poir. 2. T. ex h. Paris. 1837.
- Anacampseros* L. 2. D. 6-poll. alt. flore pallide subcærulescenti violaceo Aug. explic.
- cæruleum* Vahl. ☉. F. 3-4 poll. alt. flor. Aug.
- Ewersii* H. Hamb. 2. T.
- hybridum* L. 2. D. 6-8-poll. alt., flor. flav. Jun. expl. intr. 1810.
- oppositifolium* Sims, ex hort. Paris. 1837.
- populifolium* L. 2. D. 7 poll. alt., flor. alb. apice rubr. Aug. explic.

Rhodiola Cand. (*Rhodiola rosea* L.) 2. D.
sempervivoides Fisch. 2. T. ex h. Bonn. 1836.
sexangulare L. 2. D. fl. flav.
spurium M. B. 2. D.
Telephium L. 2. D. 1 uln. fl. alb. Aug. explic.
β. purpur. 1808. herbar. species.

Selinum

austriacum Jacqv. (*montanum* W.) 2. D. 1826, et ex h. Paris. 1837.
Carvifolia L. (*Angelica* Spr. *Coneoselinum* Reichb. Fr.) 2. D. bipedal.
decipiens Wendl. 2. T. F.

Sempervivum

arboreum L. 2. T.
aureum Chr. Sm. 2. T. intr. 1834 ex h. Bonn.
barbatum Chr. Sm. (*spathulat.* Horn.) 2. T.
cæspitosum Chr. Sm. (*barbat.* Horn.) 2. T. fere pedale, flor. flav. mens. Maj. expl.
tectorum L. (Taklök) 2. D.
tortuosum Ait. 2. T.
urbicum Horn. 2. T. F.
villosum Ait. (2. H. Gryph.) 2. 2. T.
 species.

Senebiera

pinnatifida Cand. v. *Coronopus didymus*.

Senecio

egyptius L. 2. D. intr. 1810, deinde etiam ex h. Paris. 1837. cur. Cl. J. Ag.
arabicus L. 2. D. ex h. Par. 1837.
chrysanthemifolius, 2. D. sub h. nom. ex h. Paris. 1837.
delphinifolius Vahl. 2. D. 1835.
desquamatus Horn. 2. D. 1812.

elegans L. 2. D.
β. fl. plen. et alb.
erucifolius L. 2. D.
hieracifolius L. 2. D. ex h. Paris. 1837.
Jacobæa L. 2. D.
lyratifolius Reichb. 2. D. ex h. Paris. 1837.
ovatus W. 2. D. 3½ uln. alt., fl. flav. mens. Sept.-expl.
saracenicus L. 2. D. 3 uln. alt., flor. flav. mens. Aug. explic.
tenuifolius Jacqv. 2. D. 3-pedal. flor. flav.
triflorus L. 2. D. ex h. Kil. 1836. pedalis, flor. flav.
venustus Ait. 2. C. ex hort. Paris. 1837.
vulgaris L. 2. D. spont.

Seriola

ætnensis L. 2. D. 1810. 1812, et ex h. Paris. 1837.
alliata Biv. v. *Rodigia lavigata*.

Serratula (Skära).

alata W. suppl. 2. D. 1826, et ex h. Paris. 1837.
arvensis L. v. *Carduus*.
Behen Cand. (*Centaurea* L.) ex h. Petr. 1836.
centauroides L. 2. D. intr. 1835, et ex h. Petr. 1836.
coronata L. 2. D. 1826, et ex h. Paris. 1837.
tinctoria L. (Ångskära) 2. D. species.

Seseli

corsicum Led. (*ammoides* Spr.) 2. D. 1836.
Hippomarathrum L. 2. D. ex h. Paris. 1837.
montanum L. 2. D. 3-pedale, flor. Aug.
varium Trev.; D. ex h. Paris. 1837.

Sesleria

echinata Host (*Echinaria capitata* Desf. H. Paris. *Cenchrus*

capitatus L.) ☉. D. ex h. Paris. 1837. flor. Aug.
elongata Host. 2. D. 12--20
 poll. alt.; flor. Aug.

Setaria P. B. v. *Panici* species.

Sherardia
arvensis L. ☉. D.

Sibthorpia
peregrina Spr. v. *Disandra prostrata*.

Sicyos
angulata L. ☉. D. 1810, et ex h. Paris. 1837.
 species.

Sida

Abutilon L. ☉. D. flor. flav. angustissima; ex h. Par. 1837.
carpinifolia L. fil. 2. C. ex h. Paris. 1837.

Dilleniana W. (*Anoda* Cav.) ☉. D. ex h. Paris. 1837.
dioica Cav. (*Napaea* L. *Sida scabra*) 2. D. 3-4 uln. alt. flor. parv., alb. mens. Aug. explic.

hastata W. (*S. cristata* β. L.) ☉. T. F. sesquipedal. fl. violaceo-rubr. l. lilacin. mens. Aug. explic.

incarnata Spr. (*Anoda* Kunth.) ☉. 2. T. ex hh. Hafniae 1836, et Paris. 1837.

indica L. ☉. C. 1835.
mollis Ort. 2. T. ex h. Paris. 1837.

Napaea W. (*Napaea laevis* L.) 2. D. 3; uln. alt., flor. parv. alb. mens. Sept. expl.

parviflora Cand. ☉. C. 1820.
tiliaefolia Fisch. ☉. D. 1822.
triangularis Humb. ☉. D. 1820.
triloba Cav. 2. T.
ulmifolia Cav. ☉. C. 1808.
 species.

Sideritis

brutia Tenor.; ex hort. Paris. 1837.

canariensis L. 2. T. 1810.

montana L. ☉. D. 1808.

perfoliata L. 2. 2. D. F. ex h. Paris. 1837.

romana L. ☉. D. 1810 et ex h. Paris. 1837.

rubro-punctata Retz. Pugill. observ. botan. 1810.

scordoides L. 2. D. flor. flav. lab. super. alb.

syriaca L. 2. 2. T. 1810, et ex h. Paris. 1837. an eadem ac *S. brutia*? florentem non vidi.

species 2. T.

Siegesbeckia

flosculosa Herit. ☉. D. T. 1810 et ex h. Paris. 1837.

orientalis L. (*triangularis* Cav.) ☉. D. T. 1808; — ex h. Paris. 1837. — *bipedalis*, flor. pauc. flav.

parviflora v. *Galinsogea*.

Silene

egyptiaca L. ex hort. Crac. 1836.

altaica Pers. (*jeniseensis* W.) ☉. D. 1815. 1835-1837. sesquipedal.

antirrhina L. ☉. D. 1810. 1815 et ex h. Crac. 1837.

apetala W. ☉. D. 1815 et ex h. Crac. 1836.

Armeria L. ☉. D. β. fl. alb.

Atocion Jacqv. (*orahidea* L.) ☉. D. 1810-1837.

Behen L. ☉. D. 1834. 1835.

bellidifolia Jacqv. ☉. D. 1834. 1835.

catholica Oth. (*Cucubalus* L.) 2. D.

cerastoides L. ☉. D. 1835 e seminib. in h. Lund. 1834 matur.

chloræfolia Sm. 2. 2. F. ex h. Petrop. 1836.

colorata Schousb. ☉. D. 10 poll. alt., fl. rubr. mens. Jul. explic.

compacta Horn. ♂. 2. D. 3-pedalis. flor. purp. anther. cærul. mens. Jun. explic. intr. 1835.
conica L. ♂. D. 1834. 1835.
davurica. ♂. D. ex h. Paris. 1837.
depressa M. B. 1822.
Dilleniana. ♂. D. ex h. Paris. 1837.
discolor Sm. in h. olim. cult.
flavescens Kit. ♂. 2. D. 1810 et ex h. Paris. 1837.
fruticosa L. ♂. T. F. 1835 1 uln. alt.; flor. pallid. rubr. mens. Jul. explic.; intr. 1810.
hispida Desf. ♂. D. ex hort. Paris. 1837.
inflata Sm. variet. angustifolia Tenor. 2. D.
juvenalis Del. ♂. ex h. Gryph. 1836.
latifolia Horn. 2. olim in h. cult.
laxiflora Brot. ♂. D. ex h. Gryph. 1836.
legionensis Lag. ♂. ex hort. Gryph. 1836.
livida W. 2. D. ex h. Paris. 1837.
longiflora Ehrh. 2. D. 1810. 1815 et ex h. Par. 1837.
lusitanica L. ♂. D. 1808. 1815. secund. herb. nostr.
mollissima Sm. Spr. (*Cucubalus* L. *Viscago* Horn.) 2. D. 1819-1835 bipedalis flor. alb. mens. Jun. explic.
Muscipula L. ♂. D. 1835 et ex h. Paris. 1837.
nocturna L. ♂. D. 1807-1815. 1834.
nutans L. 2. D. 8-10 pollic. alt., flor. alb.
ornata Ait. ♂. ♂. F. 1824 et ex h. Bonn. 1836.
patula Desf. 2. F. D. 1824. 1826.
pelidna Reichb. ♂. ex h. Gr. 1836.
pendula L. ♂. D. 1815. 1822. herb. nostr.

petraea Kit. 2. D. 1826 et ex h. Paris. 1837.
picta Pers (*reticulata* Desf.) ♂. D. 1827, et ex h. Paris. 1837.
Pourretii Ait. ♂. D. ex hort. Gryph. 1837.
præcox. ♂. D. ex hort. Paris. 1837.
quinquevulnæra L. ♂. D. 6-8 poll. alt., floret Jul., petal. albidis centro purpur.
rubella L. ♂. D. 1815 et ex h. Kil. 1836.
sardoa Moris. ♂. ex h. Gryph. 1837. forte eadem ac. S. *lusitanica*.
sericea All. ♂. D. 1815 et ex h. Paris. 1837.
stricta L. ♂. D. 14 poll. alt., flor. parv. ros. mens. Jul. explic.; intr. 1834.
stylosa. 2. D. intr. 1833 ex h. Bonn.; in nostr. floruit 1835; caul. 8 pollic. alt., flor. sord. alb.
supina M. B. 2. D. 1824.
svavis Schrad. ♂. ex h. Kil. 1836.
tridentata Desf. ♂. D. 1834. 1835.
species.

Siler

aquilegifolium Gärtn. v. *Laserpitium*.

Silphium

connatum L. 2. D. 3 uln. alt. flor., flav.
perfoliatum L. 2. D. circit. 3 uln. alt., flor. flav. Aug. explic.
trifoliatum L. 2. D. 1826.

Silybum (*Silybrum*).

marianum Gärtn. v. *Carduus*.

Sinapis (Senap),

- alba L. ☉. D.
 Allioni Jacqv. ☉. D. bipedal.
 flor. flav. mens. Aug. explicat. intr. 1810.
 arvensis L. ☉. D.
 brassicata L. ☉. D. 1808. secund. herbar. Fallén.
 nigra L. ☉. D. 1822. 1836. flor. flav.
 orientalis L. ☉. D. flor. flav. intr. 1810.
 β. grisea 1812.
 taurica Fisch. ☉. D. 1835.
 virgata Presl. (*laevigata* L.) ☉. D. 1835.

Sinningia

- villosa H. Hamb. 2. ♀. C.

Sison

- Ammi L. ☉. D. 1810; etiam ex h. Paris. 1837.
canadense L. v. *Myrrhis*.
Podagraria Spr. v. *Egopodium*.

Sisymbrium

- Alliaria* Scop. Fr. v. *Erysimum*.
 atrovirens Horn. ☉. T. bipedale, flor. flav. mens. Jun. explic.
austriacum Jacqv. Fr. (*Sisymb. Irio* Retz.) ☉. D. 1819. 1824. flor. flav.
 bursifolium L. ☉. D. ex h. Paris. 1837. flor. alb. — (*Hesperis dentata* L.).
 gallicum. W. ☉. D. ex hort. Crac. 1836; flor. alb.
laevigatum W. (*tenuifolium* L.?) 2. D. 1812.
 officinale Scop. (*Erysimum* L.) ☉. D. spont.
 persicum Spr. ☉. ♂. D. ex h. Kil. 1836. fl. pallide flavis.
 Sophia L. ☉. D. spont.
 strictissimum L. 2. D. 4-pedale et ultra, flor. flav. versus fin. Jun. explic.

Sisyrinchium

- album 2. T. ex h. Crac. 1836.
 Bermudiana L. 2. T.
 californicum Spr. (*Marica* Ker.) 2. T. scap. 7, folia 12 pollices alt., flor. flav.; intr. 1835.
 chinense; ex h. Crac. 1836.
 convolutum Nocc. 2. T. 1823, herb. nostr. flor. flav.
 grandiflorum Cav. 2. T. ex h. Chiswich. 1837.
 laxum Link. 2. T. ex h. Paris. 1837.
 micranthum Cav. 2. T. ex h. Crac. 1836.
 striatum Cav. (*Marica* Ker.) 2. T. F. 1822, et ex h. Paris. 1837, flor. pallide flavis brunneo-striat.
 tenuifolium Humb. 2. T. 1810. 1822, et ex h. Par. 1837.

Sium

- Bulbocastanum* Spr. v. *Bunium*.
 Falcaria L. 2. D. sesquipedal., flor. alb.
 lancifolium M. B. 2. D. 2 uln. alt., flor. alb. mens. Aug. explic.; intr. 1835.
 peucedanoides Spr. 2. ex h. Hafn. 1827 et 1837.
 siculum L. 2. ex h. Par. 1837.

Smilax

- rotundifolia L. ♀. T. intr. 1837.

Smyrniun

- Olusatrum L. ♂. 2. D. ex h. Paris. 1837.
 perfoliatum L. ♂. D. ex hort. Paris. 1837.

Solanum

- atro-sanguineum. ♀. T. ex h. Paris. 1837.
 cerasiforme. ☉. T. 1822 et ex h. Paris. 1837.
 decurrens Balb. ex h. Par. 1837.
 Dulcamara L. (*Qvesved. Hållbär*) ♀. D. spont.
 flavum Kit. ☉. D. ex h. Paris. 1837.
Humboldtii W. v. *Lycopersicon esculentum* β.
 lycioides L. ♀. C. ex h. Paris. 1837.

- lycopersicum* L. v. *Lycopersicon*
esculentum.
marginatum L. fil. p. T. ex h.
Crac. 1836.
Melongena L. (*esculentum* Du-
nal et Spr. Svet. Äggplan-
ta) ☉. C. 1½ uln. alt.
flor. lilacin.
β. fruct. violaceo.
nigrum L. ☉ D. spont.
β. humile Bernh. Fr.
oleraceum Rich. ☉. ex hort.
Paris. 1837.
pendulum R. et P. ex h. Crac.
1837.
persicum W. 4. p. D. sub
hoc nomin. semina missa
1835. ex h. Petropol.
Pseudocapsicum L. p. F. 1 uln.
alt. flor. alb. anther. flav.
pterocaulon Dun. ☉. D. ex
hort. Paris. 1837. an var.
S. nigri?
pyriforme Wilbr. v. *Lycopersi-*
con.
sodomeum L. p. T. ex h. Pa-
ria. 1837.
villosum Lam. Fr. Fl. Sc. ☉. D.
species.

Solidago (Gullris).

- altissima* L. 4. D. olim horti
incola, hodie exsul.
ambigua Ait. 4. D. in hort.
olim culta, sed ibi diu-
tius non exstat.
canadensis L. 4. D. 2½ uln.
alt. flor. aur. Aug. Sept.
flexicaulis Spr. (*latifolia* L.?)
4. D. fere 1½ uln. alt.;
flor. flav.
gigantea Ait. 4. D. 3 uln. alt.;
flor. flav.
hirta W. 4. D. ex h. Paris.
1837.
humilis Pursh. ex hort. Paris.
1837.
lanceolata L. 4. D. 1½ uln.
alt. flor. flav. Sept. Oct.
lateriflora L. 4. D. 1 uln. alt.;
fl. flav. inodor. Aug. Sept.

- mexicana* L. (*multiflora* Poir.)
4. D. 1 uln. alt. fl. flav.
minuta L. 4. D. 1½ uln. alt.
flor. flav. Sept.
nutans, ex h. Paris. 1837.
patula W. 4. D. 1½ uln. alt.
flor. flav. Sept.
procera Ait. 4. D. 1808. 1820;
3 uln. alt. flor. flav.
rigida L. 4. D. circiter 1½
uln. alt.

Sonchus

- arvensis* L. 4. D. 1½ uln. alt.
caucasicus Horn. 4. D. 1836.
floridanus L. ♂. D. ex hort.
Paris. 1837.
fruticosus L. (*sgarrosus* Cand.
Spr.) p. T. 2 uln. alt. fl.
flav.
hyoserifolius Horn. p. T. intr.
1834.
macrophyllus W. 4. D.
oleraceus L. (Mjölkstistel) ☉.
D. spont 1½ uln. alt.
pinnatus Ait. p. T.
tingitanus Lam. ☉. D. 18-20
poll. alt. flor. flav. basi
brunn. Aug. expl.

Sophora

- alopecuroides* L. 4. T. F. 1834
et ex h. Petr. 1836.
australis L. v. *Baptisia*.
flavescens Ait. T. ex h. Petr.
1836.
japonica L. p. D. 4 uln. alt.;
intr. 1835 in novum. ar-
bor., e semin. enat.
tetraptera Ait. et Forst. v.
Edwardsia.

Sorbus (Rönn).

- Aria* Pers. Fr. (*Crataegus* L.
Pyrus Spr. Svet. Oxel.) p. D.
Aucuparia L. (*Pyrus* Spr.) p. D.
β. fructu luteo.; in novo
arhoreto 4 uln. alta.
hybrida L. Fr. (*Pyrus* Spr.)
p. D.

Sorghum

bicolor W. (*Holcus* L.) ☉. D. 1810, et h. Hafn. 1836.

cernuum W. ☉. D. 1810 et ex h. Petr. 1836.

rubens Horn. ☉. D. 1834 et ex h. Hafn. 1836. & vulgaris variet. videtur.

saccharatum Pers. ☉. D. ex h. Hafn. 1836. sub nom.

Andropogon Sorghum Brot. ex h. Kilens. 1836, etiam

communicat. idem videtur.

vulgare Pers. (*Holcus Sorghum* L.) ☉. D. 2 uln. alt.

Sparaxis

tricolor Ker. (*Ixia* Curt.) 2. T. intr. 1834, etiam ex h. Paris. 1837.

Sparrmannia

africana Thunb. ½. T. 3 uln. et ultra alt. flores albos, pulchr., uberrim. quotannis profert.

Spartina

cynosuroides Roth. (*Dactylis* L. fil.) 2. D. ex h. Paris. 1837.

Spartium (Hareris).

ætnense Biv. (*Genista* Spr.) ½. T. ex h. Bonn. 1836.

juncum L. (*Genista* Scop. Spr.) ½. F. etiam sub Dio 2 annos in hort. jam duravit, a frigore hiemis non læsum.

multiflorum Ait. (*Genista alba* Spr.) ½. F. ex h. Hamb. 1837.

radiatum L. ½. F. ex h. Par. 1837.

scoparium L. (*Genista* Lam. Spr.) ½. D. 3 uln. alt. flor. flav. uberr. Jun. expl. in Paradislyckan.

spinosum L. (*Cytisus* Spr.) ½. T. F.

Spergula

arvensis L. ☉. D. spont.

nodosa L. ☉. D. spont.

Spermaceo

hirta L. ☉. C. ex h. Hafn. 1837.

longifolia Aubl. ☉. C. ex h. Hafn. 1837.

stricta L. v. *Bigeloviu*.

tenuior L. ☉. C. pedalis, flor. parv., alb.; ex h. Hafniæ 1837.

Spinacia (Spenat).

oleracea L. ☉. D.

Spiræa

Aruncus L. 2. D.

chamædryfolia L. ½. D.

crenata L. ½. D. 3 uln. alt., flor. alb. inodor. mens. Maj. explic.; in Paradislyckan.

Filipendula L. 2. D. flor. Jun. β. flor. plen. 1826.

hypericifolia L. ½. D. 3 uln. alt. flor. alb. mens. Jun. explicat.

lævigata L. ½. D. in nov. arboret. 1½ uln. alt. fl. parv., alb. mens. Maj. explic.

nutans. ½. T. ex h. Chiswich 1837.

opulifolia L. ½. D. 3-4 uln. alt. fl. alb., inodor. mens. Jun. explicat.

reflexa. ½. D. sub hoc nom. ex h. Hamb. 1834 miss. — hodie 3 uln. alt.

rosmarinifolia. ½. F. 1 uln. alt., flor. parv., alb. inodor. mens. Jun. explic.

rotundifolia. ½. F. 1836. -- sub Dio 1837.

salicifolia L. ½. D. 1-2 uln. alt., flor. pallide rubr. vix odor., Jun. — Sept. Paradislyckan etc.

sorbifolia L. ½. D. 3 uln. alt. in arbor. et Paradislyckan.

thalictroides L. ½. D. intr. 1834 ex h. Hamb.; hodie in nov. arbor. 1½ uln. alt., flor. parv., alb. mens. Jun. expl.

tomentosa L. ½. D. Retz. in Act. Holm. 1798. p. 75. hodie horti profuga.

Ulmaria L. 2. D. flor. Aug.
expl. grat. odor.
β. flor. plen. olim in hort.
cult.
species 2. D. in arboreto.

Stachys

alpina L. 2. D. 1826.
annua L. 2. D.
arabica Horn. 2. T. ex hort.
Paris. 1837.
arenaria Vahl. 2. F. ex hort.
Paris. 1837.
circinata Herit. 2. T. 1819.
(*Stach. velutina* W. 1837 ex
h. Paris. communic. forte
eadem).
corsica Pers. 2. F. ex h. Crac.
1836.
cretica L. 2. D. ex h. Gryph.
1836 (*Stach. dasyantha* Ra-
fin. secund. Spr. eadem).
dasyantha Rafin. 2. ex h. Gr.
1836.
excelsa Desf. 2. ex h. Gryph.
1837.
germanica L. 2. D. *bipedalis*
et ultra, flor. parv. rubi-
cund.
grandidentata. sub hoc nom.
ex h. Bonn. 1836.
hispidula Pursh. 2. D. olim hort.
incola.
iberica M. B. (*scordifolia* W.)
2. F. 1822.
intermedia Ait. (*sibirica* Link.)
♂. 2. D. 1826; etiam ex
hb. Paris. et Gryph. 1837.
lanata Jacqv. 2. D. 24-27 poll.
alt. fl. brunco-rubr. mens.
Jun. explic.; odor plantæ
ut in plerisq. teter.
palustris L. 2. D. sponte.
pubescens Tenor. (*St. annua*
var. Spr.) ex hort. Paris.
1837.
recta L. 2. D. flor. luteo-alb.
salviæfolia Tenor. ♂. ex hort.
Paris. 1837.
sylvatica L. 2. D. sporadic
in hort. 1836.

tomentosa L. 2. ex h. Gryph.
1836.

Stachytarpheta

angustifolia Vahl. 2. C. ex h.
Hafn. 1836.
prismatica Vahl. 2. C. 1810
et ex h. Hafn. 1837.

Stapelia

brevirostris H. Hamb. 2. C.
deflexa Jacqv. (*Gonostemma*
Haw.) 2. C.
divaricata Mass. 2. C.
diffusa. 2. C.
hamata Jacqv. 2. C. 1837.
hirsuta L.? (*ambigua* Mass.) 2.
C. flor. magn. atro-violac.
grandiflora Mass. 2. C.

Staphylea (Pimpernöt).

pinnata L. 2. D. 8 uln. alt.,
fl. alb. mens. Jun. expl.;
fructum quotannis in hor-
to maturat.
trifoliata L. 2. D. 6 uln. alt.;
fructum quam prior ra-
rius maturat, frigori ma-
gis impatiens.

Statice

Alliacea Cav. v. *Armeria*.
Armeria L. v. *Armeria maritima*.
Besseriana. ex h. Hafn. 1827
et Crac. 1837.
echioides L. ♂. F.
Gmelini W. 2. D. ex h. Hafn.
1827.
Limonium L. 2. D.
reticulata L. 2. D. F. 1½ uln.
alt. flor. parv., rubicund.
inodor. mens. Aug. explic.
scoparia Pall. 2. D. F. 1½ uln.
alt.; flor. Aug. Sept.
sinuata L. 2. F. ex h. Crac.
1837.
species. 2. T.

Stellaria

graminea L. 2. D. spont.
Holostea L. 2. D. *pedalis*, fl.
alb. Jun.

media With. (*Aisne* L.) ☉. D. spont.

Stenactis (*Aster*).

speciosa Lindl. 2. T. D. bipedalis et ultra flor. disci flav., radii lilacin. medio mens. Jun. explic. usque ad initium Novemb. dur.; intr. 1834 ex h. Londin.

Stenochilus

maculatus Ker. 2. T. ex hort. Kew. 1837. cur. Cl. J. Ag.

Sternbergia

lutea Ker. (*Amaryllis* L.) 2. F.

Stevia

alba 2. F. forte sequentis.
ovata Lag. 2. F. 1813. 1835.
1½ uln. alt. flor. alb.
pedata Cav. v. *Hymenopappus*.
serrata Cav. 2. F. 1806. — 1810. etc.

Stipa (Fjädergräs).

Aristella L. 2. F. ex h. Par. 1837.
capillata L. 2. D. 1810 et ex h. Paris. 1837.
gigantea Lag. 2. F. ex hort. Hafn. 1835.
pennata L. 2. D. pedalis. — intr. 1810.
seminuda Vahl. ☉. D. ex h. Hafn. 1835.
splendens Trin. (*membranacea* Pursh.) F. intr. 1835.
tortilis Desf. ☉. D. ex hort. Crac. 1837. forte eadem ac *St. seminuda* Danorum.

Streptocarpus

Rhexii Lindl. 2. C. T. Scapus spitham., flor. dilute cærul. striis violac.; in-odor.; mens. Aug. expl.; introd. 1834 ex h. Bonn.

Struthiopteris

europæa Sw. Fr. (*Struthiopteris germanica* W.) 2. D. in hort. nostro ad summum bipedalis; in insula vero Lybeck lacus Krageholmensis Scaniae spont. saltem 6-pedalem et abundantem vidi.

Stylidium

adnatum R. Br. 2. T. ex h. Kew. 1837.
fruticosum R. Br. 2. T. vix 4-pollicare, flor. parv., rubicund., inodor. versus finem mens. Jun. explic.; intr. 1833 ex h. Hamb.; in nostro primum floruit 1835.

Styphelia

ovalifolia Spr. 2. T. sub nomine *Acrotiche ovalifolia* R. Br. ex h. Kew. 1837. commun. cur. Cl. J. Ag.
Richei Labill. (*Leucopogon Richei* R. Br.) 2. T. ex h. Kew. 1837.

Succisa

alpina Spr. (*Scabiosa* L. *Cephalaria* Schrad.) 2. D. 3-pedalis et ultra; flor. albo-flavis.
centauroides Spr. (*Scabiosa corniculata* Kit.?) 2. D. ex h. Hafn. 1836. bipedalis, flor. albescenti flavis.
lævigata Spr. (*Scabiosa* Kit. *Scab. coriacea* W.) 2. D. 1810. 1826, et ex hort. Hafn. 1835.
pratensis Mönch. (*Scabios. Succisa* L.) 2. D.
syriaca Spr. (*Scabiosa* L.) ☉. D. 1810, et ex h. Paris. 1837.
tatarica Spr. (*Scabiosa* L. *Scabiosa elata* Horn.) ♂. D. sæpe 3 uln. et ultra alt., flor. magn. sulphureis.
transsylvanica Spr. (*Scabiosa*

L.) (3) D. ex hort. Paris.
1837.

Suocovia

balgarica Med. (*Bunius* L.) (3).

F. 1821; etiam ex h. h.
Hafn. et Paris. 1837.

Suninea (*Centaurea*?).

mollis. sub hoc nom. ex hort.
Paris. 1837.

Sutherlandia

frutescens R. Br. (*Colutea* L.)

♂. F. T. 1 uln. alt.; flor.
coccin.; ex h. Hamb. 1836.

Symphoria

glomerata Pursh. (*Symphoricar-*
pos vulgaris Horn. *Lonicera*
Symphoricarpus L.) ♂. D.
fere 2 uln. alt., florib. ru-
bicund. post leniores hie-
mes sæpe jam mens. Maj.
explicat. (1835 floruit d.
25 Febr. sub dio); olim
etiam in hort. culta et de-
nuo intr. 1834 ex hort.
Hamb.

racemosa Pursh. ♂. D. 2 uln.
alt., flor. parv. rubicund.
inod. Aug. explic.

Symphoricarpus

vulgaris Horn. v. *Symphoria glo-*
merata.

Symphytum

asperissimum M. B. 2. D. 2
uln. alt.; flor. azur. ante
explication. ruf., inodor.
Jun. — Aug. in Paradis-
lyckan copiose.

bullatum Jacq. (*tauricum* W.)
2. D. pedale et bipedal.
flor. albo-luteis, Maj. ex-
plicat.

officinale L. 2. D.
♂. flor. alb.

Syringa (Syrén).

chinensis L. ♂. D. 2½ uln. alt.
florib. lilacin. mens. Jun.
explic. in Plantat. Para-
dislyckan.

persica L. ♂. D. 2-3 uln. alt.;
florib. lilacin., odore mi-
tiori quam in *S. vulgaris*.
Arboret.

villosa Vahl. (*Ligustrum sinen-*
se Lour.) ♂. T.

vulgaris L. ♂. D. 4-8 uln. alt.
in Arboret.
♂. flor. alb.
γ. fl. lilacino.

Tagetes (Sammetsros).

erecta L. (3) D. sesquipedal.
flor. magnis, flav. sed co-
lor. et plenit. var.

Fevillæi sub hoc nom. ex h.
Kil. 1836.

lucida Cav. 2. F. ex h. Par.
1837.

patula L. (3) D. flor. color.
var. Jun. explic.

♂. flor. plen.
runcunculoides, ex hort. Crac.
1837.

tenuifolia Cav. (3) T. 1814.

Talinum

patens W. ♂. C. 1806. 1810.
horti incola, hodie exsul.

Tamarix

gallica L. ♂. F. 1 uln. alt.,
ægre coeli nostri hiemes
perfert.

germanica L. ♂. F. D., in h.
olim frequentius culta.

Tamus

Elephantopus Herit. v. *Dioscorea*.

Tanacetum (Rensan).

Balsamita L. v. *Balsamita vul-*
garis.

horeale, sub hoc nom. ex h.
Paris. 1837.

canariense Cand. ♂. ex hort.
Paris. 1837.

vulgare L. 2. D. 3-pedale;
flor. flav. fragrant., Aug.
explic.

♂. crispum. 3-pedale.

Taxus (Tax. Idgran).

baccata L. ♂. D. 15 uln. alt.

canadensis W. & D. ex hort.

Hamb. 1836.

hibernica & D. 13 uln. alt.;
intr. 1833 ex h. Hamb.

Tecoma

capensis v. *Bignonia*.

radicans Juss. (*Bignonia* L.)

& D. nostro coelo jam ad-
sucta, in Arboret. sub dio
cum successu colitur; intr.
1834 ex h. Hamb.

Telephium

Imperati L. & D. ex hort.

Hafn. 1827 et Par. 1837.

Tellima

grandiflora Dougl. v. *Mitella*.

Tetragonia

cristallina Herit. ☉. T. ex h.
Paris. 1837.

expansa Ait. ☉. T. D. 1810.
1820 et ex h. Par. 1837;
planta decumb. spitham.
floret. Aug.

Tetragonolobus

conjugatus Link. v. *Lotus*.

purpureus Mönch. v. *Lotus*.

siliquosus Roth. v. *Lotus*.

Teucrium

Arduini L. & D. F. ex h. Paris.
1837.

bracteatum Desf. & D. F. 1812.

Chamaedrys L. & D. F. flor.
rubr.

fruticans L. & D. F. 1820, herb.
nostr.

lucidum L. & D. F. ex h. Paris.
1837.

Marum L. (Kattgräs) & D. T. F.
florib. rubr.; tota planta
graveolens.

masiliense L. & D. F. 1810. 1812.

sibiricum L. v. *Nepeta*.

Thalictrum

appendiculatum H. Gryph. & D.
1837 ex h. Gryph. a

Cel. Hornsch. communic.

aquilegifolium L. & D. 1 uln.
alt.; in Fägelång spontan.

proterius; flor. ros. anther.
flav. Jul. Aug.

carolinianum Walt. & D. ex
h. Parib. 1837.

crenatum & D. ex h. Parib.
1837.

discolor Horn. & D.

elatum Jacq. & D.

exaltatum H. Gryph. & D. ex
h. Gryph. 1837.

flavum L. & D.

floribundum Schrö. & D. ex
h. Gryph. 1836.

foliosum Cand. & D. T. ex hort.
Chiswich. 1837.

glucum Desf. & D. ex hort.
Gryph. 1837.

medium Jacq. (*odnetum* W.)
& D. 2 uln. alt., flor.
parv. virescentib. anther.
flavis; Aug.

mitrocarpum. & D. ex h. Gryph.
1837.

minus L. & D. in hort. olim
cult.

nigricans Jacq. & D.

nutans Desf. & D. 1833, etiam
ex h. Gryph. 1837.

petaloideum L. & D.

purpurascens L. & D. 1827,
et ex h. Paris. 1837.

speciosum Mill. & D. ex h.
Hafn. 1827 et Gr. 1837.

specios.

Thapsia

garganica L. & D. F. ex hort.
Paris. 1837.

Thea (Thebuske),

Bohea L. (*chinensis* Sims, Spr.)
& D. T. 20 poll. alt.

viridis L. (*chinensis* var. Spr.)
& D. T. 1 uln. alt. intr. 1835
ex h. Hamb.

Thespesia

populnea Corr. Spr. v. *Hibiscus*.

Thlaspi

arvense L. ☉. D. spont.

Bursa Pastoris L. v. *Capsella*.
campestris L. v. *Lepidium*.
sativum Crantz. v. *Lepidium*.
saxatile L. v. *Aethionema*.

Thrinchia

hirta Roth. (*Leontodon* L. *Apar-*
gia Scop.) 2. D. 1807.

tuberosa Cand. (*Leontod.* L.
Apargia W.) 2. D. 1824.

Thryallis

brachystachya. 2. C. ex hort.
 Chiswich. 1837.

Thuia

occidentalis L. 2. D. ante 42
 annos in hortum introd.,
 jam 14 uln. alt. fl. mens.
 Mart.

orientalis L. 2. D. 6 uln. alt.,
 trunci diamet. 5-poll. Hie-
 mes nostras, etiam inten-
 siores perferre videtur, sed
 apices ramulorum quotan-
 nis frigore ferruginantur;
 flor. mens. Mart.

sphaeroides Rich. (*Cupressus thy-*
oides L.) 2. F. intr. 1836
 ex h. Hamb.

Thunbergia

capensis Retz. 2. C.

Thymus (Timjan).

alpinus L. 2. D. 1821.

filiformis Ait. 2. 2. T. ex h.
 Paris. 1837

inodorus Desf. ex hort. Crac.
 1836.

patavinus Ait. 2. 2. D. ex hh.
 Hafn. 1836, Paris. 1837.
 flor. rubr.

Teneriffæ Pers. (*terebinthaceus*
 W.) 2. 2. T.

vulgaris L. 2. D.

Tiaridium

indicum Lehm. (*Heliotropium*
 L.) 2. C.

Tigridia

Pavonia Juss. (*Ferraria Pavo-*
nia L.) 2. T. flor. specios.

Tilia (Lind).

alba Horn. (*argentea* Cand.) 2.
 D. 25 uln. alt., trunci
 diametro prope radicem
 20-poll.

canadensis Mx. (*americana* L.
glabra Vent.) 2. D. 9 uln.
 alt., intr. 1826. Folia
 maxima.

europæa L. 2. D. 28 uln. alt.
 Paradislyckan.

2. 2. asplenifolia; 6 uln. alt.
 in Arboret.

grandiflora Ehrh. 2. D.

laxiflora Mx. 2. D. valde affi-
 nis priori.

Tillandsia

amoena Link. 2. C. floruit in
 Caldar. 1838 a d. 15 Jan.
 ad 6 Febr.

Tolpis

barbata Gaertn. (*Crepis* L. *Dre-*
punia Desf. Cand.) 2. D.
 8 poll. alt. flor. flav. Aug.
coronopifolia Spr. (*Crepis* Desf.)
 2. D. ex h. Kil. 1836.

Tordylium

Anthriscus L. v. *Torilis*.

Torilis

Anthriscus Gmel. (*Tordylium*
 L. *Caucalis* W.) 2. 2. D.
 3-pedal., flor. roseo-albis.
helvetica Gmel. (*Caucalis arven-*
sis Huds. *Scandix infesta*
 L.) 2. D. ex h. Hafniæ
 1837. sesquipedal., flor.
 flavescenti albis.

Tournefortia

fruticosa R. Br. (*Messerschmidia*
 L.) 2. T. 5-pedalis; flor.
 parv. virescenti albis, in-
 odor. d. 10 Aug. explic.
heliotropioides Hook. 2. T. ex
 h. Paris. 1837.

Trachelium

cæruleum L. 2. F. 1810. 1821,
 etiam ex h. Crac. 1836 ses-
 quipedal., fl. cærulescent.

Tradescantia

erecta Cav. 2. T. D. 6-10 poll.
 alt., flor. cærul. violaceo
 antheris flav. mens. Jun.
 explic.

Sellowii (*Sellowiana* Pohl.) 2.
 T. F. sub hoc nom. ex h.
 Crac. 1836.

subaspera Ker. ex hort. Paris.
 1837.

virginica L. 2. D. pedal. flor. cyan.
species.

Tragopogon

(Bockskägg. Salsofi).

coloratus 2. ♂. D. ex hort.

Gryph. 1837.

crocifolius L. ♂. D. sesquipedal., flosculis subviolaceis.

dubius Horn. Spr. ♂. D. sesquipedal. floscul. ochroleucis.

major Jacq. ♂. D. 1833. 1836. floscul. flav.

orientalis L. ♂. D. 1833 et ex h. Gryph. 1836 flosculis aureis.

picroides L. v. *Urospermum*.

porrifolius L. ♂. D. floscul. rubr. l. violaceis.

pratensis L. ♂. D. spont. sesquipedal. flor. flav.

roseus Trev. ♂. D. floscul. roseis, ex h. Gryph. 1836.

Tragopyrum

lanceolatum M. B. (*Polygonum frutescens* L.) 2. T. F. D.

Trevirania

coccinea W. Spr. (*Cyrilla pulchella* Herit.) 2. C. spitham., flor. coccineis.

Trianthema

monogynum L. 2. C. ex hort. Paris. 1837. flor. parvis.

Trichocladus

crinitus Pers. (*Dahlia crinita* Thunb.) 2. T. ex h. Kew. 1837.

Trichodium (*Agrostis*).

Trichosanthes

palmata H. Hamb. 2. C. intr. 1834.

Tridax

procumbens L. v. *Balbisia elongata*.

Trientalis

Trifolium

(Klöver. Våpling).

albidum Retz. 2. D. ex hort. Paris. 1837.

alexandrinum L. 2. D. 1824 et ex h. Paris. 1837.

alpestre L. 2. D. flor. rubr.

clypeatum L. 2. D. 1810, et ex h. Paris. 1836.

elegans Sav. 2. (2.) D. 1836.

erinaceum M. B. D. ex hort. Paris. 1837.

europæum. ♂. D. ex h. Paris. 1836, seminib. educat.

filiforme L. 2. D. spont. flor. citrin. Jun.

flexuosum Jacq. (*medium* L.) 2. D. nuper etiam ex h.

Paris. 1837. sesquipedal. flor. purp. Jun.

fragiferum L. 2. D. (etiam circa Lund ubique spont.) spitham. fl. rubicund. parum odor. Jul. — Sept.

hybridum L. 2. D. sesquiped. flor. pallid. ros. Jun.

incarnatum L. 2. D. 1812--1834. iterum ex h. Gryph. 1837. flor. incarnat.

intermedium Guss. 2. D. ex h. Paris. 1837.

lappaceum L. 2. D. 1810. 1815, et ex hort. Paris. 1837.

libanoticum H. Lips. D. 1835; 18-22 poll. alt. flor. rubr. inodor. Jul. explic.

Lupinaster L. (*Pentaphyllum Lupinaster* Pers.) 2. D. 1808. 1810.

maritimum Sm. 2. ♂. D. ex h. Paris. et Gryph. 1837.

Melilotus etc. v. *Meliloti* species.

Michelianum Sav. 2. D. ex h. Paris. 1837.

montanum L. 2. D. pedal. fl. alb. Jun.

ochroleucum L. 2. D. 1822, et nova vice ex h. Paris. 1837, flor. luteis.

pallidum Kit. 2. D. 1812 et denuo ex h. Paris. 1837.

pannonicum L. 2. D. 1813.

pratense L. ④. D. spont.
repens L. ④. D. spont.
resupinatum L. ④. D. 1810.
 1812, iterumque ex hort.
 Paris. 1837. flor. purpur.
scahrum L. ④. D. ex h. Pa-
 ris. 1836. flor. parv. albe-
 scentibus.
speciosum W. D. ex h. Paris.
 1836.
squarrosus L. ④. D. 1810.
 1815 et ex h. Paris. 1837.
striatum L. ④. D. flor. parv.
 incarnat.
stellatum L. ④. D. 1815 et ex
 h. Paris. 1837.
suaveolens W. 1821.
tomentosum L. ④. D. ex h.
 Paris. 1837.

Trigonella

arcuata Mey. ④. D. ex h. Kil.
 1836.
azurca. sub hoc nom. ex hort.
 Petrop. 1836.
Calliceras Fisch. ④. D. 1835.
 1836. pedalis et ultra, fl.
 flav. inodor. Jul.
cancellata Fisch. ④. D. ex h.
 Paris. 1837.
corniculata W. ④. D. 1812.
 1815 etc.
elegans. ④. D. ex hort. Paris.
 1837.
Fischeriana. ④. D. ex hort.
 Paris. 1836.
Foenum græcum L. ④. D. pe-
 dalis et ultra; florib. alb,
 Jul. explic.
gladiata Stev. ex h. Petropol.
 1836.
hamosa L. ④. D. ex h. Paris.
 1837.
hybrida Pourr. ④. (④.) D. F.
 ex h. Paris. 1837.
monantha. sub hoc nom. ex h.
 Petrop. 1836.
monspeliaca L. ④. D. 1814
 et ex h. Paris. 1837.

polycerata L. ④. D. 1824 et
 ex h. Paris. 1837.
prostrata Cand. ④. D. ex h.
 Paris. 1837.
spinosa L. ④. D. ex h. Kil.
 1835.
striata L. suppl. (tenuis Fisch.)
 ④. D. 1835. pedalis, flor.
 parv. flav. inodor. Jul.
uralensis Ledeb. ④. ex hort.
 Kil. 1836.
species.

Tripsacum

dactyloides L. ④. (④.) D. F.
 ex h. Paris. 1837.
hermaphroditum L. (*Cenechrus le-
 vigatus* Trin.) ④. C. 1806-
 1821 etc.

Triticum (Hvete).

caninum Schreb. ④. D.
dasyanthum Led. ④. D.
densiflorum W. ④. D.
junceum L. (var. *gigantum*
 Retz.) ④. D.
monococcon L. ④. D. 1812.
 herbar. nostr.
repens L. (Qvickrot) ④. D. sp.;
 vitium.
Spelta L. ④. D. 1812. 1815. herb.
Zea. ④. D. ex h. Paris. 1837.

Tritonia (Waitzia Wilbr.)

refracta Ker. v. *Gladiolus*.
squalida Ker. Spr. (*Ixia* Ait.
Ixia crocata Ker.) ④. T.
 ex h. Paris. 1837.

Trollius

asiaticus L. ④. F. sesquipedal.
 flor. flav. extus virescent.
 Jun.
europæus L. (Bullerblomster)
 ④. D.

Tropæolum

(Indiansk Krasse).
majus L. ④. D. flor. crocei;
 Jul. — Novembr.

β. atro-sanguineum ex h.
Hamb. 1834.
minus L. ☉. D. flor. auran-
tiaco-flav. Aug.

horto copiose crevit, dein-
de extincta, nuper iterum
introducta; verno tempo-
re florens.

Tulipa (Tulpan).

Celsiana Red. 2. F. olim horti
incola, hodie profuga.
Gesneriana L. 2. D. plures va-
rietates colore et plenitu-
dine adsunt.
sylvestris L. 2. D. 10-15 pol-
licar. flor. flav. subnutan-
tes, mens. Maj. explic.
In graminos. horti no-
stri ut plurimum Lunden-
sium copiose spontanea,
plerumque sterilis.

Turritis

glabra L. (*Arabis lucida* Spr.)
♂. D. ex h. Crac. 1836.
3-pedal. flor. parv. albo-
flav. Jul.
hirsuta L. (*Arabis* Scop.) ♂.
D. sesquipedal. flor. alb.
Jun.

Tussilago

alba L. 2. D. 6-pollicar. flor.
alb., Mart. — Maj.
Anandria L. - (*Chaptalia* Spr.
Perdicium B. Br.) 2. D.
1822. et ex h. Hafn. 1837.
Farfara L. (Hästhof) 2. D.
fragrans Vill. 2. D. - Spi-
tham., fl. albo-rubicund.
fragrant. mens. April. (in-
terd. Dec. aut Jan.) expl.

Olim in Tepidar. ser-
vata, jam sub Dio in hor-
to culta bene viget et quo-
tannis abund. floret; etiam
in Paradislycan.

nivea Vill. (*paradoxa* Retz.) 1806
in hort. culta.

Petasites L. 2. D. in hortis
Lundens. vitium.

spuria Retz. 2. D. ante plu-
res annos fere neglecta in

Ulex

europæus L. ♀. F. D. 1½ uln.
alt. in plantatione Para-
dislycan.

strictus ♀. F. 1 uln. alt.; in-
trod. 1834 ex h. Hamb.

Ulmus (Ahn).

americana L. ♀. D. circa ann.
1798 in horto nostr. cul-
ta, hodie extincta.

betuloides ♀. D. 4 uln. alt. in
Arboret.

campestris L. ♀. D.

β. montana Sm. 3 uln. alt.
in hort.

corylifolia ♀. D. in Arboret.

effusa W. ♀. D. olim in hort.
cult, ibi diutius non ex-
stat.

gigantea ♀. D. in nov. Arbor.
tantum 5 uln. alt.

pumila W. ♀. D. 13 uln. alt.,
trunci diamet. mox supra
terram 12-poll., flor. Apr.
Maj.

suberosa Mönch. ♀. D.

Usula ♀. D. Arbor 28 uln.
alt. trunci diametro cir-
citer 1½ uln. sub hoc no-
mine in horto exstat, flo-
res non proferens.

Uropetalum

serotinum Ker. (*Hyacinthus* L.)
2. T. 1826, et ex h. Kil.
1836.

Urospermum

picroides Desf. (*Tragopogon* L.
Arnopogon W.) ☉. D.
1811, etiam ex h. Paris.
1837; flor. flav. involucr.
viridi.

Urtica (Nässla).

canadensis L. 2. D. 1½ uln.
alt., flor. albesc. Jun., in-
nocua, nec urens

cannabina L. 2. D. 1808-1820.
valde urens.

dioica L. 2. D. spont.

Dodartii L. ☉. D. 1808 etc.

pilulifera L. (*bularica* L.) ☉.

D. valde urens.

urens. L. ☉. D. spont.

Usteria

Barclayana v. *Maurundia*.

scandens Cav. v. *Maurundia sem-
perflorens*.

Uvularia

perfoliata L. 2. D.

Valantia

Aparine Horn. ☉. D. 1836.
sesquipedal. flor. alb.

articulata L. (*Galium* R. et Sch.)

☉. F. 1812.

cruciata L. v. *Galium*.

Cucullaria L. (*Galium* Spr.) ☉.

D. sub hoc nom. in hort.
cult. 1810.

spuria Horn. ☉. D. ex hort.
Hafn. 1837.

Valeriana

angustifolia Cav. v. *Centranthus
ruber* β.

Caleitrapa L. v. *Centranthus*.

corniculata. ☉. D. ex h. Par.
1837.

officinalis L. (Wenderot) 2.

D. 3-pedal. fl. albo-rubic.

Phu L. 2. D. 2 uln. alt., fl.
alb., mens. Jun. explic.

pyrenaica L. 2. D. ex h. Hafn.
1837.

rubra L. v. *Centranthus*.

Velezia

rigida L. ☉. F. ex h. Paris.
1837.

Vella

annua L. ☉. D. circiter 8-pol-
licaris, flor. pallide flav.

Veratrum (Prustrot).

album L. 2. D. 1810 — hodie
abest.

nigrum L. 2. D. 3-pedale et
ultra; flor. Aug.

Verbascum (Kongsljus).

austriacum Schrad. (*orientale*
M. B.) ♂. D. ex h. Hafn.
1836.

blattarioides Lam. (*virgatum*
Spr.) ♂. D. ex h. Hafn.
1837.

compactum M. B. ♂. D. ex h.
Petr. 1836.

condensatum Schrad. ♂. ex h.
Hafn. 1836. flor. flav.

glabrum W. β. F. ex h. Hafn.
1836.

gnaphaloides M. B. ♂. D. ex
h. Paris. 1837.

Lychnitis L. 2. D. 1821 herb.
nostr. flor. albesc.

nigrum L. 2. D. 3-pedal. et
ultra. Aug.

ovatum Schrad. D. ex h. Par.
1837.

phoeniceum L. ♂. D. 1822. herb.
nostr. flor. cyan.

pyramidatum M. B. ♂. D. ex
h. Paris. 1837.

repandum W. ♂. D. ex hort.
Hafn. 1837.

rugulosum W. (*australe* Schrad.).
♂. 2. D. 3½ uln. alt., flo-
re magn. toto flav. stamin.

3 hirsut. 2 glabr.; Aug.

speciosum Schrd. ♂. D. ex h.
Hafn. 1837.

Verbena (Jernört).

angustifolia Mx. 2 ex h. Pa-
ris. 1837.

Aubletia Jacqv. ☉. ♂. F. 1810
et ex h. Crac. 1836. flor.
rubr.

Valerianella v. Fodia.

- banariensis* L. 2. T. F. flor. caerulea.
bracteosa Mx. ①. 2. F. D. 1810 et ex h. Par. 1837.
caroliniana L. 2. F. ex hort. Lund. herbar. nostr.
gacilis sub hoc nom. ex h. Paris. 1837.
hispida R. et P. ①. T. ex h. Paris. 1837.
littoralis Humb. T. ex h. Par. 1837.
multifida R. et P. 2. C. ex h. Paris. 1837.
nodiflora L. 2. C. ex h. Lund. herb. nostr.
officinalis L. 2. D. 28 poll. alt., flor. parv. rubic. l. lilacin. mens. Aug. expl.
stricta Vent. (*rigens* Mx.) 2. T. ex h. Paris. 1837.
triphylla Herit. v. *Lippia*.
urticaefolia L. 2. D. herbar. nostr.
- caucasica* M. B. 2. D. ex h. Paris. 1837.
ceratocarpa 2. D. ex hb. Hafn. et Paris. 1837.
Chamaedrys L. 2. D. spont.
crenulata Hoffm. (*villosa* Spr.) 2. D. 1 uln. alt. florib. late caerulea. Jun. explic.
decussata Ait. 2. F.
didyma Tenor. ①. D. plant. pedalis procumb. flor. caerulea. alb. Jul. Aug.
elatio Ehrh. 2. D. nimis affinis. *V. longifolia*; 2½ uln. alt., florib. caerulea. mens. Sept. explic.
fruticulosa L. 2. F. 1819 et ex h. Gryph. 1836.
Gentianoides Vahl. 2. D. sesquipedal., flor. albo-caerulea. Jun. explic.
glabra Ehrh. 2. D. ex hort. Gryph. 1836 flor. caerulea. 2. fl. alb. 1819.
grandis Fisch. 2. D. 1835 et ex h. Gryph. 1837, 2½ uln. alt., florib. caerulea. Aug. explic.
hederifolia L. ①. D. spont. copios. 4-poll. flor. Mart. — Octobr.
incana Luxm. 2. D. ex hort. Lund. 1833 herbar. nostr.
laciniata 2. D. 1833 et ex h. Hafn. 1837.
latifolia L. 2. D. 26 pollic. alt., fl. late caerulea. mens. Jun. explic.
longifolia L. 2. D. 24—26 poll. alt. flor. cyan. versus fin. Jun. explic.
maritima L. 2. D. flor. caerulea. 2. fl. alb.
media Schrad. 2. D. 1833 ex h. Hafn. 1837. flor. caerulea. 2. latifolia ex h. Gryph. 1836.
multifida 2. D. ex h. Paris. 1837.
officinalis L. (Erenpris) 2. D. flor. caerulea.
orientalis Ait. 2. D. 22 poll. alt., flor. late caerulea. versus finem. Jun. explic.

- paniculata* L. 2. D. 24 poll.
 alt., flor. late carul. Jun.
pinnata L. 2. D. ex h. Bonn.
 1836. flor. carul.
prostrata L. 2. D. 16-18 poll.
 alt., flor. carul. Jun.
romana. ☉. D. ex h. Paris.
 1837.
serpyllifolia L. 2. D. spont.
sibirica L. 2. D. 2 uln. et
 ultr. alt. flor. dilute vio-
 laceo-rubr. inodor. mens.
 Jun. explic.
taurica W. 2. D. (var. *orien-
 talis* secund. Spr.) sesqui-
 pedal. flor. late carul. Jun.
Teucrium L. (*dentata* Schmidt)
 2. D. 13-14 pollic. alt.
 flor. late carul. Jun.
urticaefolia L. 2. D. 1819. her-
 bar. nostr.
verna L. ☉. D. spont.
virginica L. 2. D. 1½-2 uln.
 alt., flor. rubicundo-alb.
 Sept.
speciosa ex h. Paris.

Vesicaria (*Alyssum* Linn.)

- cretica* Poir. 2. F. ex h. Hafn.
 1827 et Crac. 1836.
sinuata Poir. 2. D. 1822
 et ex h. Paris. 1837. flor.
 flav.
utriculata Lam. 2. F. florib.
 pallide flav.

Viburnum

- dentatum* L. 2. D.
edule Pursh. 2. D.
Lantana L. 2. D. 3-5 uln.
 alt.; flor. alb. odore nau-
 seoso, mens. Jun. explic.
 β. *minor*
Lentago L. 2. D. 1 uln. alt.;
 in nov. Arbor. adhuc non
 floruit.
Opulus L. 2. D. 2 uln. alt.
 flor. simpl. albo-virescent.
 β. *roseum* (Südbollar) 4-5

- uln. alt. florib. alb.; in
 horto et in Paradislykan.
prunifolium L. 2. D. vix 3
 uln. alt., flor. parv. alb.
 inodor. mens. Jun. explic.
 introd. in nov. Arboret.
 1837.

- Tinus* L. 2. T. F. 1 uln. et
 ultra alt., flor. alb. l. ru-
 bicundo-albis, Januar. —
 Mart.

Vicia (Vicker).

- amoena* Fisch. ☉. D. ex hort.
 Crac. 1836.
angustifolia Roth. Fr. ☉. D.
 flor. purpur.
atropurpurea Desf. ☉. D. ex
 h. Hafn. 1827 et Paris.
 1837.
Bacsa Münch. ☉. D. ex hort.
 Kil. 1835.
benghalensis L. ☉. D. ex h.
 Crac. 1836.
bithynica L. ☉. D. 1810 et
 ex h. Paris. 1837.
dumetorum L. 2. D. flor. pur-
 pur. Jun. explicatis, in
 Paradislykan.

Ervilia W. v. *Ervum*.

Faba L. ☉. D.

- β. *seminib.* stramineis. 4-
 pedal. flor. odor. grat.
 γ. *seminib.* atro-rubentib.
 bipedal. flor. Jul.
globosa Retz. ☉. D. in hort.
 cult. 1812-1815. 1834.
gracilis Loisel. ☉. D. ex hort.
 Paris. 1837.
graminea Sm. ☉. ex h. Crac.
 1836.
helvetica. sub hoc nom. ex h.
 Crac. 1836.
hybrida L. ☉. D. 1813. 1815
 et iterum ex h. Par. 1837.
 flor. flavescentib.
linifolia, sub hoc nom. ex h.
 Crac. 1836.
lutea L. ☉. D. 8 pollic., flor.
 albo-lut., inodor. mens.
 Jul. explic. 1837; introd.
 1810.
monantha Retz. (*calcarata* Desf.)
 ☉. D. 1806-1820; etiam
 ex h. Crac. 1836.

- marbonensis* L. ☉. D. 1835.
polyphylla Desf. ☉. D. ex h.
 Kilens. 1836. Similis *V.*
Cracca.
purpurea Stev. ☉. D. ex h.
 Crac. 1836.
sativa L. ☉. D.
segetalis Thuill. ex h. Crac.
 1836.
sepium L. 2. D. sesquipedal.
 flor. rubro-violac. Jun.
serratifolia Jacq. ☉. D. in h.
 Lund. 1810. 1814.
sordida Kit. ☉. D. 1815 et ex
 h. Paris. 1837. flor. sor-
 dide flav.
 β. flor. flav.
sylvatica L. 2. D.
villosa Roth. ☉. D. Simil. *V.*
Cracca; floruit in horto
 alb. Aug. usque ad Octob.
 1837.
 species.

Vieusseuxia

pavonia Cand. Spr. v. *Iris*.

Vinca (Vintergrön).

herbacea Kit. 2. D. fl. caerulea.
major L. β. D. fl. magna, cae-
 rulea in medio mens. Maj.
 explic. — Sub tegmine
 muri hibernae etiam inten-
 siores sine discrimine su-
 stulit.

minor L. β. D. flor. caerulea in
 med. mens. April. explic.
 β. fol. variegat.

γ. flor. plen.

rosea L. β. C. sesquiped. flor.
 roseis.

β. fl. alb. Maj.

Viola

altaica Pall. 2. fl. albo; ex h.
 Gryph. 1836.

appendiculata Cand. ☉. ex h.
 Gryph. 1836.

canina L. 2. D.

hederacea Labill. ex h. Paris.
 1837.

hirta L. 2. D. spont. inodor.

lanceifolia Bess. 2. ex h. Hafn.
 1837.

mirabilis L. 2. D. in Para-
 dislyckan.

odorata L. (Luck-Fioler) 2. D.
 flor. cyan. April. Maj.

β. flor. albid. plen.

tricolor L. (Styfmors-Fioler)

☉. ♂. D. sponte.

β. *arvensis* Wahl. spont.

γ. *rothomagensis* Pers.

δ. flor. purpureo-atro.

species.

Viscago

molissima Horn. v. *Silene*.

Vitex

Agnus castus L. β. F. D. Usque
 ad 1836 in Tepidar. no-
 stro servata, deinde in h.
 sub Dio culta bene viguit.

Vitis (Vinstock. Vinranka).

hederacea Ehrh. (*Ampelopsis* Mx.

Hedera 5-folia L.) β. D.

folia autumnale coerulea.

Labrusca L. β. olim in horto
 sub Dio culta.

palmata Vahl. β. D.

incisa Jacq. β. F. D.

vinifera L. β. D. floret Jun.

vulpina L. β. D. in hort. olim
 culta.

Vogelia paniculata Horn. v. Nestia.

Wahlenbergia

(*Campanula*).

elongata Schrad. ☉. T. ex h.
 Kil. 1836.

gracilis Cand. fl. ☉. T. ex
 hh. Bonn. 1836 et Paris.
 1837.

pendula Schrad. ☉. T. ex h.
 Kil. 1836, sesquipedal.,
 flor. parv. pendul.

Waldsteinia

geoides W. 2. D. spithamea,

caespitos., flor. flav. jam
d. 26 April. explicat. intr.
1824 ex h. Hamb.

aliquot decennia in hort.
cult., jam extincta.

Wedelia

perfoliata W. v. *Melampodium*.

Xanthium

orientale L. fl. ☉. D. ex h.
Hafn. 1827 et Par. 1837.
strumarium L. ☉. D. 3-peda-
le; flor. glaucescenti viri-
dih., mens. Aug. explic.

Xanthorrhiza

apiifolia Herit. ♀. Olim in h.
sub Dio plantata, sed sine
successu. Cfr. Retz. in Act.
Holm. 1798. p. 78.

Xanthoxylon

fraxineum W. ♀. D. fere bi-
pedale, in novo Arboret.

Xeranthemum

(*Pappersblomma* Hortulan).

annuum L. ☉. D. bipedale et
ultra; floret Aug. — Nov.
♂. flor. alb.

fulgidum L. v. *Elichrysum*.

inapertum Horn. ☉. D. 26
poll. alt., flor. subulacinis,
Aug. explicat.

orientale W. (*Chardinia xeran-
themoides* Desf.) ☉. ♀. F.
intr. 1834 ex h. Bonn.;
deinde ex h. Petrop. 1838.
etiam miss.

Xerotes

longifolia R. Br. v. *Lamandra*.

Ximenesia

encelioides Cav. ♀. ☉. F. D.
20 pollic. alt.; flor. flav.
Aug. explic.

Xylophylla

Arbuscula W. (*Phyllanthus spe-
ciosus* Jacqv.) ♀. C. ante

Yucca

Gloriosa L. ♀. T. F.
filamentosa L. ♀. ♀. T.

Zacintha

verrucosa Gärtn. ☉. 1811. 1815
et ex h. Kil. 1836; peda-
lis, flor. subflavescent.

Zanthorrhiza v. Xan- thorrhiza.

Zanthoxylon. v. Xantho- xylon.

Zea (Msis).

Curagua Molin. ☉. ex h. Kil.
1836.

Mays L. (Turkiskt-Hvete) ☉.
D. 2 uln. alt., flor. Aug.;
plures varietates in horto
coluntur.

Zephyranthes

grandiflora Spr. v. *Amaryllis*
minuta.

Zingiber (Ingefära).

officinale Rosch. (*Amomum Zin-
giber* L.) ♀. C. bipedale
et ultra.

Zinnia

ambigua Link. ☉. D. ex hort.
Kil. 1836.

aurea Booth. ☉. D. ex hort.
Hamb. 1834 et Crac. 1836;
16 poll. alt., florib. flav.
Aug.

elegans Jacqv. ☉. T. D. 1 uln.
alt., flor. specios. purpur.,
carneis, coccin., Jun. —
Oct.; 1810 et iterum intr.
1833 ex h. Hamb.

grandiflora. ☉. D. ex h. Crac.
1837.

multiflora L. ☉. D. 1811.
1834 et ex h. Kil. 1836.
1½ uln. alt., flor. coccin.
aut. flav. Aug. — Octob.

pauciflora L. ☉. D. 1808.
tenuiflora Jacqv. (*revoluta* Cav.
Booth) ☉. D. 1½ uln. alt.;
flor. coccin., Aug. — Oct.

verticillata Andr. ☉. D. 1806;
quæ sub hoc nom. ex hor-
tis aliquoties communicata
ad alias species referenda.

Ziziphora

capitata L. ☉. D. sub spi-
tham., flor. lilacin.

hispanica L. ☉. D. 1833 et
ex h. Hafn. 1835.

tenuior L. ☉. D. 1810 et ex
hh. Petropol. 1834. Hafn.
et Crac. 1836. flor. albis.
species.

Zizyphus

Paliurus W. v. *Paliurus au-
stralis*.

Zoëgea

Leptaurea L. ☉. T. fere bi-
pedalis; flor. aureis; ex h.
Paris. 1837.

Zygophyllum

species.

Corrigenda.

Pag. 1. l. 7 (ad dextram). 1836. l. 1826.

„ 9. l. 21. Antherium. l. Anthericum.

„ 12. l. 8. spont. delect.

„ 36. l. 3 (ab infra). Dorychnium. l. Dorycnium.

„ 52. l. 3 (ad dextram). Illecebrum. l. Illecebrum.

„ 81. l. 23 (ad dextram) pendunculata l. pedunculata.
leviores mend. typ. Ipse corrig. b. Lector.

the 1990s, the number of people in the world who are undernourished has declined from 1.1 billion to 800 million. The number of people who are malnourished has declined from 1.5 billion to 1 billion. The number of people who are obese has increased from 100 million to 300 million. The number of people who are overweight has increased from 100 million to 300 million. The number of people who are obese and overweight has increased from 100 million to 300 million. The number of people who are obese and overweight has increased from 100 million to 300 million.

1. *Principles of Mathematics* by David Hilbert. This book is a classic work in the foundations of mathematics, discussing the principles of arithmetic and geometry.

• • • • •





